

COMP 2211 Exploring Artificial Intelligence
K-Nearest Neighbor - Population and Sample Standard Deviation

Huiru Xiao

Materials Provided by Dr. Desmond Tsoi
Department of Computer Science & Engineering
The Hong Kong University of Science and Technology, Hong Kong SAR, China

Standard Deviation

- Standard deviation is one of the most common ways to measure the spread of values in a dataset.
- There are two different types of standard deviation you can calculate, depending on the type of data you are working with.
  - Population standard deviation
  - Sample standard deviation

Population Standard Deviation

- We calculate the population standard deviation when the dataset we are working with represents the entire population, i.e. every value that we are interested in.
- The formula to calculate a population standard deviation, denoted as σ, is:

$$\sigma = \sqrt{\frac{\sum_{i=1}^{n}(x_i - \mu)^2}{n}}$$

where

- $\sum$: Summation
- x_i: The ith value in the dataset
- μ: The population mean
- n: The population size

Sample Standard Deviation

- We calculate the sample standard deviation when the dataset we are working with represents a sample taken from a large population of interest.
- The formula to calculate a sample standard deviation, denoted as s, is:

$$s = \sqrt{\frac{\sum_{i=1}^{n}(x_i - \overline{x})^2}{n-1}}$$

where

- $\sum$: Summation
- x_i: The ith value in the dataset
- $\overline{x}$: The sample mean
- n: The sample size

Population vs. Sample Standard Deviation

- The difference between the population and the sample standard deviation: When calculating the sample standard deviation, we divided by n-1 instead of n.
- Because when we calculate the sample standard deviation, we tend to underestimate the true variability in the population. In other words, our estimate of the true population standard is biased.
- To correct this bias, we divide by n-1. This makes the sample standard deviation an unbiased estimate of the population standard deviation.

That's all!

Any questions?