

Set Theory

Discrete Mathematic

Anders Kalhauge



Fall 2017



Set primitives

- Notation

- Ordered pairs and Cartesian product

- Relations and Functions

Venn diagrams and set operations

Subsets

- Set equality

- Empty set and Powersets

- Equality laws for sets

- Set properties

Set primitives

Notation

Ordered pairs and Cartesian product

Relations and Functions

Venn diagrams and set operations

Subsets

Set equality

Empty set and Powersets

Equality laws for sets

Set properties

The sets of ...

- \mathbb{R} all real numbers
- \mathbb{R}^+ all positive (not 0) real numbers
- \mathbb{R}^- all negative (not 0) real numbers
- \mathbb{R}^{nonneg} all nonnegative (positive or 0) real numbers
- \mathbb{Z} all integers
- \mathbb{N} or \mathbb{Z}^+ all positive integers
- \mathbb{N}_0 or \mathbb{Z}^{nonneg} all nonnegative integers
- \mathbb{Q} all rational numbers (quotients)

Set-roster notation

$$\begin{aligned} A &= \{1, 2, 3\} \\ &= \{3, 1, 2\} \\ &= \{1, 2, 2, 3, 3, 3\} \end{aligned}$$

$$B = \{10, 11, 12, \dots, 119\}$$

Set-builder notation

$$M = \{x \in S \mid P(x)\}$$

$$A = \{x \in \mathbb{Z} \mid -2 < x < 7\}$$

$$B = \{x \in \mathbb{N} \mid x^2 < 10\}$$

$$C = \{x \in \mathbb{R}^+ \mid \log(x) < 10\}$$

What are the members of:

$$\{x \in \mathbb{N} \mid x^2 + 20 < 100\}$$

defined as sets

The ordered pair

$$(a, b) \neq (b, a) \neq (a, b, c)$$

defined as sets

The ordered pair

$$(a, b) \neq (b, a) \neq (a, b, c)$$

can be defined as

$$\{\{a\}, \{a, b\}\}$$

defined as sets

The ordered pair

$$(a, b) \neq (b, a) \neq (a, b, c)$$

can be defined as

$$\{\{a\}, \{a, b\}\}$$

which is not the same as

$$\{\{b\}, \{b, a\}\}$$

The pairs

$$(a, b)$$

and

$$(c, d)$$

are equal if, and only if

$$a = c \wedge b = d$$

Cartesian product of A and B is $A \times B$ read “ A cross B ”

$$A \times B = \{(a, b) | a \in A \wedge b \in B\}$$

Cartesian product of A and B is $A \times B$ read “ A cross B ”

$$A \times B = \{(a, b) | a \in A \wedge b \in B\}$$

Example:

$$\begin{aligned} &\{1, 2, 3\} \times \{3.4, 7.2\} = \\ &\{(1, 3.4), (1, 7.2), (2, 3.4), (2, 7.2), (3, 3.4), (3, 7.2)\} \end{aligned}$$

Cartesian product of A and B and C is $A \times B \times C$

$$(A \times B) \times C = \{((a, b), c) | (a, b) \in A \times B \wedge c \in C\}$$

$$(A \times B) \times C = \{((a, b), c) | (a \in A \wedge b \in B) \wedge c \in C\}$$

$$A \times B \times C = \{((a, b), c) | a \in A \wedge b \in B \wedge c \in C\}$$

Short hand for the pair-pair $((a, b), c)$ is called a tuple:

$$(a, b, c)$$

A relation R between two sets A and B can be defined as a subset of the cartesian product.

$$R \subseteq A \times B$$

- A is the **domain** of R
- B is the **co-domain** of R

$$x R y$$

tells that

$$(x, y) \in R$$

Given a relation *Persons*, and the sets of *IDs*, *FirstNames*, *LastNames*

$$Person \subseteq IDs \times FirstNames \times LastNames$$

Where the subset is legal combinations of the three, normally defined by the tuples in a relational database.

- The function F from set A to set B
- is a relation with
 - domain A
 - and co-domain B
- $\forall x \in A, \exists y \in B, (x, y) \in F$
- $\forall x \in A \wedge y \in B \wedge z \in B, (x, y) \in F \wedge (x, z) \in F \rightarrow y = z$

Set primitives

Notation

Ordered pairs and Cartesian product

Relations and Functions

Venn diagrams and set operations


Subsets


Set equality


Empty set and Powersets


Equality laws for sets


Set properties











Notation

Ordered pairs and Cartesian product

Relations and Functions

Venn diagrams and set operations

Subsets

Set equality

Empty set and Powersets

Equality laws for sets

Set properties

$$A \subseteq B$$

$$A \subseteq B$$



$$\forall x, x \in A, x \in B$$

$$A \not\subseteq B$$



What should go here?

$$A \not\subseteq B$$



$$\exists x, x \in A, x \notin B$$

Proof outline

- suppose x is a particular, but arbitrarily chosen element of A
- prove that x is an element of B

$$A \subset B$$

$$A \subset B$$



$$A \subseteq B \wedge \exists x \in B, x \notin A$$

Definition of set equality

Definition of set equality

$$A = B$$



$$A \subseteq B \wedge B \subseteq A$$

Proof outline

- prove that $A \subseteq B$
- prove that $B \subseteq A$

Create a program that

- Takes two sets A and B represented as **arrays**
- Evaluates whether the sets are:
 - $A \subset B = -1$
 - $A = B = 0$
 - $A \supset B = 1$
 - otherwise -2

```
fun compareSets(a: IntArray, b: IntArray): Int {  
    if (a.size == 0) return if (b.size == 0) 0 else -1  
    if (b.size == 0) return 1  
    var result = 0  
    var ia = 0  
    var ib = 0  
    while (ia < a.size && ib < b.size) {  
        if (a[ia] == b[ib]) { ia++; ib++ }  
        else if (a[ia] > b[ib]) {  
            if (result == 1) return -2  
            ib++; result = -1 // a is subset  
        }  
        else if (a[ia] < b[ib]) {  
            if (result == -1) return -2  
            ia++; result = 1 // a is superset  
        }  
    }  
    if (ia == a.size && ib == b.size) return result;  
    return -2 // sets  
}
```

$$\emptyset = \{\}$$

There is only one empty set
The set with no elements

Partitions of sets

Sets A and B are disjoint if

$$A \cap B = \emptyset$$

Partions $\{A_1, A_2, A_3, \dots, A_n\}$ of set A

$$\square A = \bigcup_{i=1}^n A_i$$

$$\square \forall a, b \in \{1, 2, 3, \dots, n\}, A_a \cap A_b = \emptyset$$

$$\mathcal{P}(A)$$

The set of all subsets of A

$$\mathcal{P}(\{a, b, c\}) = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}, \{a, b, c\}\}$$

$$(A \cup B)^C = A^C \cap B^C$$

$$(A \cup B)^C = A^C \cap B^C$$

$$(A \cap B)^C = A^C \cup B^C$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$A \subseteq B \rightarrow A \cap B = A$$

$$A \subseteq B \rightarrow A \cap B = A$$

$$A \subseteq B \rightarrow A \cup B = B$$

Inclusion of intersection:

$$A \cap B \subseteq A$$

Inclusion in union:

$$A \subseteq A \cup B$$

Transitivity of subsets

$$A \subseteq B \wedge B \subseteq C \rightarrow A \subseteq C$$

of set operations

Intersection:

$$x \in X \cap Y \iff x \in X \wedge x \in Y$$

Union:

$$x \in X \cup Y \iff x \in X \vee x \in Y$$

Difference:

$$x \in X - Y \iff x \in X \wedge x \notin Y$$

Complement:

$$x \in X^C \iff x \notin X$$

Ordered Pair:

$$(x, y) \in X \times Y \iff x \in X \wedge y \in Y$$