

RWorksheet_Laguda#3a

Rey-Ann S. Laguda

2023-10-11

#1. VECTORS #A

```
first_eleven <- LETTERS[1:11]
first_eleven
```

```
## [1] "A" "B" "C" "D" "E" "F" "G" "H" "I" "J" "K"
```

#B

```
odd_letters <- LETTERS[seq(1,25,2)]
odd_letters
```

```
## [1] "A" "C" "E" "G" "I" "K" "M" "O" "Q" "S" "U" "W" "Y"
```

#C

```
vowels <- c(letters[which(letters %in% c("a","e","i","o","u"))],
            LETTERS[which(LETTERS %in% c("A","E","I","O","U"))])
vowels
```

```
## [1] "a" "e" "i" "o" "u" "A" "E" "I" "O" "U"
```

#D.

```
last5letter <- tail(letters, 5)
last5letter
```

```
## [1] "v" "w" "x" "y" "z"
```

#E

```
letterbetween15to24 <- letters[15:24]
letterbetween15to24
```

```
## [1] "o" "p" "q" "r" "s" "t" "u" "v" "w" "x"
```

#NUMBER 2 #A.

```
city <- c("Tuguegarao City", "Manila", "Iloilo City", "Tacloban", "Samal Island", "Davao City")
city
```

```
## [1] "Tuguegarao City" "Manila"           "Iloilo City"      "Tacloban"
```

```
## [5] "Samal Island"    "Davao City"
```

#B

```
temp <- c(42, 39, 34, 34, 30, 27)
temp
```

```
## [1] 42 39 34 34 30 27
```

#C

```
df <- data.frame(city, temp)
df
```

```
##           city temp
## 1 Tuguegarao City  42
## 2           Manila  39
## 3      Iloilo City  34
## 4       Tacloban  34
## 5     Samal Island  30
## 6       Davao City  27
```

```
#D
```

```
names(df) <- c("City", "Temperature")
df
```

```
##           City Temperature
## 1 Tuguegarao City         42
## 2           Manila         39
## 3      Iloilo City         34
## 4       Tacloban         34
## 5     Samal Island         30
## 6       Davao City         27
```

```
#E
```

```
str(df)
```

```
## 'data.frame':   6 obs. of  2 variables:
##  $ City          : chr  "Tuguegarao City" "Manila" "Iloilo City" "Tacloban" ...
##  $ Temperature: num  42 39 34 34 30 27
```

```
#F
```

```
df[3:4,]
```

```
##           City Temperature
## 3 Iloilo City         34
## 4   Tacloban         34
```

```
#G
```

```
highest_temp_city <- df$City[which.max(df$Temperature)]
lowest_temp_city  <- df$City[which.min(df$Temperature)]
highest_temp_city
```

```
## [1] "Tuguegarao City"
```

```
lowest_temp_city
```

```
## [1] "Davao City"
```

```
#2 MATRICES #A
```

```
m <- matrix(c(1:8,11:14),ncol = 4, nrow = 3)
m
```

```
##      [,1] [,2] [,3] [,4]
## [1,]    1    4    7   12
## [2,]    2    5    8   13
## [3,]    3    6   11   14
```

```
#B
```

```
matrix(c(1:8,11:14), ncol = 4, nrow = 3)*2
```

```
##      [,1] [,2] [,3] [,4]
## [1,]    2    8   14   24
## [2,]    4   10   16   26
## [3,]    6   12   22   28
```

```
#C
```

```
matrix(c(1:8,11:14), ncol = 4, nrow = 3)[2,]
```

```
## [1]  2  5  8 13
```

```
#D
```

```
matrix(c(1:8,11:14), ncol = 4, nrow = 3)[1:2,c(3,4)]
```

```
##      [,1] [,2]
## [1,]    7   12
## [2,]    8   13
```

```
#E
```

```
matrix(c(1:8,11:14), ncol = 4, nrow = 3) [3,c(2,3)]
```

```
## [1]  6 11
```

```
#F
```

```
matrix(c(1:8,11:14), ncol = 4, nrow = 3)[,4]
```

```
## [1] 12 13 14
```

```
#G
```

```
mat <- matrix(c(1:8,11:14)*2, ncol = 4, nrow = 3)
rownames(mat) <- c("isa", "dalawa", "tatlo")
colnames(mat) <- c("uno", "dos", "tres", "quatro")
mat
```

```
##      uno dos tres quatro
## isa      2  8  14    24
## dalawa   4 10  16    26
## tatlo    6 12  22    28
```

```
#H
```

```
dim(m) <- c(6,2)
```

```
m
```

```
##      [,1] [,2]
## [1,]    1    7
## [2,]    2    8
## [3,]    3   11
## [4,]    4   12
## [5,]    5   13
## [6,]    6   14
```

```
#3. ARRAYS #A
```

```
values <- c(1,2,3,6,7,8,9,0,3,4,5,1)
array_3d <- array(values, dim = c(2,4,3))
values
```

```
## [1] 1 2 3 6 7 8 9 0 3 4 5 1
```

```
array_3d
```

```
## , , 1
```

```
##
```

```
##      [,1] [,2] [,3] [,4]
```

```
## [1,]    1    3    7    9
```

```
## [2,]    2    6    8    0
```

```
##
```

```
## , , 2
```

```
##
```

```
##      [,1] [,2] [,3] [,4]
```

```
## [1,]    3    5    1    3
```

```
## [2,]    4    1    2    6
```

```
##
```

```
## , , 3
```

```
##
```

```
##      [,1] [,2] [,3] [,4]
```

```
## [1,]    7    9    3    5
```

```
## [2,]    8    0    4    1
```

```
#B
```

```
#B: The array has three(3) dimensions.
```

```
#C
```

```
dimnames(array_3d) <- list(c("a","b"),
```

```
LETTERS[1:4],c("1st-Dimensional Array", "2nd-Dimensional Array", "3rd-Dimensional Array"))
```

```
array_3d
```

```
## , , 1st-Dimensional Array
```

```
##
```

```
##   A B C D
```

```
## a 1 3 7 9
```

```
## b 2 6 8 0
```

```
##
```

```
## , , 2nd-Dimensional Array
```

```
##
```

```
##   A B C D
```

```
## a 3 5 1 3
```

```
## b 4 1 2 6
```

```
##
```

```
## , , 3rd-Dimensional Array
```

```
##
```

```
##   A B C D
```

```
## a 7 9 3 5
```

```
## b 8 0 4 1
```