

clone2vec
default, ≥ 3 cells
clone2vec
forced, ≥ 2 cells
Clonotrace
default, ≥ 10 cells
Clonotrace
forced, ≥ 2 cells
Benisse
≥ 2 cells
Time per dataset
Gut GC reporter
RBD reporter
NP reporter
GC + plasma cells
Plasmodium d10-42
Plasmodium d0-14
Human LN vaccine
Spontaneous GC
Human COVID-19
LARRY d2
LARRY d4/d6
Peak memory per dataset
Gut GC reporter
RBD reporter
NP reporter
GC + plasma cells
Plasmodium d10-42
Plasmodium d0-14
Human LN vaccine
Spontaneous GC
Human COVID-19
LARRY d2
LARRY d4/d6
Time with 1-16 cores (RBD)
1 core
2 cores
4 cores
8 cores
16 cores
Time on GSE195673 subsets
557 clones (5 %)
1,114 clones (10 %)
2,785 clones (25 %)
5,570 clones (50 %)
< 3 clones ≥ 10
≥ 10 cells
no BCR
not planned
10^{-1} 10^{0} 10^{1} 10^{2} 10^{3} 10^{4} 10^{5}
times Threadfin (log)
time: faster 0, slower 10
memory: less 2, more 8
cores: faster 0, slower 5
data size: faster 0, slower 4
times Threadfin (log)
time: faster 0, slower 11
memory: less 2, more 8
cores: faster 0, slower 5
data size: faster 0, slower 4
times Threadfin (log)
time: faster 0, slower 7
memory: less 0, more 7
cores: faster 0, slower 5
data size: faster 0, slower 4
times Threadfin (log)
time: faster 0, slower 11
memory: less 0, more 11
cores: faster 0, slower 5
data size: faster 0, slower 4
times Threadfin (log)
time: faster 0, slower 8
memory: less 3, more 6
cores: faster 0, slower 3
data size: faster 0, slower 4
hollow: a Clonotrace run whose clone distances used one R session
triangle: Clonotrace memory of the main R session only (a lower bound)