

NP–CGG | GSE305877 | WT day 14: 13,727 QC cells, 508 profiles; one identifiable mouse
A
scRNA clusters
Cell UMAP 2
Cell UMAP 1
B
GC score
RNA score
C
LZ score
D
Cycling score
E
Paired receptor coverage
Paired BCR
No eligible pair
F
Programme map-back
P1 P2 P3
G
Clone mean GC score
Clone UMAP 2
Clone UMAP 1
H
Programme stability
Bootstrap Jaccard
0.59
0.37
0.77
P1
P2
P3
RNA scores are descriptive overlays. 6,959 cells have eligible paired BCRs; 3,182 belong to displayed profiles. P1/P2 stability <0.75.
NP–OVA plate study | GSE246382 | day 14: 884 cells, 96 profiles; 11 mice
I
Recorded sorting gates
DZ
Myc+ LZ
LZ
PC
J
Sequence-derived W33L
germline W33
W33L
K
Mouse identity
M1 M4 M8 S3
M2 M5 S1 S4
M3 M6 S2
L
Profiled clone members
In ≥2-cell profile
Other cells
M
Clone PC-gate fraction
Fraction
N
Clone callable W33L fraction
O
W33L vs captured PC fraction
n = 74
Captured PC fraction
Callable W33L fraction
P
Programme map-back
Grey: unavailable annotation or no displayed profile. W33L fractions use callable cells only. Fractions describe captured cells, not output rates.