

பாகம் எடுத்துவிடப்பட்டது (கூறு சொல்) செப்டெம்பர் 2018 முதலில்.

கலை குருகு மாண்புமிகு (புது தாலி) மே 2018

General Certificate of Education (Adv. Level) Examination, August 2011

**வணக்கம் எல்லைய  
இனைந்த கணிதம்** II  
**Combined Mathematics** II

10 E II

08.08.2018 / 08:30 - 11:40

ஏட ஆண்டு  
மூன்று மணித்தியாலம்  
*Three hours*

அடுக்க விவரத்தை வெளியிட	- பின்தான் 10 மி மேலதிக வாசிப்பு நீரும் - 10 நிமிடங்கள்
<b>Additional Reading Time</b>	<b>10 minutes</b>

**Use additional reading time to go through the question paper, select the questions and decide on the questions that you give priority in answering.**

**Instructions:**

**Index Number** \_\_\_\_\_

- \* This question paper consists of two parts;  
**Part A** (Questions 1 – 10) and **Part B** (Questions 11 – 17)
  - \* **Part A:**  
Answer **all** questions. Write your answers to each question in the space provided. You may use additional sheets if more space is needed.
  - \* **Part B:**  
Answer **five** questions **only**. Write your answers on the sheets provided..
  - \* At the end of the time allotted, tie the answer scripts of the two parts together so that Part A is on top of Part B and hand them over to the supervisor.
  - \* You are permitted to remove only Part B of the question paper from the Examination Hall.
  - \* In this question paper,  $g$  denotes the acceleration due to gravity.

**For Examiners' Use only**

**(10) Combined Mathematics II**

<b>Part</b>	<b>Question No.</b>	<b>Marks</b>
<b>A</b>	1	
	2	
	3	
	4	
	5	
	6	
	7	
	8	
	9	
	10	
<b>B</b>	11	
	12	
	13	
	14	
	15	
	16	
	17	
<b>Total</b>		
<b>Percentage</b>		

<b>Paper I</b>	
<b>Paper II</b>	
<b>Total</b>	
<b>Final Marks</b>	

## **Final Marks**

In Numbers	
In Words	

## **Code Numbers**

<b>Marking Examiner</b>	
<b>Checked by:</b>	1
	2
<b>Supervised by:</b>	

## Part A


1. Two particles  $A$  and  $B$  of masses  $2m$  and  $m$  respectively, moving on a smooth horizontal table along the same straight line towards each other with the same speed  $u$ , collide directly. Just after collision, the particle  $A$  comes to rest. Show that the coefficient of restitution is  $\frac{1}{2}$ , and that the magnitude of the impulse exerted on  $B$  due to collision is  $2mu$ .

A hand-drawn illustration of a large, stylized bird or dragon-like creature, possibly a Roc, standing on a rocky cliff edge. The creature has a long neck, a large beak, and a patterned body. It is surrounded by a yellow aura and is positioned above a small, dark, circular object.

2. A particle is projected from a point on the horizontal ground, at an angle  $\alpha$  ( $0 < \alpha < \frac{\pi}{2}$ ) to the horizontal, with initial speed  $u = \sqrt{2gR}$ , where  $R$  is the horizontal range of the projectile on the ground. Show that the angle between the two possible initial directions of projection is  $\frac{\pi}{3}$ .

3. A particle  $P$  of mass  $m$  and a particle  $Q$  of mass  $\lambda m$  are attached to the two ends of a light inextensible string which passes over a smooth fixed pulley. The system is released from rest, with the string taut, as shown in the figure. The particle  $P$  moves downwards with acceleration  $\frac{g}{2}$ . Show that  $\lambda = \frac{1}{3}$ .

If the particle  $P$  strikes a horizontal inelastic floor with speed  $v$  and the particle  $Q$  never reaches the pulley, find the time taken by the particle  $Q$  to reach the maximum height from the instant when the particle  $P$  struck the floor.



4. A car of mass 1200 kg moves with the engine shut off, down a straight road of inclination  $\alpha$  to the horizontal, where  $\sin \alpha = \frac{1}{30}$ , at a certain constant speed. Taking the acceleration due to gravity  $g = 10 \text{ ms}^{-2}$ , find in newtons, the resistance to the motion of the car.


Find the power of the engine in kilowatts, when the car ascends the same road, under the same resistance with an acceleration  $\frac{1}{6} \text{ ms}^{-2}$ , at the instant when its speed is  $15 \text{ ms}^{-1}$ .

5. In the usual notation, let  $3\mathbf{i}$  and  $2\mathbf{i}+3\mathbf{j}$  be the position vectors of two points  $A$  and  $B$  respectively, with respect to a fixed origin  $O$ . Let  $C$  be the point on the straight line  $OB$  such that  $\hat{OCA} = \frac{\pi}{2}$ . Find  $\overrightarrow{OC}$  in terms of  $\mathbf{i}$  and  $\mathbf{j}$ .
- .....  
 .....

6. A uniform rod  $AB$  of length  $2a$  and weight  $W$ , is held in equilibrium by a light inextensible string  $BC$ , and by a horizontal force  $P$  applied at the end  $A$ , as shown in the figure. Given that the rod makes an angle  $45^\circ$  with the horizontal, show that angle  $\theta$  that the string  $BC$  makes with the horizontal is given by  $\tan \theta = 2$ .

Find the tension in the string, in this position, in terms of  $W$ .

.....  
 .....



7. Let  $A$  and  $B$  are two events in a sample space  $S$ . With the usual notation,  $P(A) = \frac{1}{3}$ ,  $P(B) = \frac{1}{4}$  and  $P(A \cap B) = \frac{1}{6}$ . Find  $P(A|B')$ ,  $P(A' \cap B')$  and  $P(B'|A')$ , where  $A'$  and  $B'$  denote complementary events of  $A$  and  $B$  respectively.

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

8. A bag contains 4 red balls and 3 black balls which are identical in all aspects except for their colour. Four balls are drawn from the bag at random, one at a time, without replacement. Find the probability that

- the balls drawn are of the same colour,
- the balls drawn, on any two consecutive draws are of different colours.

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

9. Five positive integers each of which is less than 8, have only one mode. Their mean, mode and median are in the ratios 6:10:5. Find these five integers.

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

10. Temperature in a certain city was recorded daily for 20 days. For this set of data the mean  $\mu$  and the standard deviation  $\sigma$  were calculated as  $28^{\circ}\text{C}$  and  $4^{\circ}\text{C}$  respectively. However, after detecting that two of the above temperatures have been mistakenly entered as  $35^{\circ}\text{C}$  and  $21^{\circ}\text{C}$ , they were later corrected as  $25^{\circ}\text{C}$  and  $31^{\circ}\text{C}$ . Find the correct values of  $\mu$  and  $\sigma$ .

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

.....  
 .....

**Department of Examinations, Sri Lanka**

ඇංග්‍රීසු පොදු සාහිත්‍ය පථ (ලේඛන රෝග) පිළිගෙය, 2018 අගෝස්තු

கல்வி போகுத் தாதுப் பதிர் (உயிர் துப்புவணம், 2018 குறைப்

General Certificate of Education (Adv. Level) Examination, August 2018

யൂ-ട്ടുബ് റിലീഫ്  
കുറഞ്ഞത് കണിതമ്  
**Combined Mathematics**

10 E II

## Part B

\* Answer five questions only.

(In this question paper,  $g$  denotes the acceleration due to gravity.)

- 11.(a) A lift moving in a mine pit of depth  $4d$  metres begins to move vertically downwards from rest from a point  $A$ , at time  $t = 0$ . First, it moves a distance  $d$  metres with constant acceleration  $\frac{g}{2} \text{ m s}^{-2}$ , and then it moves with the velocity attained at the end of that motion, for another distance of  $d$  metres. The lift then moves the remaining distance with constant retardation so as to come to rest exactly at the point  $B$  at a distance  $4d$  metres below  $A$ .

Sketch the velocity-time graph for the motion of the lift.

Hence, find the total time taken by the lift to move down from A to B.


- (b) A ship is sailing due North with uniform speed  $u \text{ km h}^{-1}$ , relative to earth. At a certain instant a boat  $B_1$  is observed at an angle  $\beta$  East of South, from the ship at a distance  $p \text{ km}$  from the path of the ship. At the same instant, a boat  $B_2$  is observed Westward at a distance  $q \text{ km}$  from the ship. Both boats sail in straight line paths, with uniform speed  $v (> u) \text{ km h}^{-1}$  relative to earth, intending to intercept the ship. Sketch in the same diagram, velocity triangles to determine the paths of the boats relative to earth.

Show that the path of the boat  $B_1$ , relative to earth makes an angle  $\beta - \sin^{-1}\left(\frac{u \sin \beta}{v}\right)$  West of North, and find the path of the boat  $B_2$ , relative to earth.

Let  $\beta = \frac{\pi}{3}$  and  $v = \sqrt{3}u$ . Show that if  $3q^2 > 8p^2$ , then the boat  $B_1$  will intercept the ship before the boat  $B_2$ .

- 12.(a) The trapezium  $ABCD$  such that  $AB = a$  and  $\hat{B}AD = \frac{\pi}{6}$ , shown in the figure, is a vertical cross-section through the centre of gravity of a smooth uniform block of mass  $2m$ . The lines  $AD$  and  $BC$  are parallel, and the line  $AB$  is a line of greatest slope of the face containing it. The block is placed with the face containing  $AD$  on a smooth horizontal floor. A particle  $P$  of mass  $m$  is placed at the point  $A$  and it is given a velocity  $u$  along  $\overrightarrow{AB}$ , where  $u^2 = \frac{7ga}{3}$ , as shown in the figure. Show that the retardation of  $P$  relative to the block is  $\frac{2g}{3}$  and find the velocity of the particle  $P$  relative to the block when the particle  $P$  reaches  $B$ .

Also, there is a small hole on the upper face of the block at the point  $E$  on  $BC$  such that  $BE = \frac{\sqrt{3}a}{2}$ . By considering the motion relative to the block, show that the particle  $P$  will fall into the hole at  $E$ .



(b) One end of a light inextensible string of length  $a$  is attached to a fixed point  $O$  and the other end to a particle  $P$  of mass  $m$ . The particle hangs at rest vertically below  $O$ , and it is given a horizontal velocity of magnitude  $u = \sqrt{kag}$ , where  $2 < k < 5$ . Show that, when the string has turned through an angle  $\theta$  and still taut, the speed  $v$  of the particle is given by  $v^2 = (k - 2)ag + 2ag \cos \theta$ .

Find the tension in the string, in this position.

Deduce that the string becomes slack when  $\theta = \alpha$ , where  $\cos \alpha = \frac{2-k}{3}$ .

13. A particle  $P$  of mass  $m$  is attached to two ends of two equal light elastic strings, each of natural length  $a$  and modulus  $mg$ . The free end of one string is attached to a fixed point  $A$ , and the free end of the other string is attached to a fixed point  $B$  which is at a distance  $4a$  vertically below  $A$ . (See the figure.) Show that, with both strings taut, the particle will be in equilibrium at a distance  $\frac{5a}{2}$  below  $A$ .

The particle  $P$  is now raised to the mid-point of  $AB$  and is gently released from rest in that position. When both strings are taut and the length of the string  $AP$  is  $x$ , show that  $\ddot{x} + \frac{2g}{a} \left( x - \frac{5a}{2} \right) = 0$ .


Re-write this equation in the form  $\ddot{X} + \omega^2 X = 0$ , where  $X = x - \frac{5a}{2}$  and  $\omega^2 = \frac{2g}{a}$ .

Using the formula  $\dot{X}^2 = \omega^2(c^2 - X^2)$ , find the amplitude  $c$  of this motion.

At the instant when the particle  $P$  reaches its lowest position, the string  $PB$  is cut.

Show that the particle reaches its highest position in the new motion when  $x = a$ .

Show further that the total time taken by the particle  $P$  to move from its initial position at  $x = 2a$ , a distance  $a$  downwards, and then a distance  $\frac{a}{2}$  upwards is  $\frac{\pi}{3} \sqrt{\frac{a}{2g}} (3 + \sqrt{2})$ .



- 14.(a) Let  $OAB$  be a triangle,  $D$  be the mid-point of  $AB$  and  $E$  be the mid-point of  $OD$ . The point  $F$  lies on  $OA$  such that  $OF : FA = 1 : 2$ . The position vectors of  $A$  and  $B$  with respect to  $O$  are  $\mathbf{a}$  and  $\mathbf{b}$  respectively. Express the vectors  $\overrightarrow{BE}$  and  $\overrightarrow{BF}$  in terms of  $\mathbf{a}$  and  $\mathbf{b}$ .

Deduce that  $B$ ,  $E$  and  $F$  are collinear and find the ratio  $BE : EF$ .

Find the scalar product  $\overrightarrow{BF} \cdot \overrightarrow{DF}$  in terms of  $|\mathbf{a}|$  and  $|\mathbf{b}|$  and show that if  $|\mathbf{a}| = 3|\mathbf{b}|$ , then  $\overrightarrow{BF}$  is perpendicular to  $\overrightarrow{DF}$ .

- (b) A system of forces in the  $Oxy$ -plane consists of three forces  $3Pi + 2Pj$ ,  $2Pi - Pj$  and  $-Pi + 2Pj$  acting at the points  $(-a, 2a)$ ,  $(0, a)$  and  $(-a, 0)$  respectively, where  $P$  and  $a$  are positive quantities measured in newtons and metres, respectively. Show that the clockwise moment of the system about the origin  $O$  is  $12Pa$  N m.


Also, show that the system is equivalent to a single resultant force of magnitude  $5PN$  and find its direction and the equation of the line of action.

Now, an extra force is introduced to the system so that the new system is equivalent to a couple with clockwise moment  $24Pa$  N m. Find the magnitude, direction and the equation of the line of action of the extra force.

- 15.(a) A uniform rod  $AB$  of weight  $W$  and length  $2a$ , has its end  $A$  placed on a rough horizontal ground and the end  $B$  against a smooth vertical wall. The rod lies in a vertical plane perpendicular to the wall and makes an angle  $\theta$  with the horizontal, where  $\tan \theta = \frac{3}{4}$ . A particle of weight  $W$  is attached to the point  $C$  on the rod with  $AC = x$ . The rod with the particle is in equilibrium. The coefficient of friction between the rod and the ground is  $\frac{5}{6}$ . Show that  $x \leq \frac{3a}{2}$ .


- (b) The framework shown in the adjoining figure is made of five light rods  $AB$ ,  $BC$ ,  $AC$ ,  $CD$  and  $AD$  freely jointed at their ends. It is given that  $AB = a$ ,  $BC = 2a$ ,  $AC = CD$  and  $\angle CAD = 30^\circ$ . A load of weight  $W$  hangs at  $D$  and the framework is in equilibrium in a vertical plane with  $AB$  horizontal and  $AC$  vertical, supported by vertical forces  $P$  and  $Q$  acting respectively at  $A$  and  $B$  in the directions indicated in the figure. Find the value of  $Q$ , in terms of  $W$ .

Draw a stress diagram using Bow's notation and hence, find the stresses in the five rods and state whether these stresses are tensions or thrusts.



16. Show that the centre of mass of a uniform solid hemisphere of radius  $a$  is at a distance  $\frac{3}{8}a$  from its centre.


A hemispherical portion of radius  $a$  is carved out from a uniform solid right circular cylinder of radius  $a$ , height  $a$  and density  $\rho$ . Now, a uniform solid hemisphere of radius  $a$  and density  $\lambda\rho$  is fastened at its circular face to the circular face of the remaining portion of the cylinder, so that the two axes of symmetry coincide, as shown in the adjoining figure. Show that the centre of mass of the body  $S$  thus formed, lies on its axis of symmetry at a distance  $\frac{(11\lambda + 3)a}{4(2\lambda + 1)}$  from  $O$ , the centre of the rim.



Let  $\lambda = 2$  and let  $A$  be a point on the circular rim of the body  $S$ .

This body  $S$  is kept in equilibrium against a rough vertical wall by means of a light inextensible string with one end attached to the point  $A$  and the other end to a fixed point  $B$  on the vertical wall. In this equilibrium position, the axis of symmetry of  $S$  is perpendicular to the wall and the hemispherical surface of  $S$  touches the wall at the point  $C$  at a distance  $3a$  vertically below the point  $B$ . (See the adjoining figure.) The points  $O$ ,  $A$ ,  $B$  and  $C$  lie on a vertical plane perpendicular to the wall.

If  $\mu$  is the coefficient of friction between the hemispherical surface of  $S$  and the wall, show that  $\mu \geq 3$ .



17.(a) All applicants who apply for a certain job in an institute are required to sit for an aptitude test. Those who obtain A grades in this aptitude test are selected for the job, and the rest of the applicants have to face an interview. In a survey, it was found that 60% of the applicants get A grades, and of these, 40% are females. From the applicants who face the interview, only 10% get selected and 70% of them are females.

Find the probability that

- a male is selected for this job,
- a male who has been selected for the job has obtained an A grade for the aptitude test.

(b) Waiting times (in minutes) before being treated of 100 patients at a certain hospital are collected. The following table gives the distribution of values obtained by subtracting 20 minutes from each of these times and dividing each of the resulting differences by 10.

Range of values	Number of patients
-2 – 0	30
0 – 2	40
2 – 4	15
4 – 6	10
6 – 8	5

Estimate the mean and the standard deviation of the distribution given in this table.

Hence, estimate the mean  $\mu$  and the standard deviation  $\sigma$  of the waiting times of the 100 patients.

Also, estimate the coefficient of skewness  $\kappa$  defined by  $\kappa = \frac{\mu - M}{\sigma}$ , where  $M$  is the mode of the waiting times of the 100 patients.

\*\*\*