

நிலை எண் காலை தா (குமார் சென்டீ, 2016 காலை காலை) குமார் துறைப் பகுதி (ஏ.ஐ. பி.) மூலம், 2016 காலை General Certificate of Education (Adv. Level) Examination, August 2016

ஸங்கிள்வத மாணிக்ய	II
இணைந்த கணிதம்	II
Combined Mathematics	II

10 E II

ஒட்டு நூற்று
மண்ணு மணித்தியாலும்
Three hours

Index Number

Instructions:

- * This question paper consists of two parts;
Part A (Questions 1–10) and **Part B** (Questions 11–17)
 - * **Part A:**
Answer **all** questions. Write your answers to each question in the space provided. You may use additional sheets if more space is needed.
 - * **Part B:**
Answer **five** questions **only**. Write your answers on the sheets provided.
 - * At the end of the time allotted, tie the answer scripts of the two parts together so that Part A is on top of Part B and hand them over to the supervisor.
 - * You are permitted to remove **only** Part B of the question paper from the Examination Hall.
 - * In this question paper, g denotes the acceleration due to gravity.

For Examiners' Use only

(10) Combined Mathematics II

Part	Question No.	Marks
A	1	
	2	
	3	
	4	
	5	
	6	
	7	
	8	
	9	
	10	
B	11	
	12	
	13	
	14	
	15	
	16	
	17	
	Total	
	Percentage	

Paper I	
Paper II	
Total	
Final Marks	

Final Marks

In Numbers	
In Words	


Code Numbers

Marking Examiner	
Checked by:	1
	2
Supervised by:	

Part A


1. A particle of mass m hangs in equilibrium at one end of a light inextensible string of length l whose other end is tied to a fixed point O . Another particle of mass $2m$ collides horizontally with velocity u with the first particle and coalesces with it. Find the velocity with which the composite particle begins to move.

Show that if $u = \sqrt{gl}$, then the composite particle reaches a maximum height of $\frac{2l}{9}$ above its initial level.




2. A particle P of mass m and a particle Q of mass $3m$ move on a smooth horizontal table along the same straight line towards each other with speeds $5u$ and u respectively, as shown in the figure. After their impact, P and Q move away from each other with speeds u and v respectively.

Find v in terms of u , and show that the coefficient of restitution between P and Q is $\frac{1}{3}$.



3. A particle P , projected horizontally with velocity u given by $u = \frac{3}{2} \sqrt{ga}$ from a point A at the edge of a step of a fixed stairway perpendicular to that edge, moves under gravity. Each step is of height a and length $2a$ (see the figure). Show that the particle P will not hit the first step below A , and it will hit the second step below A at a horizontal distance $3a$ from A .



4. A car of mass M kg moves along a straight level road against a resistance of constant magnitude R N. At an instant when the car is moving at speed v m s $^{-1}$, its acceleration is a m s $^{-2}$. Show that the power of its engine at this instant is $(R + Ma)v$ W.

The car then moves with a constant speed v_1 m s $^{-1}$ against a resistance of the same constant magnitude R N up a straight road inclined at an angle α to the horizontal, working at the same power. Show that $v_1 = \frac{(R + Ma)v}{R + Mg \sin \alpha}$.




5. In the usual notation, let $\mathbf{a} = 3\mathbf{i} + 4\mathbf{j}$, $\mathbf{b} = 4\mathbf{i} + 3\mathbf{j}$ and $\mathbf{c} = \alpha\mathbf{i} + (1 - \alpha)\mathbf{j}$, where $\alpha \in \mathbb{R}$.


Find (i) $|a|$ and $|b|$.

(ii) $a \cdot c$ and $b \cdot c$ in terms of a .

If the angle between a and c is equal to the angle between b and c , show that $\alpha = \frac{1}{2}$.



6. One end of a light inextensible string of length $2l$ is attached to the highest point of a thin smooth rigid circular wire of radius $a (> \sqrt{2}l)$ which is fixed in a vertical plane. A small smooth bead of weight w , which is free to move along the wire, is attached to the other end of the string. The bead is in equilibrium with the string taut, as in the figure. Mark the forces acting on the bead and show that the tension of the string is $\frac{2wl}{a}$.



7. Let A and B be two events of a sample space Ω . In the usual notation, $P(A) = p$, $P(B) = \frac{p}{2}$ and $P(A \cup B) - P(A \cap B) = \frac{2p}{3}$, where $p > 0$. Find $P(A \cap B)$ in terms of p .
Deduce that if A and B are independent events, then $p = \frac{5}{6}$.

Deduce that if A and B are independent events, then $p = \frac{5}{6}$.

8. A bag contains 6 white balls and n black balls which are equal in all respects, except for colour. Two balls are taken out at random from the bag, one after the other, without replacement. The probability that the first ball is white and the second ball is black is $\frac{4}{15}$. Find the value of n .

9. The mean of three distinct integers less than 11 is 7. When two more integers are taken, the mean of all five integers is 5. Also, the only mode of these five integers is 3. Find the five integers.

10. An arrow is shot at a rotating circular target-board consisting of five equal sectors numbered 1, 2, 3, 4 and 5. The number of times the arrow hits each of the sectors is given in the following frequency table, where p and q are constants.

Number	1	2	3	4	5
Frequency	1	p	q	5	2

If the mean and the variance of the above data are given to be 3 and $\frac{6}{5}$ respectively, find the values of p and q .

கிடைத் தமிழ்நாடு/முழுப் பதிப்பு/கொலையெடு/All Rights Reserved]

විද්‍යාත් නො සැකිනු යුතු (Good bye) Session, 2016 වෛශ්‍ය පිටපත

கல்விப் போக்கு துறைப் பகுதி (பார் து)ப் பிர'சூ, 2016 ஒன்று

General Certificate of Education (Adv. Level) Examination, August 2016

கலைக்கு மாணிக்கை II
இணைந்த கணிதம் II
Combined Mathematics II


10 E II

PART B

* Answer five questions only.

(In this question paper, g denotes the acceleration due to gravity.)


- 11.(a) A particle P of mass m is connected to a particle Q of mass $3m$ by a light inextensible string passing over a small smooth pulley fixed at a height $3h$ above an inelastic horizontal floor. Initially the two particles are held at a height h above the floor with the string taut, and released from rest. (See the adjoining figure.) Applying Newton's second law separately to the motions of P and Q , show that the magnitude of acceleration of each particle is $\frac{g}{2}$.



- Using these graphs, show that $t_0 = 2\sqrt{\frac{h}{g}}$ and find t_1 in terms of g and h .

Show further that the particle P reaches a maximum height $\frac{5h}{2}$ above the floor.

- (b) A straight river of breadth a flows with uniform speed u . The points A and C are situated on opposite banks of the river such that the line AC is perpendicular to the direction of flow of the river. Also, a stationary buoy B is fixed in the middle of the river, on the upstream side of AC such that ABC is an equilateral triangle. (See the adjoining figure.) A boat moving with speed v ($> u$) relative to water starts off from A and moves until it reaches B . Then it moves from B to C . Sketch the velocity triangles for the motions of the boat from A to B and from B to C .




Show that the speed of the boat in its motion from A to B is $\frac{1}{2}(\sqrt{4v^2 - u^2} - \sqrt{3}u)$ and find its speed in the motion from B to C .

Hence, show that the total time taken by the boat for the paths AB and BC is $\frac{a\sqrt{4v^2 - u^2}}{\frac{v^2 - u^2}{v^2 + u^2}}$.


- 12.(a) The triangle ABC in the figure is a vertical cross-section through the centre of gravity of a uniform wedge of mass $2m$. The line AB is a

line of greatest slope of the face containing it and $\hat{ABC} = \frac{\pi}{4}$. The wedge is placed with the face containing BC on a rough horizontal floor. The face containing AB is smooth. A particle of mass m is held on AB as in the figure and the system is released from rest. It is given that the wedge moves in the direction of \vec{BC} and that



the magnitude of the frictional force exerted on the wedge by the floor is $\frac{R}{6}$, where R is the magnitude of the normal reaction exerted on the wedge by the floor. Obtain equations which are sufficient to determine R , in terms of m and g .

(b) OAB in the figure is a circular sector of radius a subtending an angle $\frac{\pi}{6}$ at the centre O with OA vertical. It is a cross-section perpendicular to the axis of a smooth cylindrical sector fixed with its axis horizontal. One end of a light inextensible string passing over a small smooth pulley fixed at B is attached to a particle P of mass $3m$ and the other end is attached to a particle Q of mass m . Initially, the particle P is held at A and the particle Q hangs freely at the horizontal level of O . The system is released from rest in this position, with the string taut. When OP makes an angle θ ($0 < \theta < \frac{\pi}{6}$) with the upward vertical, show that $2a\dot{\theta}^2 = 3g(1 - \cos \theta) + g\theta$ and that the tension in the string is $\frac{3}{4}mg(1 - \sin \theta)$, and find the normal reaction on the particle P .



13. One end of a light elastic string of natural length a and modulus of elasticity $4mg$ is tied to a fixed point O and the other end to a particle P of mass m . The particle P is released from rest at O . Find the velocity of the particle P when it passes through the point A , where $OA = a$.

Show that the length of the string $x (\geq a)$ satisfies the equation $\ddot{x} + \frac{4g}{a} \left(x - \frac{5a}{4} \right) = 0$.

Taking $X = x - \frac{5a}{4}$, express the above equation in the form $\ddot{X} + \omega^2 X = 0$, where $\omega (> 0)$ is a constant to be determined.

Assuming that $\dot{X}^2 = \omega^2 (c^2 - X^2)$, find the amplitude c of this simple harmonic motion.

Let L be the lowest point reached by the particle P . Show that the time taken by P to move from A to L is $\frac{1}{2} \sqrt{\frac{a}{g}} \left\{ \pi - \cos^{-1} \left(\frac{1}{3} \right) \right\}$.

At the instant when the particle P is at L , another particle of mass λm ($1 \leq \lambda < 3$) is gently attached to P . Show that the equation of motion of the composite particle of mass $(1+\lambda)m$ is $\ddot{x} + \frac{4g}{(1+\lambda)a} \left\{ x - (5+\lambda) \frac{a}{4} \right\} = 0$.

Show further that the composite particle performs complete simple harmonic motion with amplitude $(3-\lambda) \frac{a}{4}$.

14. (a) The position vectors of two points A and B with respect to an origin O are \mathbf{a} and \mathbf{b} respectively, where O, A and B are not collinear. Let C be the point such that $\overrightarrow{OC} = \frac{1}{3}\overrightarrow{OB}$ and let D be the point such that $\overrightarrow{OD} = \frac{1}{2}\overrightarrow{AB}$. By expressing \overrightarrow{AC} and \overrightarrow{AD} in terms of \mathbf{a} and \mathbf{b} , show that $\overrightarrow{AD} = \frac{3}{2}\overrightarrow{AC}$.

Let P and Q be the points on AB and OD respectively, such that $\overrightarrow{AP} = \lambda \overrightarrow{AB}$ and $\overrightarrow{OQ} = (1-\lambda) \overrightarrow{OD}$, where $0 < \lambda < 1$. Show that $\overrightarrow{PC} = 2\overrightarrow{CQ}$.

- (b) In a parallelogram $ABCD$, let $AB = 2$ m and $AD = 1$ m, and let $\hat{\angle}BAD = \frac{\pi}{3}$. Also, let E be the mid-point of CD . Forces of magnitudes 5, 5, 2, 4 and 3 newtons act along AB, BC, DC, DA and BE respectively, in the directions indicated by order of the letters. Show that their resultant force is parallel to \overrightarrow{AE} , and find its magnitude.

Also, show that the line of action of the resultant force meets AB produced at a distance $\frac{3}{2}$ m from B .

An additional force acting through C is now added to the above system of forces so that the resultant force of the new system is along \overrightarrow{AE} . Find the magnitude and direction of the additional force.


15. (a) Four equal uniform rods, each of weight w_1 , are smoothly jointed at their ends to form a rhombus $ABCD$. The mid-points of BC and CD are connected by a light rod such that $B\hat{A}D = 2\theta$. Each of the joints B and D carries equal loads of weight w_2 . The system, hanging symmetrically from the joint A , is in equilibrium in a vertical plane with the light rod horizontal. Show that the thrust in the light rod is $2(2w_1 + w_2) \tan \theta$.

- (b) The adjoining figure represents a framework of five light rods AB , BC , CD , AC and AD , smoothly jointed at the ends.

It is given that $AC = CB$ and $B\hat{A}C = 30^\circ = A\hat{D}C$. The framework is smoothly hinged at D . A weight W is suspended at the joint B and the framework is kept in equilibrium in a vertical plane with AB horizontal and AD vertical, by a horizontal force of magnitude X acting at A .


Using Bow's notation, draw stress diagrams for the joints B , C and A in the same figure.

Hence, find the value of X and the stresses in all rods, distinguishing between tensions and thrusts.




16. Show that the centre of mass of a uniform semi-circular lamina of radius r and centre O is at a distance $\frac{4r}{3\pi}$ from O .

As shown in the adjoining figure, a uniform plane lamina L is made by rigidly attaching a rectangle $ABCD$ to a square $PQRS$ such that DC and PQ lie on the same line with their mid-points coinciding, and removing a semi-circular region XYZ of radius $\frac{a}{2}$ centred at the mid-point T of RS . It is given that $AB = a$ and $AD = PQ = 2a$. Show that the centre of mass of the lamina L lies on the axis of symmetry at a distance ka from RS , where $k = \frac{238}{3(48 - \pi)}$.



As shown in the adjoining figure, the lamina L is in equilibrium on a rough plane inclined at an angle α to the horizontal with its plane vertical and the edge PS on a line of greatest slope such that the point P lies below S . Show that $\tan \alpha < (2 - k)$ and $\mu \geq \tan \alpha$, where μ is the coefficient of friction between the lamina and the inclined plane.



17.(a) An unbiased cubical die A shows 1, 2, 3, 3, 4, 5 on its six separate faces. The die A is tossed twice. Find the probability that the sum of the two numbers obtained is 6.

Another die B , identical to A in all respects except for the numbers on the faces, shows 2, 2, 3, 4, 4, 5 on its six separate faces. The die B is tossed twice. Find the probability that the sum of the two numbers obtained is 6.

Now, the two dice A and B are put in a box. One die is taken out of the box at random and tossed twice. Given that the sum of the two numbers obtained is 6, find the probability that the die taken out of the box is the die A .

(b) The mean and the standard deviation of n numbers x_1, x_2, \dots, x_n are μ_1 and σ_1 respectively, and the mean and the standard deviation of m numbers y_1, y_2, \dots, y_m are μ_2 and σ_2 respectively. Let the mean and the standard deviation of all of these $n + m$ numbers be μ_3 and σ_3 respectively.

$$\text{Show that } \mu_3 = \frac{n\mu_1 + m\mu_2}{n + m}.$$

$$\text{Let } d_1 = \mu_3 - \mu_1. \text{ Show that } \sum_{i=1}^n (x_i - \mu_3)^2 = n(\sigma_1^2 + d_1^2).$$

$$\text{By taking } d_2 = \mu_3 - \mu_2, \text{ write down a similar expression for } \sum_{j=1}^m (y_j - \mu_3)^2.$$

$$\text{Deduce that } \sigma_3^2 = \frac{(n\sigma_1^2 + m\sigma_2^2) + (nd_1^2 + md_2^2)}{n + m}.$$

The number of copies sold per day, during the first 100 days after publishing a new book, had mean 2.3 and variance 0.8. During the next 100 days, the number of copies sold per day had mean 1.7 and variance 0.5. Find the mean and the variance of the number of copies sold per day during the first 200 days.
