

Lecture 20: Single Proportion Test

Chapter 6.1

Question for Today

According to a poll done by the New York Times/CBS News in June 2012, only about 44% of the American public approved of the Supreme Court's performance.

Question for Today

According to a poll done by the New York Times/CBS News in June 2012, only about 44% of the American public approved of the Supreme Court's performance.

The sample proportion $\hat{p} = 0.44$ is **point estimate** of p : the true proportion of the American public who approves.

Question for Today

According to a poll done by the New York Times/CBS News in June 2012, only about 44% of the American public approved of the Supreme Court's performance.

The sample proportion $\hat{p} = 0.44$ is **point estimate** of p : the true proportion of the American public who approves.

What are some next things to ask?

Question for Today

According to a poll done by the New York Times/CBS News in June 2012, only about 44% of the American public approved of the Supreme Court's performance.

The sample proportion $\hat{p} = 0.44$ is **point estimate** of p : the true proportion of the American public who approves.

What are some next things to ask?

- ▶ What was n ?
- ▶ What is the **SE** of $\hat{p} = 44\% = 0.44$?
- ▶ What is the sampling distribution of \hat{p} ?

Question for Today

Just like with \bar{x} , if we want to use the normal model to

- ▶ build confidence intervals via z^*
- ▶ conduct hypothesis tests via the normal tables

we need the **sampling distribution** of \hat{p} to be nearly normal.

Question for Today

Just like with \bar{x} , if we want to use the normal model to

- ▶ build confidence intervals via z^*
- ▶ conduct hypothesis tests via the normal tables

we need the **sampling distribution** of \hat{p} to be nearly normal.

This happens when the population distribution of 0's and 1's is not too strongly skewed. As the sample size $n \rightarrow \infty$, this is less of an issue by the CLT.

Question for Today

Just like with \bar{x} , if we want to use the normal model to

- ▶ build confidence intervals via z^*
- ▶ conduct hypothesis tests via the normal tables

we need the **sampling distribution** of \hat{p} to be nearly normal.

This happens when the population distribution of 0's and 1's is not too strongly skewed. As the sample size $n \rightarrow \infty$, this is less of an issue by the CLT.

Note:

$$\hat{p} = \frac{x_1 + \dots + x_n}{n} = \frac{1}{n} \sum_{i=1}^n x_i$$

where each of the x_i 's are 0/1 success/failure **Bernoulli** random variables.

Conditions for Sampling Dist'n of \hat{p} Being Nearly Normal

The sampling distribution of the sample proportion \hat{p} based on sample size n is nearly normal when

Conditions for Sampling Dist'n of \hat{p} Being Nearly Normal

The sampling distribution of the **sample proportion** \hat{p} based on sample size n is nearly normal when

- ▶ The observations are independent: the 10% rule

Conditions for Sampling Dist'n of \hat{p} Being Nearly Normal

The sampling distribution of the **sample proportion** \hat{p} based on sample size n is nearly normal when

- ▶ The observations are independent: the 10% rule
- ▶ We expect to see at least 10 successes and 10 failures in our sample. This is called the **success-failure condition**:
 - ▶ $np \geq 10$
 - ▶ $n(1 - p) \geq 10$

Conditions for Sampling Dist'n of \hat{p} Being Nearly Normal

If conditions are met, then the sampling distribution of \hat{p} is nearly normal with

- ▶ mean p (the true population proportion)
- ▶ standard error

$$SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$$

Conditions for Sampling Dist'n of \hat{p} Being Nearly Normal

If conditions are met, then the sampling distribution of \hat{p} is nearly normal with

- ▶ mean p (the true population proportion)
- ▶ standard error

$$SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$$

Note the similarity of the previous formula for the sample mean \bar{x} :

$$SE_{\bar{x}} = \frac{\sigma}{\sqrt{n}} = \sqrt{\frac{\sigma^2}{n}}$$

What p to use?

But we **don't know** what p is. So what p do we use

- ▶ to check the success/failure condition?

- ▶ for the $SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$?

For

What p to use?

But we **don't know** what p is. So what p do we use

- ▶ to check the success/failure condition?

- ▶ for the $SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$?

For

- ▶ Confidence intervals: plug in the **point estimate** \hat{p} of p

What p to use?

But we **don't know** what p is. So what p do we use

- ▶ to check the success/failure condition?

- ▶ for the $SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$?

For

- ▶ Confidence intervals: plug in the **point estimate** \hat{p} of p
- ▶ Hypothesis tests: plug in the **null value** p_0 from $H_0 : p = p_0$

Confidence Intervals

Going back to the poll: $\hat{p} = 0.44$ based on $n = 976$. What is a 95% confidence interval?

Confidence Intervals

Going back to the poll: $\hat{p} = 0.44$ based on $n = 976$. What is a 95% confidence interval?

Check the conditions and find SE using $p = \hat{p}$

Confidence Intervals

Going back to the poll: $\hat{p} = 0.44$ based on $n = 976$. What is a 95% confidence interval?

Check the conditions and find SE using $p = \hat{p}$

- ▶ $976 < 10\%$ of 313 million \Rightarrow independence

Confidence Intervals

Going back to the poll: $\hat{p} = 0.44$ based on $n = 976$. What is a 95% confidence interval?

Check the conditions and find SE using $p = \hat{p}$

- ▶ $976 < 10\%$ of 313 million \Rightarrow independence
- ▶ Defining a success as a person approving of the job done by the Supreme Court:

Confidence Intervals

Going back to the poll: $\hat{p} = 0.44$ based on $n = 976$. What is a 95% confidence interval?

Check the conditions and find SE using $p = \hat{p}$

- ▶ $976 < 10\%$ of 313 million \Rightarrow independence
- ▶ Defining a success as a person approving of the job done by the Supreme Court:
 - ▶ $976 \times \hat{p} = 976 \times .44 = 429$ successes ≥ 10
 - ▶ $976 \times (1 - \hat{p}) = 976 \times .56 = 547$ failures ≥ 10

Confidence Intervals

Going back to the poll: $\hat{p} = 0.44$ based on $n = 976$. What is a 95% confidence interval?

Check the conditions and find SE using $p = \hat{p}$

- ▶ $976 < 10\%$ of 313 million \Rightarrow independence
- ▶ Defining a success as a person approving of the job done by the Supreme Court:
 - ▶ $976 \times \hat{p} = 976 \times .44 = 429$ successes ≥ 10
 - ▶ $976 \times (1 - \hat{p}) = 976 \times .56 = 547$ failures ≥ 10

$$SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}} \approx \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} = \sqrt{\frac{0.44(1-0.44)}{976}} = 0.016$$

Confidence Intervals

A 95% confidence interval using the normal model has $z^* = 1.96$,
thus:

$$\text{point estimate} \pm 1.96 \times SE$$

Confidence Intervals

A 95% confidence interval using the normal model has $z^* = 1.96$, thus:

$$\text{point estimate} \pm 1.96 \times SE$$

In our case

$$\hat{p} \pm 1.96 \times SE_{\hat{p}} = 0.44 \pm 1.96 \times 0.016 = (0.409, 0.471)$$

Hypothesis Tests

Thomas Carcetti is running for mayor of Baltimore. His campaign manager **claims** he has more than 50% support of the electorate.

Hypothesis Tests

Thomas Carcetti is running for mayor of Baltimore. His campaign manager **claims** he has more than 50% support of the electorate.

The Baltimore Sun collects a random sample of $n = 500$ likely voters and finds that 52% support him. Does this provide convincing evidence for the claim of Carcetti's manager at the 5% significance level?

Hypothesis Tests

The hypothesis test is, with the null value $p_0 = 0.5$

$$\begin{array}{l} H_0 : p = p_0 \\ \text{vs} \quad H_A : p > p_0 \end{array}$$

Hypothesis Tests

The hypothesis test is, with the null value $p_0 = 0.5$

$$\begin{array}{l} H_0 : p = p_0 \\ \text{vs} \quad H_A : p > p_0 \end{array}$$

Check the conditions and find SE using $p = p_0$

Hypothesis Tests

The hypothesis test is, with the null value $p_0 = 0.5$

$$\begin{array}{l} H_0 : p = p_0 \\ \text{vs} \quad H_A : p > p_0 \end{array}$$

Check the conditions and find SE using $p = p_0$

- ▶ $500 < 10\%$ of the population of Baltimore \Rightarrow independence

Hypothesis Tests

The hypothesis test is, with the null value $p_0 = 0.5$

$$\begin{array}{l} H_0 : p = p_0 \\ \text{vs} \quad H_A : p > p_0 \end{array}$$

Check the conditions and find SE using $p = p_0$

- ▶ $500 < 10\%$ of the population of Baltimore \Rightarrow independence
- ▶ Success-failure condition
 - ▶ $np_0 = 500 \times 0.5 = 250 \geq 10$
 - ▶ $n(1 - p_0) = 500 \times (1 - 0.5) = 250 \geq 10$

Hypothesis Tests

The hypothesis test is, with the null value $p_0 = 0.5$

$$\begin{array}{l} H_0 : p = p_0 \\ \text{vs} \quad H_A : p > p_0 \end{array}$$

Check the conditions and find SE using $p = p_0$

- ▶ $500 < 10\%$ of the population of Baltimore \Rightarrow independence
- ▶ Success-failure condition
 - ▶ $np_0 = 500 \times 0.5 = 250 \geq 10$
 - ▶ $n(1 - p_0) = 500 \times (1 - 0.5) = 250 \geq 10$

$$SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}} \approx \sqrt{\frac{p_0(1-p_0)}{n}} = \sqrt{\frac{0.5(1-0.5)}{500}} = 0.022$$


Hypothesis Tests

$$z = \frac{\text{point estimate } \hat{p} - \text{null value } p_0}{SE_{\hat{p}}} = \frac{0.52 - 0.50}{0.022} = 0.89$$

Hypothesis Tests

$$z = \frac{\text{point estimate } \hat{p} - \text{null value } p_0}{SE_{\hat{p}}} = \frac{0.52 - 0.50}{0.022} = 0.89$$


p-value is 0.1867. In the original %'age scale:



Hypothesis Tests

$$z = \frac{\text{point estimate } \hat{p} - \text{null value } p_0}{SE_{\hat{p}}} = \frac{0.52 - 0.50}{0.022} = 0.89$$

p-value is 0.1867. In the original %'age scale:



Hence we do **not** reject the null hypothesis, and we do not find convincing evidence to support the campaign manager's claim.

Next Time

Same as with the jump from

$$\mu \text{ to } \mu_1 - \mu_2$$

i.e. from one to two-sample tests for means, we make the jump from

$$p \text{ to } p_1 - p_2$$

i.e. from one to two-sample tests for proportions.