

## Lecture 7: Probability

### Chapter 2.x

1 / 16

## Outcomes

2 / 16

## Disjoint AKA Mutually Exclusive Outcomes

3 / 16

## Addition Rule of Probability

4 / 16

## General Addition Rule of Probability

5 / 16

## Sample Space and the Complement of Events

6 / 16

## Independence

Two processes are **independent** if knowing the outcome of one provides no useful information about the outcome of the other. Otherwise they are dependent.

Consider:

1. Die rolls
2. You get a movie recommendation from your friend Robin, but then their significant other Sam also recommends it.
3. You compare test scores from two Grade 9 students in the same class. Then same school. Then same school district. Then same city. Then same state.

## Independence

## Conditional Probability

9 / 16

### Example

Let's suppose I take a random sample of 100 Midd kids to study their smoking habits.

|        | Smoker | Not Smoker | Total |
|--------|--------|------------|-------|
| Male   | 19     | 41         | 60    |
| Female | 12     | 28         | 40    |
| Total  | 31     | 69         | 100   |

10 / 16

## Put It Together! Independence and Conditional Prob.

11 / 16

### Gambler's Fallacy: Roulette



You can bet on individual numbers, sets of numbers, or **red vs black**. Let's assume no 0 or 00, so that  $P(R) = P(B) = \frac{1}{2}$ .

12 / 16

## Gambler's Fallacy: Roulette

One of the biggest cons in casinos: **spin history boards**.



Let's ignore the numbers and just focus on what color occurred.

Note: the white values on the left are **black** spins.

13 / 16

## Gambler's Fallacy: Roulette

Let's say you look at the board and see that the last 4 spins were **red**. You will always hear people say "**Black is due!**"

Ex. on the 5th spin people think:

$$P(B_5 \mid R_1 \text{ and } R_2 \text{ and } R_3 \text{ and } R_4) >$$

$$P(R_5 \mid R_1 \text{ and } R_2 \text{ and } R_3 \text{ and } R_4)$$

14 / 16

## Gambler's Fallacy: Roulette

But assuming the wheel is not rigged, spins are independent i.e.  
 $P(A|B) = P(A)$ . So:


$$P(B_5|R_1 \text{ and } R_2 \text{ and } R_3 \text{ and } R_4) = P(B_5) = \frac{1}{2}$$

$$P(R_5|R_1 \text{ and } R_2 \text{ and } R_3 \text{ and } R_4) = P(R_5) = \frac{1}{2}$$

15 / 16

## Next Time

Discuss the Normal Distribution



16 / 16