

# Lecture 17: Paired Data and Difference of Two Means

Chapter 5.1-5.2

# Goals for Today

- ▶ Define statistical power
- ▶ Difference of Means
- ▶ Note on Practical vs Statistical Significance

# The 8 Types of Questions

Here are the 8 broad types of questions we can answer with statistical methods (confidence intervals and hypothesis tests) in this class:

# The 8 Types of Questions

Here are the 8 broad types of questions we can answer with statistical methods (confidence intervals and hypothesis tests) in this class:

1. What is the mean value  $\mu$ ?

# The 8 Types of Questions

Here are the 8 broad types of questions we can answer with statistical methods (confidence intervals and hypothesis tests) in this class:

1. What is the mean value  $\mu$ ?
2. Are the means of two groups  $\mu_1$  and  $\mu_2$  equal or not?

# The 8 Types of Questions

Here are the 8 broad types of questions we can answer with statistical methods (confidence intervals and hypothesis tests) in this class:

1. What is the mean value  $\mu$ ?
2. Are the means of two groups  $\mu_1$  and  $\mu_2$  equal or not?
3. What is the mean paired difference  $\mu_{diff}$ ?

# The 8 Types of Questions

Here are the 8 broad types of questions we can answer with statistical methods (confidence intervals and hypothesis tests) in this class:

1. What is the mean value  $\mu$ ?
2. Are the means of two groups  $\mu_1$  and  $\mu_2$  equal or not?
3. What is the mean paired difference  $\mu_{diff}$ ?
4. What is the proportion  $p$  of “successes”?

# The 8 Types of Questions

Here are the 8 broad types of questions we can answer with statistical methods (confidence intervals and hypothesis tests) in this class:

1. What is the mean value  $\mu$ ?
2. Are the means of two groups  $\mu_1$  and  $\mu_2$  equal or not?
3. What is the mean paired difference  $\mu_{diff}$ ?
4. What is the proportion  $p$  of “successes”?
5. Are the proportions of “successes” of two groups  $p_1$  and  $p_2$  equal or not?



# The 8 Types of Questions

Here are the 8 broad types of questions we can answer with statistical methods (confidence intervals and hypothesis tests) in this class:

1. What is the mean value  $\mu$ ?
2. Are the means of two groups  $\mu_1$  and  $\mu_2$  equal or not?
3. What is the mean paired difference  $\mu_{diff}$ ?
4. What is the proportion  $p$  of “successes”?
5. Are the proportions of “successes” of two groups  $p_1$  and  $p_2$  equal or not?
6. Are the means  $\mu_1, \dots, \mu_k$  of  $k$  groups **all** equal or not?

# The 8 Types of Questions

Here are the 8 broad types of questions we can answer with statistical methods (confidence intervals and hypothesis tests) in this class:

1. What is the mean value  $\mu$ ?
2. Are the means of two groups  $\mu_1$  and  $\mu_2$  equal or not?
3. What is the mean paired difference  $\mu_{diff}$ ?
4. What is the proportion  $p$  of “successes”?
5. Are the proportions of “successes” of two groups  $p_1$  and  $p_2$  equal or not?
6. Are the means  $\mu_1, \dots, \mu_k$  of  $k$  groups **all** equal or not?
7. Are we observing counts that we were expected?


# The 8 Types of Questions

Here are the 8 broad types of questions we can answer with statistical methods (confidence intervals and hypothesis tests) in this class:

1. What is the mean value  $\mu$ ?
2. Are the means of two groups  $\mu_1$  and  $\mu_2$  equal or not?
3. What is the mean paired difference  $\mu_{diff}$ ?
4. What is the proportion  $p$  of “successes”?
5. Are the proportions of “successes” of two groups  $p_1$  and  $p_2$  equal or not?
6. Are the means  $\mu_1, \dots, \mu_k$  of  $k$  groups **all** equal or not?
7. Are we observing counts that we were expected?
8. Are two categorical variables independent?

# Are the means of two groups $\mu_1$ and $\mu_2$ equal or not?

Example from Chapter 5.2: Did men (n=45) run faster than women (n=55)?



# Difference in Means

We are interested in the difference of two population means

$\mu_w - \mu_m$  where

# Difference in Means

We are interested in the difference of two population means

$\mu_w - \mu_m$  where

- ▶  $\mu_w$  is the mean time for women
- ▶  $\mu_m$  is the mean time for men

# Difference in Means

We are interested in the difference of two population means

$\mu_w - \mu_m$  where

- ▶  $\mu_w$  is the mean time for women
- ▶  $\mu_m$  is the mean time for men

The data:

	men	women
$\bar{x}$	87.65	102.13
$s$	12.5	15.2
$n$	45	55

# Difference of Means

We now recreate all the elements of Chapter 4 using this new population parameter  $\mu_w - \mu_m$ :



# Difference of Means

We now recreate all the elements of Chapter 4 using this new population parameter  $\mu_w - \mu_m$ :

1. Determine a point estimate of  $\mu_w - \mu_m$ .

# Difference of Means

We now recreate all the elements of Chapter 4 using this new population parameter  $\mu_w - \mu_m$ :

1. Determine a point estimate of  $\mu_w - \mu_m$ .
2. Show the normality of the sampling distribution: mean and SE

# Difference of Means

We now recreate all the elements of Chapter 4 using this new population parameter  $\mu_w - \mu_m$ :

1. Determine a point estimate of  $\mu_w - \mu_m$ .
2. Show the normality of the sampling distribution: mean and SE
3. Build a confidence interval

# Difference of Means

We now recreate all the elements of Chapter 4 using this new population parameter  $\mu_w - \mu_m$ :

1. Determine a point estimate of  $\mu_w - \mu_m$ .
2. Show the normality of the sampling distribution: mean and SE
3. Build a confidence interval
4. Conduct hypothesis tests

# Difference of Means

We now recreate all the elements of Chapter 4 using this new population parameter  $\mu_w - \mu_m$ :

1. Determine a point estimate of  $\mu_w - \mu_m$ .
2. Show the normality of the sampling distribution: mean and SE
3. Build a confidence interval
4. Conduct hypothesis tests

First, the point estimate for  $\mu_w - \mu_m$  is the sample difference of means

$$\bar{x}_w - \bar{x}_m = 102.13 - 87.65 = 14.48$$

# Normality of Sampling Distribution

If the sample means  $\bar{x}_1$  and  $\bar{x}_2$

# Normality of Sampling Distribution

If the sample means  $\bar{x}_1$  and  $\bar{x}_2$

- ▶ each meet the criteria for having nearly normal sampling distributions

# Normality of Sampling Distribution

If the sample means  $\bar{x}_1$  and  $\bar{x}_2$

- ▶ each meet the criteria for having nearly normal sampling distributions
- ▶ also the observations from the two samples are independent



# Normality of Sampling Distribution

If the sample means  $\bar{x}_1$  and  $\bar{x}_2$

- ▶ each meet the criteria for having nearly normal sampling distributions
- ▶ also the observations from the two samples are independent

then the difference in sample means  $\bar{x}_1 - \bar{x}_2$  will also have a nearly normal sampling distribution...

# Normality of Sampling Distribution

with

# Normality of Sampling Distribution

with

- ▶ mean  $\mu_1 - \mu_2$

# Normality of Sampling Distribution

with

- ▶ mean  $\mu_1 - \mu_2$
- ▶ estimated standard error

$$SE_{\bar{x}_1 - \bar{x}_2} = \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

# Normality of Sampling Distribution

with

- ▶ mean  $\mu_1 - \mu_2$
- ▶ estimated standard error

$$SE_{\bar{x}_1 - \bar{x}_2} = \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}$$

Note the different  $s^2$ 's and sample sizes.

# Normality of Sampling Distribution

We verify the conditions:

# Normality of Sampling Distribution

We verify the conditions:

- ▶ Because each sample consists of less than 10% of their respective populations (men: 45 of 7192 and women: 55 of 9732).

# Normality of Sampling Distribution

We verify the conditions:

- ▶ Because each sample consists of less than 10% of their respective populations (men: 45 of 7192 and women: 55 of 9732).
- ▶ The observations for both groups don't look too skewed.



# Normality of Sampling Distribution

We verify the conditions:

- ▶ Because each sample consists of less than 10% of their respective populations (men: 45 of 7192 and women: 55 of 9732).
- ▶ The observations for both groups don't look too skewed.
- ▶ Each sample has at least 30 observations (rule of thumb).

# Normality of Sampling Distribution

We verify the conditions:

- ▶ Because each sample consists of less than 10% of their respective populations (men: 45 of 7192 and women: 55 of 9732).
- ▶ The observations for both groups don't look too skewed.
- ▶ Each sample has at least 30 observations (rule of thumb).
- ▶ The samples are independent (not paired or linked in any way).

# Normality of Sampling Distribution

We verify the conditions:

- ▶ Because each sample consists of less than 10% of their respective populations (men: 45 of 7192 and women: 55 of 9732).
- ▶ The observations for both groups don't look too skewed.
- ▶ Each sample has at least 30 observations (rule of thumb).
- ▶ The samples are independent (not paired or linked in any way).

the sampling distribution is Normal with mean  $= \mu_w - \mu_m$  and

$$SE_{\bar{x}_w - \bar{x}_m} = \sqrt{\frac{15.2^2}{55} + \frac{12.5^2}{45}} = 2.77$$

# Confidence Interval

A 95% confidence interval for  $\mu_1 - \mu_2$  is

$$(\text{point estimate for } \mu_1 - \mu_2) \pm 1.96 \times SE$$

# Confidence Interval

A 95% confidence interval for  $\mu_1 - \mu_2$  is

$$\begin{aligned} &(\text{point estimate for } \mu_1 - \mu_2) \pm 1.96 \times SE \\ &(\bar{x}_1 - \bar{x}_2) \pm 1.96 \times SE_{\bar{x}_1 - \bar{x}_2} \end{aligned}$$

# Confidence Interval

A 95% confidence interval for  $\mu_1 - \mu_2$  is

$$\begin{aligned} &(\text{point estimate for } \mu_1 - \mu_2) \pm 1.96 \times SE \\ &(\bar{x}_1 - \bar{x}_2) \pm 1.96 \times SE_{\bar{x}_1 - \bar{x}_2} \end{aligned}$$

So for the Cherry Blossom Run data, a 95% CI for  $\mu_w - \mu_m$  is:

$$14.48 \pm 1.96 \times 2.77 = [9.05, 19.91]$$

## Next Time

- ▶ Hypothesis test for differences in means
- ▶ Paired differences
- ▶ One sample t-test