

Lecture 20: Single Proportion Test

Chapter 6.1

Quiz 8

Question 1: According to the article, why do scientists even bother with correlational/observational studies, when no notions of causality can be established?

Quiz 8

Question 1: According to the article, why do scientists even bother with correlational/observational studies, when no notions of causality can be established?

Answer: One reason is that correlational studies are excellent starting points for deciding which hypotheses to evaluate with the more rigorous randomized controlled experiment.

Quiz 8

Question 2: The article argues that various scientific disciplines should set professional labeling standards for material discussed in the media... Rank the four possible labels in order of how much credence the public should give them, from lowest to highest.

Quiz 8

Question 2: The article argues that various scientific disciplines should set professional labeling standards for material discussed in the media... Rank the four possible labels in order of how much credence the public should give them, from lowest to highest.

Answer:

3. preliminary result
1. large-scale observational study
4. large-sample randomized controlled test
2. well-established scientific law that we know how to apply in a wide range of conditions

Causality

What is causality? How do we establish it?

- ▶ http://nfs.unipv.it/nfs/minf/dispense/patgen/lectures/files/disease_causality.html
- ▶ <http://bayes.cs.ucla.edu/BOOK-2K/>

Question for Today

According to a poll done by the New York Times/CBS News in June 2012, only about 44% of the American public approves of the Supreme Court's performance.

Question for Today

According to a poll done by the New York Times/CBS News in June 2012, only about 44% of the American public approves of the Supreme Court's performance.

The sample proportion $\hat{p} = 0.44$ is **point estimate** of p : the true (population) proportion of the American public who approves.

Question for Today

According to a poll done by the New York Times/CBS News in June 2012, only about 44% of the American public approves of the Supreme Court's performance.

The sample proportion $\hat{p} = 0.44$ is **point estimate** of p : the true (population) proportion of the American public who approves.

Let's assume the sampling procedure was representative. What's the first thing to ask?

- ▶ What was n ?
- ▶ In other words, what is the **SE** of $\hat{p} = 44\%$?

Question for Today

Just like with \bar{x} , if we want to use the normal model to

- ▶ build confidence intervals via z^*
- ▶ conduct hypothesis tests via the normal tables

we need the **sampling distribution** of \hat{p} to be nearly normal.

Question for Today

Just like with \bar{x} , if we want to use the normal model to

- ▶ build confidence intervals via z^*
- ▶ conduct hypothesis tests via the normal tables

we need the **sampling distribution** of \hat{p} to be nearly normal.

This happens when the population distribution of 0's and 1's is not too strongly skewed. But as before, as the sample size $n \rightarrow \infty$, this is less of an issue by the CLT.

Conditions for Sampling Dist'n of \hat{p} Being Nearly Normal

The sampling distribution of the sample proportion \hat{p} based on a sample of size n is nearly normal when

Conditions for Sampling Dist'n of \hat{p} Being Nearly Normal

The sampling distribution of the **sample proportion** \hat{p} based on a sample of size n is nearly normal when

- ▶ The observations are independent: the 10% rule

Conditions for Sampling Dist'n of \hat{p} Being Nearly Normal

The sampling distribution of the **sample proportion** \hat{p} based on a sample of size n is nearly normal when

- ▶ The observations are independent: the 10% rule
- ▶ We expect to see at least 10 successes and 10 failures in our sample. This is called the **success-failure condition**: that both
 - ▶ $np \geq 10$
 - ▶ $n(1 - p) \geq 10$

Conditions for Sampling Dist'n of \hat{p} Being Nearly Normal

If conditions are met, then the sampling distribution of \hat{p} is nearly normal with

- ▶ mean p (the true population proportion)
- ▶ standard error

$$SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$$

Conditions for Sampling Dist'n of \hat{p} Being Nearly Normal

If conditions are met, then the sampling distribution of \hat{p} is nearly normal with

- ▶ mean p (the true population proportion)
- ▶ standard error

$$SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$$

Note the similarity of the previous formula for the sample mean \bar{x} :

$$SE_{\bar{x}} = \frac{\sigma}{\sqrt{n}} = \sqrt{\frac{\sigma^2}{n}}$$

Standard Error of \hat{p}

But we **don't know** what p is. So what p do we use

- ▶ to check the conditions?
- ▶ for the $SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$?

How do we verify success/failure conditions?

Standard Error of \hat{p}

But we **don't know** what p is. So what p do we use

- ▶ to check the conditions?
- ▶ for the $SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$?

How do we verify success/failure conditions?

- ▶ Confidence intervals: plug in the **point estimate** \hat{p} of p

Standard Error of \hat{p}

But we **don't know** what p is. So what p do we use

- ▶ to check the conditions?
- ▶ for the $SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$?

How do we verify success/failure conditions?

- ▶ Confidence intervals: plug in the **point estimate** \hat{p} of p
- ▶ Hypothesis tests: plug in the **null value** p_0 from $H_0 : p = p_0$

Confidence Intervals

Going back to the poll: $\hat{p} = 0.44$ based on $n = 976$. What is a 95% confidence interval?

Confidence Intervals

Going back to the poll: $\hat{p} = 0.44$ based on $n = 976$. What is a 95% confidence interval?

Check the conditions and find SE using $p = \hat{p}$

Confidence Intervals

Going back to the poll: $\hat{p} = 0.44$ based on $n = 976$. What is a 95% confidence interval?

Check the conditions and find SE using $p = \hat{p}$

- ▶ $976 < 313 \text{ million} \Rightarrow \text{independence}$

Confidence Intervals

Going back to the poll: $\hat{p} = 0.44$ based on $n = 976$. What is a 95% confidence interval?

Check the conditions and find SE using $p = \hat{p}$

- ▶ $976 < 313 \text{ million} \Rightarrow$ independence
- ▶ Defining a success as a person approving of the job done by the Supreme Court:

Confidence Intervals

Going back to the poll: $\hat{p} = 0.44$ based on $n = 976$. What is a 95% confidence interval?

Check the conditions and find SE using $p = \hat{p}$

- ▶ $976 < 313 \text{ million} \Rightarrow$ independence
- ▶ Defining a success as a person approving of the job done by the Supreme Court:
 - ▶ $976 \times \hat{p} = 976 \times .44 = 429 \text{ successes} \geq 10$
 - ▶ $976 \times (1 - \hat{p}) = 976 \times .56 = 547 \text{ failures} \geq 10$

Confidence Intervals

Going back to the poll: $\hat{p} = 0.44$ based on $n = 976$. What is a 95% confidence interval?

Check the conditions and find SE using $p = \hat{p}$

- ▶ $976 < 313 \text{ million} \Rightarrow$ independence
- ▶ Defining a success as a person approving of the job done by the Supreme Court:
 - ▶ $976 \times \hat{p} = 976 \times .44 = 429 \text{ successes} \geq 10$
 - ▶ $976 \times (1 - \hat{p}) = 976 \times .56 = 547 \text{ failures} \geq 10$

$$SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}} \approx \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} = \sqrt{\frac{0.44(1-0.44)}{976}} = 0.016$$

Confidence Intervals

A 95% confidence interval using the normal model has $z^* = 1.96$,
thus:

$$\text{point estimate} \pm 1.96 \times SE$$

Confidence Intervals

A 95% confidence interval using the normal model has $z^* = 1.96$, thus:

$$\text{point estimate} \pm 1.96 \times SE$$

In our case

$$\hat{p} \pm 1.96 \times SE_{\hat{p}} = 0.44 \pm 1.96 \times 0.016 = (0.409, 0.471)$$

Hypothesis Tests

Thomas Carcetti is running for mayor of Baltimore. His campaign manager **claims** he has more than 50% support of the electorate.

Hypothesis Tests

Thomas Carcetti is running for mayor of Baltimore. His campaign manager **claims** he has more than 50% support of the electorate.

The Baltimore Sun collects a random sample of $n = 500$ likely voters and finds that 52% support him. Does this provide convincing evidence for the claim of Carcetti's manager at the 5% significance level?

Hypothesis Tests

The hypothesis test is, with the null value $p_0 = 0.5$

$$\begin{array}{l} H_0 : p = p_0 \\ \text{vs} \quad H_A : p > p_0 \end{array}$$

Hypothesis Tests

The hypothesis test is, with the null value $p_0 = 0.5$

$$\begin{array}{l} H_0 : p = p_0 \\ \text{vs} \quad H_A : p > p_0 \end{array}$$

Check the conditions and find SE using $p = p_0$

Hypothesis Tests

The hypothesis test is, with the null value $p_0 = 0.5$

$$\begin{array}{l} H_0 : p = p_0 \\ \text{vs} \quad H_A : p > p_0 \end{array}$$

Check the conditions and find SE using $p = p_0$

- ▶ $500 < 10\%$ of the population of Baltimore \Rightarrow independence

Hypothesis Tests

The hypothesis test is, with the null value $p_0 = 0.5$

$$\begin{array}{l} H_0 : p = p_0 \\ \text{vs} \quad H_A : p > p_0 \end{array}$$

Check the conditions and find SE using $p = p_0$

- ▶ $500 < 10\%$ of the population of Baltimore \Rightarrow independence
- ▶ Success-failure condition
 - ▶ $np_0 = 500 \times 0.5 = 250 \geq 10$
 - ▶ $n(1 - p_0) = 500 \times (1 - 0.5) = 250 \geq 10$

Hypothesis Tests

The hypothesis test is, with the null value $p_0 = 0.5$

$$\begin{array}{l} H_0 : p = p_0 \\ \text{vs} \quad H_A : p > p_0 \end{array}$$

Check the conditions and find SE using $p = p_0$

- ▶ $500 < 10\%$ of the population of Baltimore \Rightarrow independence
- ▶ Success-failure condition
 - ▶ $np_0 = 500 \times 0.5 = 250 \geq 10$
 - ▶ $n(1 - p_0) = 500 \times (1 - 0.5) = 250 \geq 10$

$$SE_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}} \approx \sqrt{\frac{p_0(1-p_0)}{n}} = \sqrt{\frac{0.5(1-0.5)}{500}} = 0.022$$


Hypothesis Tests

$$z = \frac{\text{point estimate } \hat{p} - \text{null value } p_0}{SE_{\hat{p}}} = \frac{0.52 - 0.50}{0.022} = 0.89$$

Hypothesis Tests

$$z = \frac{\text{point estimate } \hat{p} - \text{null value } p_0}{SE_{\hat{p}}} = \frac{0.52 - 0.50}{0.022} = 0.89$$


p-value is 0.1867.



Hypothesis Tests

$$z = \frac{\text{point estimate } \hat{p} - \text{null value } p_0}{SE_{\hat{p}}} = \frac{0.52 - 0.50}{0.022} = 0.89$$

p-value is 0.1867.



Hence we do **not** reject the null hypothesis, and we do not find convincing evidence to support the campaign manager's claim.

Next Time

Same as with the jump from

$$\mu \text{ to } \mu_1 - \mu_2$$

i.e. from one to two-sample tests for means, we make the jump from

$$p \text{ to } p_1 - p_2$$

i.e. from one to two-sample tests for proportions.