

Lecture 16: Sample Size and Power

Chapter 4.6

Last Time: Reddie Sleep Example

Tested number of hours of sleep:

- ▶ $H_0 : \mu = 7$

- ▶ $H_A : \mu > 7$

Two-Sided Alternative Hypothesis

Say instead we had a two-sided alternative hypothesis:

- ▶ $H_0 : \mu = 7$

- ▶ $H_A : \mu \neq 7$


Two-Sided Alternative Hypothesis

Say instead we had a **two-sided alternative hypothesis**:

► $H_0 : \mu = 7$

► $H_A : \mu \neq 7$

The the p-value would be double: $2 \times 0.007 = 0.014$. Picture:



Pre-specifying α

Say Dr. Q is conducting a hypothesis tests. They start with $\alpha = 0.05$.

Pre-specifying α

Say Dr. Q is conducting a hypothesis tests. They start with $\alpha = 0.05$.

They conduct the test and get $p\text{-value} = 0.09$. They then declare “having used an $\alpha = 0.10$, we reject the null hypothesis and declare our results to be significant.”

Pre-specifying α

Say Dr. Q is conducting a hypothesis tests. They start with $\alpha = 0.05$.

They conduct the test and get $p\text{-value} = 0.09$. They then declare “having used an $\alpha = 0.10$, we reject the null hypothesis and declare our results to be significant.”

What's not honest about this approach?

Pre-specifying α

Say Dr. Q is conducting a hypothesis tests. They start with $\alpha = 0.05$.

They conduct the test and get $p\text{-value} = 0.09$. They then declare “having used an $\alpha = 0.10$, we reject the null hypothesis and declare our results to be significant.”

What's not honest about this approach?

Ronald Fisher, the creator of p-values, never intended for them to be used this way: <http://en.wikipedia.org/wiki/P-value#Criticisms>

Goals for Today

- ▶ More in depth discussion of
 - ▶ 10% sampling rule
 - ▶ Skew condition to check to use the normal model
- ▶ How big a sample size do I need?
- ▶ Statistical power
- ▶ Statistical vs practical significance

10% Sampling Rule

Question: Why do we set $n \leq 10\%$ of the population size N ?

10% Sampling Rule

Question: Why do we set $n \leq 10\%$ of the population size N ?

Intuition: Shouldn't we always sample as many people as we can?

10% Sampling Rule

Question: Why do we set $n \leq 10\%$ of the population size N ?

Intuition: Shouldn't we always sample as many people as we can?

Answer: Yes, if we only care about the mean. If we also care about the SE, then we need to be careful.

10% Sampling Rule

Question: Why do we set $n \leq 10\%$ of the population size N ?

Intuition: Shouldn't we always sample as many people as we can?

Answer: Yes, if we only care about the mean. If we also care about the SE, then we need to be careful.

Issue: Sampling without vs with replacement.

Finite Population Correction

Finite Population Correction

Conceptual and Mathematical Notions of Sampling

Conceptual: If we sample everybody, we know the true μ .

Conceptual and Mathematical Notions of Sampling

Conceptual: If we sample everybody, we know the true μ .

and

Mathematical: If $n = N$ then $FPC = \sqrt{\frac{N-n}{N-1}} = 0$ then
 $SE = \frac{\sigma}{\sqrt{n}} \times FPC = 0$

Conceptual and Mathematical Notions of Sampling

Conceptual: If we sample everybody, we know the true μ .

and

Mathematical: If $n = N$ then $FPC = \sqrt{\frac{N-n}{N-1}} = 0$ then
 $SE = \frac{\sigma}{\sqrt{n}} \times FPC = 0$

i.e.

- ▶ the sampling distribution is just one point: the true μ .
- ▶ if we repeat this procedure many times, we get the same value each time: 0 variability.

Sampling and the SE

Question: Why do we care that our SE is correct?

Sampling and the SE

Question: Why do we care that our SE is correct?

Answer: If not

- ▶ the SE in confidence intervals is off
- ▶ the z-scores of \bar{x} have the wrong denominator

Skew Condition to Check to Use Normal Model

Throughout the book, they talk about the condition for \bar{x} being nearly normal and using s in place of σ in $SE = \frac{\sigma}{\sqrt{n}}$:

Skew Condition to Check to Use Normal Model

Throughout the book, they talk about the condition for \bar{x} being nearly normal and using s in place of σ in $SE = \frac{\sigma}{\sqrt{n}}$:

- ▶ On page 164: the population distribution is not strongly skewed

Skew Condition to Check to Use Normal Model

Throughout the book, they talk about the condition for \bar{x} being nearly normal and using s in place of σ in $SE = \frac{\sigma}{\sqrt{n}}$:

- ▶ On page 164: the population distribution is not strongly skewed
- ▶ On page 167: the data are not strongly skewed

Skew Condition to Check to Use Normal Model

Throughout the book, they talk about the condition for \bar{x} being nearly normal and using s in place of σ in $SE = \frac{\sigma}{\sqrt{n}}$:

- ▶ On page 164: the population distribution is not strongly skewed
- ▶ On page 167: the data are not strongly skewed
- ▶ On page 168: the distribution of sample observations is not strongly skewed

Skew Condition to Check to Use Normal Model

Throughout the book, they talk about the condition for \bar{x} being nearly normal and using s in place of σ in $SE = \frac{\sigma}{\sqrt{n}}$:

- ▶ On page 164: the population distribution is not strongly skewed
- ▶ On page 167: the data are not strongly skewed
- ▶ On page 168: the distribution of sample observations is not strongly skewed
- ▶ On page 185: the population data are not strongly skewed

Skew Condition to Check to Use Normal Model

However, they all mean the same thing:

Skew Condition to Check to Use Normal Model

However, they all mean the same thing:

1. The **true population** distribution from which you are drawing your sample observations/data x_1, \dots, x_n is not too skewed.


Skew Condition to Check to Use Normal Model

However, they all mean the same thing:

1. The **true population** distribution from which you are drawing your sample observations/data x_1, \dots, x_n is not too skewed.
2. The histogram (visual estimate) of the sample observations/data x_1, \dots, x_n is not too skewed.


Sample Size: Thought Experiment

Say you have two distributions with $\mu = 15$ but different σ .



Sample Size: Thought Experiment

Say you have two distributions with $\mu = 15$ but different σ .



Which of the two distributions do you think will require a bigger n to estimate μ “well”?

Margin of Error

Recall our formula for a 95% confidence interval:

$$\left[\bar{x} - 1.96 \frac{s}{\sqrt{n}}, \bar{x} + 1.96 \frac{s}{\sqrt{n}} \right]$$

Margin of Error

Recall our formula for a 95% confidence interval:

$$\left[\bar{x} - 1.96 \frac{s}{\sqrt{n}}, \bar{x} + 1.96 \frac{s}{\sqrt{n}} \right]$$

The **margin of error** is half the width of the CI.

Margin of Error

Recall our formula for a 95% confidence interval:

$$\left[\bar{x} - 1.96 \frac{s}{\sqrt{n}}, \bar{x} + 1.96 \frac{s}{\sqrt{n}} \right]$$

The **margin of error** is half the width of the CI.

Say we knew the **true** standard deviation σ , then

$$\text{Margin of Error} = 1.96 \frac{\sigma}{\sqrt{n}}$$

Identify n for a Desired Margin of Error

Identify n for a Desired Margin of Error

So

Identify n for a Desired Margin of Error

So

- ▶ As σ goes up, you need more n

Identify n for a Desired Margin of Error

So

- ▶ As σ goes up, you need more n
- ▶ As z^* goes up, i.e. higher confidence level, you need more n


Identify n for a Desired Margin of Error

So

- ▶ As σ goes up, you need more n
- ▶ As z^* goes up, i.e. higher confidence level, you need more n
- ▶ As the desired margin of error goes down, you need more n

Back to Thought Experiment

For the same desired maximal margin of error m and same confidence level, we need a larger n to estimate the mean of the blue curve:



Type II Error Rate and Power

Type II Error Rate and Power