

Lecture 19: ANOVA Part I

Chapter 5.5

Previously: Conditions for Using t Distribution

We use the t distribution when you have

Previously: Conditions for Using t Distribution

We use the t distribution when you have

- ▶ n is small.

Previously: Conditions for Using t Distribution

We use the t distribution when you have

- ▶ n is small.
- ▶ Independence: $n \leq 10\%$ rule

Previously: Conditions for Using t Distribution

We use the t distribution when you have


- ▶ n is small.
- ▶ Independence: $n \leq 10\%$ rule
- ▶ Observations come from a nearly normal distribution:
 - ▶ Look at a histogram of the data (difficult when n is small)
 - ▶ Consider whether any previous experiences alert us that the data may be normal

Analysis of Variance (ANOVA)

A farmer has the choice of four tomato fertilizers and wants to compare their performance in terms of crop yield.


Analysis of Variance (ANOVA)

A farmer has the choice of four tomato fertilizers and wants to compare their performance in terms of crop yield.



Analysis of Variance (ANOVA)

A farmer has the choice of four tomato fertilizers and wants to compare their performance in terms of crop yield.



Analysis of Variance (ANOVA)

We have $k = 4$ groups AKA **levels of a factor**: the 4 types of fertilizer.

Analysis of Variance (ANOVA)

We have $k = 4$ groups AKA **levels of a factor**: the 4 types of fertilizer.

- ▶ n_i plants assigned to each of the $k = 4$ fertilizers:

n_1	n_2	n_3	n_4	total n
3	7	4	6	20

Analysis of Variance (ANOVA)

We have $k = 4$ groups AKA **levels of a factor**: the 4 types of fertilizer.


- ▶ n_i plants assigned to each of the $k = 4$ fertilizers:

n_1	n_2	n_3	n_4	total n
3	7	4	6	20

- ▶ Count the number of tomatoes on each plant


Tomato Fertilizer

We observe the following, where each point is one tomato plant.




Tomato Fertilizer

We observe the following, where each point is one tomato plant.
Plot the sample mean of each level.



Tomato Fertilizer

We observe the following, where each point is one tomato plant.
Plot the sample mean of each level. **Question:** are the mean tomato yields different?



Analysis of Variance

Say we have k groups and want to compare the k means:

$$\mu_1, \mu_2, \dots, \mu_k$$

Analysis of Variance

Say we have k groups and want to compare the k means:

$$\mu_1, \mu_2, \dots, \mu_k$$

We could do $\binom{k}{2}$ individual two-sample tests.

Analysis of Variance

Say we have k groups and want to compare the k means:

$$\mu_1, \mu_2, \dots, \mu_k$$


We could do $\binom{k}{2}$ individual two-sample tests.

Ex. for groups 1 & 2:

$$\begin{array}{ll} H_0 : & \mu_1 = \mu_2 \\ \text{vs. } H_a : & \mu_1 \neq \mu_2 \end{array}$$


Tomato Fertilizer Example

Numerator: the **between-group variation** refers to the variability **between** the levels (the 4 horizontal lines):





Tomato Fertilizer Example

Denominator: the **within-group variation** refers to the variability **within** each level (the 4 vertical arrows):



Tomato Fertilizer Example

Now compare the following two plots. Which has “more different” means?




Tomato Fertilizer Example

F Distributions


F Distributions

For $df_1 = 4$ and $df_2 = 6$, the F distribution looks like:



F Distributions

p -values are computed where “more extreme” means larger. Say the $F = 3$, the p -value is the area to the right of 3 and is computed in R: `pf(3,df1=4,df2=6,lower.tail=FALSE)`



Conducting An F -Test

The results are typically summarized in an ANOVA table:

Source of Variation	df	SS	MS	F	p -value
Between groups	$k - 1$	$SSTr$	$MSTr = \frac{SSTr}{k-1}$	$\frac{MSTr}{MSE}$	p
Within groups	$n - k$	SSE	$MSE = \frac{SSE}{n-k}$		
Total	$n - 1$	SST			

Conditions

1. The observations have to be **independent**. 10% rule.

Conditions

1. The observations have to be **independent**. 10% rule.
2. Trade off of n and **normality** of observations **within each group**.

Conditions

1. The observations have to be **independent**. 10% rule.
2. Trade off of n and **normality** of observations **within each group**.
3. Each of the groups has **constant variance** $\sigma_1^2 = \dots = \sigma_k^2 = \sigma^2$.
Check via:
 - ▶ boxplots
 - ▶ comparing the sample standard deviations s_1, \dots, s_k

Discussion of Quiz

Question 1: Why did $\frac{1}{20}$ studies yield a positive/significant result i.e. that there is a link between jelly beans and acne?

Discussion of Quiz


Question 1: Why did $\frac{1}{20}$ studies yield a positive/significant result i.e. that there is a link between jelly beans and acne?

Not that the p-value is 0.05, rather that $\alpha = 0.05$:

- ▶ significance level AKA
- ▶ type I error rate AKA
- ▶ false positive rate

i.e. we expect 1 out of 20 results to be significant even if there is no effect.

Publication Bias



Publication Bias

Publication bias: people only highlight significant/positive results.

Publication Bias

Publication bias: people only highlight significant/positive results.
From Wikipedia: “Publication bias occurs when the publication of research results **depends on their nature and direction.**”

Publication Bias

Publication bias: people only highlight significant/positive results.
From Wikipedia: “Publication bias occurs when the publication of research results **depends on their nature and direction.**”

To counter this, some prominent medical journals including

- ▶ New England Journal of Medicine
- ▶ The Lancet
- ▶ Journal of the American Medical Association

Publication Bias

Publication bias: people only highlight significant/positive results. From Wikipedia: “Publication bias occurs when the publication of research results **depends on their nature and direction.**”

To counter this, some prominent medical journals including

- ▶ New England Journal of Medicine
- ▶ The Lancet
- ▶ Journal of the American Medical Association

require registration of a trial **before** it starts so that unfavorable results are not withheld from publication.

Publication Bias

Publication bias: people only highlight significant/positive results.
From Wikipedia: “Publication bias occurs when the publication of research results **depends on their nature and direction.**”

To counter this, some prominent medical journals including

- ▶ New England Journal of Medicine
- ▶ The Lancet
- ▶ Journal of the American Medical Association

require registration of a trial **before** it starts so that unfavorable results are not withheld from publication.

Journal of **Negative Results:** <http://www.jnrbm.com/>

Publication Bias



REPRINTED WITH SPECIAL PERMISSION OF NORTH AMERICAN SYNDICATE

From: Sterne JA, Davey Smith G (2001) Sifting the evidence - What's wrong with significance tests. BMJ 322: 226231.

What α to Use?

Should I use $\alpha = 0.05$ as my significance level? Before using it, put some thought into the balance between:

What α to Use?

Should I use $\alpha = 0.05$ as my significance level? Before using it, put some thought into the balance between:

- ▶ **Type I errors.** Setting a smaller α yields a more **conservative** procedure: all things being equal, you will reject H_0 less often.

What α to Use?

Should I use $\alpha = 0.05$ as my significance level? Before using it, put some thought into the balance between:

- ▶ **Type I errors.** Setting a smaller α yields a more **conservative** procedure: all things being equal, you will reject H_0 less often.
- ▶ **Type II errors.** Setting a bigger α yields a more **liberal** procedure: all things being equal, you will reject H_0 more often.

Multiple Testing

A related issue is the statistical concept of **multiple testing**.

Multiple Testing

A related issue is the statistical concept of [multiple testing](#).

Say we are conducting many experiments, and H_0 is true for all of them.

Multiple Testing

A related issue is the statistical concept of **multiple testing**.

Say we are conducting many experiments, and H_0 is true for all of them.

If you repeat experiments many times, you're bound to get a significant result eventually just by **chance alone**.

Multiple Testing

What do people do?

Multiple Testing

What do people do? Make the α stricter! i.e.

- ▶ make the α smaller

Multiple Testing

What do people do? Make the α stricter! i.e.

- ▶ make the α smaller
- ▶ i.e. less chance the p-value is smaller than α

Multiple Testing

What do people do? Make the α stricter! i.e.

- ▶ make the α smaller
- ▶ i.e. less chance the p-value is smaller than α
- ▶ i.e. less chance of incorrectly rejecting H_0 when it is true

Multiple Testing

What do people do? Make the α stricter! i.e.

- ▶ make the α smaller
- ▶ i.e. less chance the p-value is smaller than α
- ▶ i.e. less chance of incorrectly rejecting H_0 when it is true

Use the **Bonferroni correction** to α : If you are conducting n tests, use $\alpha^* = \frac{\alpha}{n}$. You'll study its properties in HW8.