

## Lecture 15: Hypothesis Testing Part II

### Chapter 4.3

1 / 16

### Goals for Today

- ▶ Define significance level
- ▶ Tie-in p-Values with sampling distributions
- ▶ Example

2 / 16

## Type I Errors: US Criminal Justice System

Defendants must be “guilty beyond a reasonable doubt”: better to let a guilty person go free, than put an innocent person in jail.

- ▶  $H_0$ : the defendant is innocent
- ▶  $H_A$ : the defendant is guilty

thus “rejecting  $H_0$ ” is a guilty verdict  $\Rightarrow$  putting them in jail

In this case:

- ▶ Type I error = jailing an innocent person (worse)
- ▶ Type II error = letting a guilty person go free.

3 / 16

## Type II Errors: Airport Screening

An example of where Type II errors are more serious: [airport screening](#).

$H_0$  : passenger X does not have a weapon

$H_A$  : passenger X has a weapon

Failing to reject  $H_0$  when  $H_A$  is true is not “patting down” passenger X when they have a weapon.

Hence the long lines at airport security.

4 / 16

## Significance Level

5 / 16

## Thought experiment: Coin Flips

Say you flip a coin you think is fair 1000 times. Say you observe

- ▶ 501 heads? Do you think the coin is biased?
- ▶ 525 heads? Do you think the coin is biased?
- ▶ 900 heads? Do you think the coin is biased?

6 / 16

## Thought experiment: p-Values

7 / 16

## p-Values


8 / 16

## Thought experiment: Coin Flips

9 / 16

## Sampling Distribution of $\hat{p}$


Under  $H_0$  the sampling distribution of  $\hat{p}$  when  $n = 1000$  is:



10 / 16

Say we observe...


$$\hat{p} = \frac{501}{1000}$$



11 / 16

Say we observe...


$$\hat{p} = \frac{525}{1000}$$



12 / 16

Say we observe...

$$\hat{p} = \frac{900}{1000}$$




13 / 16

## Example about Sleep Habits

A US-wide poll found that college students sleep about 7 hours a night. You suspect that Midd Kids sleep more and investigate this claim at a pre-specified  $\alpha = 0.01$  level.

You sample  $n = 110$  Midd Kids and find that  $\bar{x} = 7.42$  and  $s = 1.75$  with a histogram that looks like:



14 / 16

## Example about Sleep Habits

15 / 16

## Example about Sleep Habits

**Conclusion:** we reject at the  $\alpha = 0.01$  significance level the hypothesis that the average # of hours Midd Kids sleep is 7, in favor of the hypothesis that they sleep more.

**Correct interpretation of the p-value:** If the null hypothesis is true ( $\mu = 7$ ), the probability of observing a sample mean  $\bar{x} = 7.42$  or greater is 0.007 (small).

**Incorrect interpretation of the p-value:** The probability that the null hypothesis ( $\mu = 7$ ) is true is 0.007.

16 / 16