

# Lecture 7: Probability

Chapter 2.x

# Outcomes

Probability forms the theoretical backbone of statistics. We use probability to characterize randomness.

We often frame probability in terms of a **random process** giving rise to an **outcome**.

# Outcomes

Probability forms the theoretical backbone of statistics. We use probability to characterize randomness.

We often frame probability in terms of a **random process** giving rise to an **outcome**.

Typical examples

- ▶ Die roll: 6 outcomes
- ▶ Coin Flip: 2 outcomes

# Disjoint AKA Mutually Exclusive Outcomes

# Addition Rule of Probability

# General Addition Rule of Probability

# General Addition Rule of Probability

# General Addition Rule of Probability



# Sample Space and the Complement of Events

# Independence

Two processes are **independent** if knowing the outcome of one provides no useful information about the outcome of the other. Otherwise they are dependent.

Consider:

# Independence

Two processes are **independent** if knowing the outcome of one provides no useful information about the outcome of the other. Otherwise they are dependent.

Consider:

1. Die rolls

# Independence

Two processes are **independent** if knowing the outcome of one provides no useful information about the outcome of the other. Otherwise they are dependent.

Consider:

1. Die rolls
2. You get a movie recommendation from your friend Robin, but then their significant other Sam also recommends it.

# Independence

Two processes are **independent** if knowing the outcome of one provides no useful information about the outcome of the other. Otherwise they are dependent.

Consider:

1. Die rolls
2. You get a movie recommendation from your friend Robin, but then their significant other Sam also recommends it.
3. You compare test scores from two Grade 9 students in the same class. Then same school. Then same school district. Then same city. Then same state.

# Independence

# Conditional Probability

## Example

Let's suppose I take a random sample of 100 Reed students to study their smoking habits.

	Smoker	Not Smoker	Total
Male	19	41	60
Female	12	28	40
Total	31	69	100



## Example

Let's suppose I take a random sample of 100 Reed students to study their smoking habits.

	Smoker	Not Smoker	Total
Male	19	41	60
Female	12	28	40
Total	31	69	100

- What is the probability of a randomly selected male smoking?

$$P(S|M) = \frac{P(S \text{ and } M)}{P(M)} = \frac{19/100}{60/100} = \frac{19}{60}$$

## Example

Let's suppose I take a random sample of 100 Reed students to study their smoking habits.

	Smoker	Not Smoker	Total
Male	19	41	60
Female	12	28	40
Total	31	69	100

- ▶ What is the probability of a randomly selected male smoking?

$$P(S|M) = \frac{P(S \text{ and } M)}{P(M)} = \frac{19/100}{60/100} = \frac{19}{60}$$

- ▶ What is the probability that a randomly selected smoker is female?

$$P(F|S) = \frac{P(F \text{ and } S)}{P(S)} = \frac{12/100}{31/100} = \frac{12}{31}$$

## Put It Together! Independence and Conditional Prob.

## Gambler's Fallacy: Roulette



You can bet on individual numbers, sets of numbers, or **red vs black**. Let's assume no 0 or 00, so that  $P(R) = P(B) = \frac{1}{2}$ .

# Gambler's Fallacy: Roulette

One of the biggest cons in casinos: **spin history boards**.



Let's ignore the numbers and just focus on what color occurred.  
Note: the white values on the left are **black** spins.

# Gambler's Fallacy: Roulette

Let's say you look at the board and see that the last 4 spins were red.

## Gambler's Fallacy: Roulette

Let's say you look at the board and see that the last 4 spins were red.

You will always hear people say "Black is due!"

## Gambler's Fallacy: Roulette

Let's say you look at the board and see that the last 4 spins were red.

You will always hear people say "Black is due!"

Ex. on the 5th spin people think:

$$\begin{aligned} P(B_5 \mid R_1 \text{ and } R_2 \text{ and } R_3 \text{ and } R_4) &> \\ P(R_5 \mid R_1 \text{ and } R_2 \text{ and } R_3 \text{ and } R_4) \end{aligned}$$



## Gambler's Fallacy: Roulette

But assuming the wheel is not rigged, spins are independent i.e.  
 $P(A|B) = P(A)$ . So:

## Gambler's Fallacy: Roulette

But assuming the wheel is not rigged, spins are independent i.e.  
 $P(A|B) = P(A)$ . So:

$$P(B_5|R_1 \text{ and } R_2 \text{ and } R_3 \text{ and } R_4) = P(B_5) = \frac{1}{2}$$
$$P(R_5|R_1 \text{ and } R_2 \text{ and } R_3 \text{ and } R_4) = P(R_5) = \frac{1}{2}$$

## Next Time

Discuss the Normal Distribution

