


From: Central Precocious Puberty Support [mailto:Info@csm.LP.abbvie.com]

To: John Doe <John.doe@example.com>

Subject: Central Precocious Puberty (CPP) treatment tips

Injection fears are normal—we're here to help.
[View in browser](#)



Dear [recipient name],

Thank you for signing up to receive updates from LUPRON DEPOT-PED while your child is on treatment.

By now, your child is ready to begin or has started their treatment regimen with LUPRON DEPOT-PED. Talking with them about their injection fears is important to their success on treatment. There are a few things to know about what to expect at the doctor's office.

MANAGING TREATMENT EXPECTATIONS WITH YOUR CHILD CAN PAVE THE WAY TO A SUCCESSFUL TREATMENT JOURNEY

Here are some tips to help prepare your child for their upcoming visits to the doctor and what you can expect when you get there.

BEFORE

- Help your child relax on the way to the doctor by playing a game or making up a song
- In the waiting room, having cards, a book, or something that can help distract your child can help ease some of the anxiety

DURING

- In the exam room, the nurse or medical assistant will prepare your child for the injection
- The nurse may do an activity to help your child relax their muscles
- The injection site can vary between the upper arm, buttocks, or thigh to make the experience easier for your child

AFTER

- After the injection, some pain and irritation at the site is expected. If more severe symptoms occur or develop, contact your child's doctor right away

HELPFUL WAYS TO DISTRACT YOUR CHILD IF THEY ARE FEELING ANXIOUS

If your child is scared of injections, there are some distraction techniques that can help. Here are just a few to get you started. Be sure to get creative! Remember, you know your child best, so do what works for them.

- Playing with toys
- Listening to music
- Reading books
- Watching videos

How you act can also play a big role in helping reduce anxiety. If you're anxious, your child may be anxious. If you're calm, your child will be calm.

Remember that with LUPRON DEPOT-PED, Support PLUS provides benefits, cost-saving information, and nurse support. [Learn more about Support PLUS.](#)

LEARN MORE ABOUT HOW YOU CAN ASSIST YOUR CHILD IN UNDERSTANDING THEIR TREATMENT PROCESS AND WHAT TO EXPECT AT THE DOCTOR'S OFFICE

[Learn more about starting therapy](#)

Use for LUPRON DEPOT-PED®(leuprolide acetate for depot suspension)

LUPRON DEPOT-PED 7.5 mg, 11.25 mg, and 15 mg for 1-month and 11.25 mg and 30 mg for 3-month administration are prescribed for the treatment of children with central precocious puberty (CPP).

It is not known if LUPRON DEPOT-PED is safe and effective in children under 2 years of age.

Important Safety Information for LUPRON DEPOT-PED

What is the most important information I should know about LUPRON DEPOT-PED?

- During the first 2 to 4 weeks of treatment, LUPRON DEPOT-PED can cause an increase in some hormones. During this time, you may notice more signs of puberty in your child, including vaginal bleeding. **Call your child's doctor if these signs continue after the second month of treatment with LUPRON DEPOT-PED.**
- Some people taking gonadotropin releasing hormone (GnRH) agonists like LUPRON DEPOT-PED have had new or worsened mental (psychiatric) problems. Mental (psychiatric) problems may include emotional symptoms such as:
 - Crying
 - Irritability
 - Restlessness (impatience)
 - Anger
 - Acting aggressive

Call your child's doctor right away if your child has any new or worsening mental symptoms or problems while taking LUPRON DEPOT-PED.

- Some people taking GnRH agonists like LUPRON DEPOT-PED have had seizures. The risk of seizures may be higher in people who:
 - Have a history of seizures
 - Have a history of epilepsy
 - Have a history of brain or brain vessel (cerebrovascular) problems or tumors
 - Are taking a medicine that has been connected to seizures, such as bupropion or selective serotonin reuptake inhibitors (SSRIs)

Seizures have also happened in people who have not had any of these problems. **Call your child's doctor right away if your child has a seizure while taking LUPRON DEPOT-PED.**

LUPRON DEPOT-PED should not be taken if your child is:

- Allergic to GnRH, GnRH agonist medicines, or any ingredients in LUPRON DEPOT-PED. See the end of the Medication Guide for a complete list of ingredients in LUPRON DEPOT-PED.
- Pregnant or becomes pregnant. LUPRON DEPOT-PED can cause birth defects or loss of the baby. If your child becomes pregnant, call your child's doctor.

Before your child receives LUPRON DEPOT-PED, tell their doctor about all of your child's medical conditions, including if they:

- Have a history of mental (psychiatric) problems
- Have a history of seizures
- Have a history of epilepsy
- Have a history of brain or brain vessel (cerebrovascular) problems or tumors
- Are taking a medicine that has been connected to seizures such as bupropion or selective serotonin reuptake inhibitors (SSRIs)
- Are breastfeeding or plans to breastfeed. It is not known if LUPRON DEPOT-PED passes into the breast milk.

Tell your child's doctor about all the medicines your child takes, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How will your child receive LUPRON DEPOT-PED?

- Your child's doctor should do tests to make sure your child has CPP before treating them with LUPRON DEPOT-PED.
- LUPRON DEPOT-PED is given as a single-dose injection into your child's muscle each month or every 3 months by a doctor or trained nurse. Your child's doctor will decide how often your child will receive the injection.
- Keep all scheduled visits to the doctor. If a scheduled dose is missed, your child may start having signs of puberty again. The doctor will do regular exams and blood tests to check for signs of puberty.

What are the possible side effects of LUPRON DEPOT-PED?

LUPRON DEPOT-PED may cause serious side effects. See "What is the most important information I should know about LUPRON DEPOT-PED?"

- **The most common side effects of LUPRON DEPOT-PED received 1 time each month include:**
 - Injection site reactions such as pain, swelling, and abscess
 - Weight gain
 - Pain throughout body
 - Headache
 - Acne or red, itchy rash and white scales (seborrhea)
 - Serious skin rash (erythema multiforme)
 - Mood changes
 - Swelling of vagina (vaginitis), vaginal bleeding, and vaginal discharge
- **The most common side effects of LUPRON DEPOT-PED received every 3 months include:**
 - Injection site reactions such as pain and swelling
 - Weight gain
 - Headache
 - Mood changes

These are not all the possible side effects of LUPRON DEPOT-PED. **Call your child's doctor for medical advice about side effects.**

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call **1-800-FDA-1088**.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/myAbbVieAssist to learn more.

This is the most important information to know about LUPRON DEPOT-PED. For more information, talk to your child's doctor or healthcare provider.

This email has been sent to you because you signed up to receive information about Central Precocious Puberty. If you no longer wish to receive emails from LUPRON DEPOT-PED, please [click here](#).

We respect your privacy. For more information, please view our [privacy policy](#). Please DO NOT REPLY TO THIS MESSAGE. [Contact us](#) with questions or comments.

Mail written correspondence to:
AbbVie Customer Service
Department 36M
1 N Waukegan Road
North Chicago, IL 60064-6163

© 2022 AbbVie. All rights reserved.

LUPRON DEPOT® is a registered trademark of AbbVie Inc.
US-LUPR-220054 March 2022

[PRIVACY POLICY](#) | [CONTACT US](#) | [TERMS OF USE](#)

abbvie

From: Central Precocious Puberty Support [mailto:Info@csm.LP.abbvie.com]

To: John Doe <john.doe@example.com>

Subject: Central Precocious Puberty (CPP) treatment tips

Injection fears are normal—we're here to help.

[View in browser](#)



Dear [recipient name],

Thank you for signing up to receive updates from LUPRON DEPOT-PED while your child is on treatment.

By now, your child is ready to begin or has started their treatment regimen with LUPRON DEPOT-PED. Talking with them about their injection fears is important to their success on treatment. There are a few things to know about what to expect at the doctor's office.

MANAGING TREATMENT EXPECTATIONS WITH YOUR CHILD CAN PAVE THE WAY TO A SUCCESSFUL TREATMENT JOURNEY

Here are some tips to help prepare your child for their upcoming visits to the doctor and what you can expect when you get there.

BEFORE

- Help your child relax on the way to the doctor by playing a game or making up a song
- In the waiting room, having cards, a book, or something that can help distract your child can help ease some of the anxiety

DURING

- In the exam room, the nurse or medical assistant will prepare your child for the injection
- The nurse may do an activity to help your child relax their muscles
- The injection site can vary between the upper arm, buttocks, or thigh to make the experience easier for your child

AFTER

- After the injection, some pain and irritation at the site is expected. If more severe symptoms occur or develop, contact your child's doctor right away

HELPFUL WAYS TO DISTRACT YOUR CHILD IF THEY ARE FEELING ANXIOUS

If your child is scared of injections, there are some distraction techniques that can help. Here are just a few to get you started. Be sure to get creative! Remember, you know your child best, so do what works for them.

- Playing with toys
- Listening to music
- Reading books
- Watching videos

How you act can also play a big role in helping reduce anxiety. If you're anxious, your child may be anxious. If you're calm, your child will be calm.

Remember that with LUPRON DEPOT-PED, Support PLUS provides benefits, cost-saving information, and nurse support. [Learn more about Support PLUS.](#)

LEARN MORE ABOUT HOW YOU CAN ASSIST YOUR CHILD IN UNDERSTANDING THEIR TREATMENT PROCESS AND WHAT TO EXPECT AT THE DOCTOR'S OFFICE

[Learn more about starting therapy](#)

Use for LUPRON DEPOT-PED®(leuprolide acetate for depot suspension)

LUPRON DEPOT-PED 7.5 mg, 11.25 mg, and 15 mg for 1-month and 11.25 mg and 30 mg for 3-month administration are prescribed for the treatment of children with central precocious puberty (CPP).

It is not known if LUPRON DEPOT-PED is safe and effective in children under 2 years of age.

Important Safety Information for LUPRON DEPOT-PED

What is the most important information I should know about LUPRON DEPOT-PED?

- During the first 2 to 4 weeks of treatment, LUPRON DEPOT-PED can cause an increase in some hormones. During this time, you may notice more signs of puberty in your child, including vaginal bleeding. **Call your child's doctor if these signs continue after the second month of treatment with LUPRON DEPOT-PED.**
- Some people taking gonadotropin releasing hormone (GnRH) agonists like LUPRON DEPOT-PED have had new or worsened mental (psychiatric) problems. Mental (psychiatric) problems may include emotional symptoms such as:
 - Crying
 - Irritability
 - Restlessness (impatience)
 - Anger
 - Acting aggressive

Call your child's doctor right away if your child has any new or worsening mental symptoms or problems while taking LUPRON DEPOT-PED.

- Some people taking GnRH agonists like LUPRON DEPOT-PED have had seizures. The risk of seizures may be higher in people who:
 - Have a history of seizures
 - Have a history of epilepsy
 - Have a history of brain or brain vessel (cerebrovascular) problems or tumors
 - Are taking a medicine that has been connected to seizures, such as bupropion or selective serotonin reuptake inhibitors (SSRIs)

Seizures have also happened in people who have not had any of these problems. **Call your child's doctor right away if your child has a seizure while taking LUPRON DEPOT-PED.**

LUPRON DEPOT-PED should not be taken if your child is:

- Allergic to GnRH, GnRH agonist medicines, or any ingredients in LUPRON DEPOT-PED. See the end of the Medication Guide for a complete list of ingredients in LUPRON DEPOT-PED.
- Pregnant or becomes pregnant. LUPRON DEPOT-PED can cause birth defects or loss of the baby. If your child becomes pregnant, call your child's doctor.

Before your child receives LUPRON DEPOT-PED, tell their doctor about all of your child's medical conditions, including if they:

- Have a history of mental (psychiatric) problems
- Have a history of seizures
- Have a history of epilepsy
- Have a history of brain or brain vessel (cerebrovascular) problems or tumors
- Are taking a medicine that has been connected to seizures such as bupropion or selective serotonin reuptake inhibitors (SSRIs)
- Are breastfeeding or plans to breastfeed. It is not known if LUPRON DEPOT-PED passes into the breast milk.

Tell your child's doctor about all the medicines your child takes, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How will your child receive LUPRON DEPOT-PED?

- Your child's doctor should do tests to make sure your child has CPP before treating them with LUPRON DEPOT-PED.
- LUPRON DEPOT-PED is given as a single-dose injection into your child's muscle each month or every 3 months by a doctor or trained nurse. Your child's doctor will decide how often your child will receive the injection.
- Keep all scheduled visits to the doctor. If a scheduled dose is missed, your child may start having signs of puberty again. The doctor will do regular exams and blood tests to check for signs of puberty.

What are the possible side effects of LUPRON DEPOT-PED?

LUPRON DEPOT-PED may cause serious side effects. See "What is the most important information I should know about LUPRON DEPOT-PED?"

- **The most common side effects of LUPRON DEPOT-PED received 1 time each month include:**
 - Injection site reactions such as pain, swelling, and abscess
 - Weight gain
 - Pain throughout body
 - Headache
 - Acne or red, itchy rash and white scales (seborrhea)
 - Serious skin rash (erythema multiforme)
 - Mood changes
 - Swelling of vagina (vaginitis), vaginal bleeding, and vaginal discharge
- **The most common side effects of LUPRON DEPOT-PED received every 3 months include:**
 - Injection site reactions such as pain and swelling
 - Weight gain
 - Headache
 - Mood changes

These are not all the possible side effects of LUPRON DEPOT-PED. **Call your child's doctor for medical advice about side effects.**

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

If you are having difficulty paying for your medicine, AbbVie may be able to help. Visit AbbVie.com/myAbbVieAssist to learn more.

This is the most important information to know about LUPRON DEPOT-PED. For more information, talk to your child's doctor or healthcare provider.

This email has been sent to you because you signed up to receive information about Central Precocious Puberty. If you no longer wish to receive emails from LUPRON DEPOT-PED, please [click here](#).

We respect your privacy. For more information, please view our [privacy policy](#). Please DO NOT REPLY TO THIS MESSAGE. [Contact us](#) with questions or comments.

Mail written correspondence to:
AbbVie Customer Service
Department 36M
1 N Waukegan Road
North Chicago, IL 60064-6163

© 2022 AbbVie. All rights reserved.

LUPRON DEPOT® is a registered trademark of AbbVie Inc.
US-LUPR-220054 March 2022

[PRIVACY POLICY](#) | [CONTACT US](#) | [TERMS OF USE](#)

abbvie