



LULEÅ UNIVERSITY OF TECHNOLOGY

POLAR ATMOSPHERIC PHYSICS

F7014R

---

# EISCAT Space Weather

---

*Authors*

A. Hoehne  
D. Talavera  
E.F.M. Weterings

*Supervisors*

A. Tjulin  
C.F. Enell  
V. Barabash

May 20, 2019

# Table of Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>GUISDAP software</b>	<b>3</b>
<b>3</b>	<b>Space weather event</b>	<b>6</b>
3.1	Solar flares	6
3.2	GOES	6
3.3	IMAGE	8
<b>4</b>	<b>EISCAT Madrigal</b>	<b>9</b>
<b>5</b>	<b>International Reference Ionosphere (IRI)</b>	<b>11</b>
<b>6</b>	<b>Discussion</b>	<b>12</b>
<b>7</b>	<b>Confirmation of Participation</b>	<b>12</b>

# 1 Introduction

example MISC **exampleMISC**.  
example Book **exampleBOOK**.  
example article **exampleARTICLE**

## 2 GUISDAP software

In this section the raw EISCAT data is processed using the MATLAB software with the package GUISDAP. The data analyzed in this section was obtained with the 42m radar located in Svalbard, Norway. The time frame of the analyzed data corresponds to the 27th of October, 2016, from 15:00 to 21:00.

GUISDAP have several "experiment" methods. From the EISCAT experiments manual: "An EISCAT experiment is a set of instructions telling the transmitters, receivers and digital signal processing units what to do at what time". Meaning that depending on the object of interest of the experimenter, Some of the parameters that change between experiments are the code length in bits, baud length, sampling rate, the range span (measurable height), time resolution, etc. In this particular case, the experiment "ipy" was used.

With the help of GUISDAP, a few plots showing different parameters observed in the ionosphere at the time period mentioned above. The following parameters are used by the ipy experiment. One of the obtained parameters is the raw electron data. The data obtained with GUISDAP which is data measured from the EISCAT radar, was compared with data formulated using the IRI model in the following pictures.




Figure 1: Electron density data obtained from EISCAT 2016-10-27.




Figure 2: Electron density data from the IRI model 2016-10-27

In these two images, the electron density from 0 to 600 km was plotted in order to compare the IRI model to the measurements from EISCAT. Both are showing the electron density distribution at around 18:00 hours. While both the model and measurements agree on where the maximum occurs, the model underestimates the amount of electrons by about a factor of 3. More on the IRI model will be discussed in a later chapter.



# EISCAT Scientific Association

## EISCAT SVALBARD RADAR

CP, 42m, ipy, 27 October 2016

Produced@DESKTOP-10LDCOV, 10-May-2019

Not for publication - see Rules-of-the-road




Figure 3: Plot produced analyzing raw data from EISCAT with GUIDAP

Figure 3 shows the results of analyzing 6 hours of data obtained by EISCAT from 15:00 to 21:00 on the 27th of October 2016. In the first row, we can see the electron density which matches with the plots on figure 1 and figure 2, showing a maximum around 300 km of altitude. The second row shows the electron temperature, with respect to each altitude, it is interesting to note that most of the times the variation is smooth and small, except for a small band right before 18:00 and a larger band around 20:30. The third row shows the ion temperature, again, this one is more or less constant along time with a few exceptions around 17:30 and 18:00. The fourth row showing the ion drift velocity, most of the values look like a more or less random noise, meaning that this drift velocity is not consistent. On the fifth and final row we see the data of the antenna within the time range of the data analyzed.

### 3 Space weather event

In the rest of this document preprocessed data is used from the Halloween 2003 space weather event. This event took place from the 28th up to the 29th of October, with the main two peaks at 11:10 (28-10-2003) and 20:50 (29-10-2003) **goes'x-ray'archive**. This solar weather event consisted of a series of solar flares and coronal mass ejections. The solar flare with the most energy was measured at 10:16:53 UCT. With an energy of  $6.9 \cdot 10^{25}$  Joule and a mass of  $1.6 \cdot 10^{10}$  gram **CME'list**, one of the strongest ever measured by GOES.

Satellite-based systems and communications were affected, as well as the instruments onboard **swpc-noaa**. Aircraft were advised to avoid high altitudes near the polar regions, and a one-hour-long power outage occurred in Sweden as a result of the solar activity. Auroras (figure 4) were observed at latitudes as far south as Texas and the Mediterranean countries of Europe **wiki'halloween'solar'storms**.




Figure 4: Geomagnetic storm/ Aurora on the 29th of October **spaceweather**.

#### 3.1 Solar flares

On the 28th 12:18 UCT one of the most powerful Coronal Mass Ejections (CME) in years erupted, this eruption, that caused a intense geomagnetic storm, is shown in figure 5. The solar flares erupted out of 486 giant sunspots. It was measured X17 on the Richter scale of solar flares. This means that the peak had an energy above  $7 \cdot 10^{-4}$  W/m<sup>2</sup>. It was also classified as a S3 storm, which means it has a flux of more than  $10^3$  with  $\geq 10$  MeV particles. **spaceweather**.

#### 3.2 GOES

In figure 6 the space weather data from GOES10 (10th Geostationary Operational Environmental Satellite) over the month October is shown. In figure 7 the data from the same satellite is shown over the month of November **ngdc-noaa**.

In the top plot the peak flux of the solar flares/ CMEs in watts per square meter (W/m<sup>2</sup>) between 1 and 8 Ångströms. Based on the flux the solar flare is classified. These are the classifications:

1. Class A: flux  $< 10^{-7}$
2. Class B:  $10^{-7} \leq \text{flux} < 10^{-6}$
3. Class C:  $10^{-6} \leq \text{flux} < 10^{-5}$
4. Class M:  $10^{-5} \leq \text{flux} < 10^{-4}$
5. Class X: flux  $\geq 10^{-4}$




Figure 5: Solar image of a CME using a coronagraph **spaceweather**.

In the second top plot the electrons flux, directed to Earth, is shown in  $\text{e}^- \text{cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$ . With the black line presenting electrons with an energy greater than 0.6 MeV, red above 2 MeV and green above 4 MeV. In the third plot the proton flux is shown, with the energies greater than 1 MeV for black, 5 MeV for red, 10 MeV for green, 30 MeV for pink, 50 MeV for blue, 60 MeV purple and 100 MeV for light blue. In the last plot the magnetic field strength in nT, where black (Hp) is the magnetic field vector component, points northward, perpendicular to the orbit plane which for a zero degree inclination orbit is parallel to Earth's spin axis. Red (He) is the magnetic field vector component, perpendicular to Hp and Hn and points earthward. Green (Hn) is the magnetic field vector

component, perpendicular to Hp and He and points eastward.




Figure 6: GOES10 data from the month October 2003 **ngdc-noaa**.







Figure 7: GOES10 data from the month November 2003 **ngdc-noaa**.

From the GOES data in figure 6 and 7 it can clearly be seen that there were a lot of instabilities during the end of October and the beginning of November. Starting around the 14th of October the magnetic field became more unstable and the proton flux increased. On the 18th the electron flux increased, as well as the flux of energy. The main two peaks were at the 28th and 29th of October, followed by some less, but still pretty strong, flares on the 2nd, 3rd and 4th of November. After the sixth of November the fluxes went down again.

### 3.3 IMAGE

In figure 8 the magnetometer data from multiple latitudes on Earth from the 29th of October to the 1st of November is shown. These values differ from the magnetometer data from GOES due to that GOES is orbiting Earth and these magnetometers are located on a fixed position on the Earth. It can be seen that the latitudes around the Auroral oval have the largest spikes.



(a) Perpendicular to the Y and Z axis (Hn) **image**. (b) Perpendicular to the z axis, pointing Eastward (He) **image**. (c) Axis from the center of the Earth northwards (Hp)**image**.

Figure 8: Magnetogram data from IMAGE (International Monitor for Auroral Geomagnetic Effects).

## 4 EISCAT Madrigal

In figure 9 the EISCAT data from the 30th of October 2003 is shown. There are a few white gaps which can be explained by interference like clouds in the atmosphere. On top we have the electron density, followed by the electron temperature, ion temperature, ion drift velocity and the system temperature.

Around 20:50 on the 29th of October 2003, one of the main peaks during this solar storm was measured. When this peak was over there was a dip in electron density, as can be seen from figure 9. Then, on the 30th of October there were some M-flares at 1:55 till 2:45 and 15:20 till 15:40 measured by GOES [goes'x-ray'archive](#). You can see in the EISCAT data that right after this change has been measured the densities and temperatures started increasing. The peaks are delayed by a couple of hours in this data.

When comparing the data from the 30th of October 2003 to a more quiet day, like the one in figure 3, it can be seen that especially the fluctuations are way more intense. The gradient between upper and lower in the atmosphere is still present. But everything is, on average, more intense. The perturbations go deeper into the atmosphere and with higher velocities.



# EISCAT Scientific Association

## EISCAT SVALBARD RADAR

SP FR, 32m, tau0, 30 October 2003




Figure 9: EISCAT radar data from the upper atmosphere science database Madrigal **madrigal**.

## 5 International Reference Ionosphere (IRI)

This model has been developed by an international collaboration sponsored by the Committee on Space Research (COSPAR) and the international Union of Radio Science (URSE). The development started in the late sixties in order to create a standardized model of the ionosphere by compiling empirical data from the available sources at the time. this model has been updated several times in order to keep it up to date with current measurements. The data of IRI comes ionosondes, incoherent scatter radars, satellites and sounding rockets.

A comparison was made between the data from the IRI model and the data obtained with EISCAT's incoherent scatter radar in Svalbard, Norway. This comparison takes data at around 21:01 hours on the 30th of October of 2003, as part of the Halloween solar storms.




Figure 10: Comparison of the IRI model data (red) and the one obtained from EISCAT through the Madrigal system (blue).

While both sources more or less agree where the maximum of electron density occurs. There is a big discrepancy in the values shown, of about one order of magnitude. This can be attributed to the fact that the IRI model uses a monthly average to display the electron density of a particular time and date.

## 6 Discussion

example MISC **exampleMISC**.  
example Book **exampleBOOK**.  
example article **exampleARTICLE**

## 7 Confirmation of Participation

Hereby, A. Hoehne, D. Talavera and E.F.M. Weterings declare to have participated and collaborated in this report.