

Tao Analysis - My Visualisations and Proofs

Rhitt C

January 6, 2026

Contents

1	Abstract, Motivation and Preliminaries	3
1.1	Abstract	3
1.2	Why do Analysis? - Motivation 1	3
1.3	Why do Analysis? - Motivation 2 (very important)	4
1.4	Logic Preliminaries	6
2	Construction of \mathbb{N}	7
2.1	Peano Axioms	7
2.2	Addition	10
2.3	Multiplication	10
3	ZFC Set Theory	11
3.2	Historical Axioms - Too Good to be True	11
3.1	ZFC Axioms	11
3.3	Functions	11
3.4	Images	11
3.5	Cartesian Products	11
3.6	Cardinality	11
4	Constructing \mathbb{Z} and \mathbb{Q}	12
4.1	Constructing \mathbb{Z} - Subtractive Closure of \mathbb{N}	12
4.2	Constructing \mathbb{Q} - Divisive Closure of \mathbb{Z}	12
4.3	Exponentiation of \mathbb{Q} to \mathbb{Z} Powers and the Modulus	12
4.4	Why \mathbb{Q} is Inadequate - Motivation for \mathbb{R}	12
5	Constructing \mathbb{R} - Cauchy Completion of \mathbb{Q} - The (Almost) Perfect Field	13
5.0	Aside - Dedekind Construction of \mathbb{R}	13
5.1	Cauchy Sequences	13
5.2	Defining Equivalence of Cauchy Sequences	13
5.3	Cauchy Construction of \mathbb{R}	13
5.4	Defining an Order on \mathbb{R}	13
5.5	The Least Upper Bound Property (LUBP) - Fixing the Problems of \mathbb{Q}	13
5.6	Exponentiation of \mathbb{R} to \mathbb{Q} Powers	13
6	Limits of Sequences	14
6.1	Convergence and Limit Laws	14
6.2	Defining $\mathbb{R}^[-\infty, \infty]$	14
6.3	Suprema and Infima	14
6.4	Limsup, Liminf and Limit Points	14
6.5	Some Standard Limits	14
6.6	Subsequences	14
6.7	As Far as Exponentiation of \mathbb{R} Goes - Inadequacies of \mathbb{R}	14

7	Series	15
7.1	Finite Series	15
7.2	Infinite Series	15
7.3	Series with $[0, \infty)$	15
7.4	The Problem with Rearranging Series	15
7.5	Root and Ratio Tests	15
8	Handling ∞	16
8.1	$ \mathbb{N} $ Cardinality	16
8.2	Summation on Infinite Sets	16
8.3	$ \mathbb{R} $ Cardinality	16
8.4	Axiom of Choice - Full Discussion	16
8.5	Well-Ordering of $ R $ Sets	16
9	Continuity	17
9.1	Subsets of \mathbb{R}	17
9.2	Algebra on \mathbb{R} -valued Functions	17
9.3	Limiting Values of Functions	17
9.4	Continuous Functions	17
9.5	Left and Right Limits	17
9.6	The Max Principle	17
9.7	The Intermediate Value Theorem (IVT)	17
9.8	Monotonicity	17
9.9	Uniform Continuity	17
9.10	Limits at ∞	17
10	Differentiation	18
11	Riemann Integral	19
12	Metric Spaces	20
13	Generalising Continuity to Metric Spaces	21
14	Uniform Convergence	22
15	Power Series	23
16	Fourier Series	24
17	Multivariable Differential Calc	25
18	Lebesgue Measure - oooh	26
19	Lebesgue Integration	27

1 Abstract, Motivation and Preliminaries

1.1 Abstract

I decided to write this text as a means to consolidate my personal understanding of undergraduate Real Analysis, and also to practise the skill of writing academic papers.

It is heavily inspired and guided by Terrence Tao's Analysis Vol. I and II. However, I have taken many liberties in altering formulations to my liking, completely spelling out motivating arguments before most definitions and theorems (even if this makes it clunky at times), going on large digressions, and providing visuals. I'm also including my solutions to the exercises in Tao's texts, mainly for personal reference.

As a digression on expression, in aspiration of approbation from Big M's appellation upon examination, the language register is largely formal in an attempt to adhere to writing conventions. But we undergo register shifts towards informality where appropriate, particularly in avoiding the infamous passive voice¹. This is because it would unnecessarily restrict syntactic freedom in a situational context where it is the *content* that matters far more than the expression. Thus, it is more favourable to be able to convey the content precisely and elegantly without jumping through linguistic hoops every step of the way.

1.2 Why do Analysis? - Motivation 1

The motivation for Analysis is twofold:

Firstly, the most obvious reason is to understand deeper concepts that stray far from the base intuition you get from VCE and high school in general. For example, you may have heard the result that there are more $x \in \mathbb{R}$ with $x \in (0, 1)$ than there are $p \in \mathbb{Q}$ with $p \in (-\infty, \infty)$ ². Your intuition tells you this is absurd, because clearly both are infinite! How can we have $\infty_1 > \infty_2$? Even more counterintuitive is that there are *just as many* $n \in \mathbb{N}$ as there are $x \in \mathbb{Z}$, even though clearly for every $n \in \mathbb{N}$ there are both $\pm n \in \mathbb{Z}$, and so you'd expect there to be about twice as many integers as there are naturals. It seems like mathematicians are just making stuff up at this point. But in Real Analysis, we handle ∞ *rigorously* to make such statements not only meaningful, but to fully understand the motivation for even discussing such matters in the first place. The full level of rigour makes many theorems feel discoverable, in that you properly understand how you would derive and proof them in the first place.

We will see how it answers exactly what you can and can't do with ∞ . E.g. Consider the matrix

$$M = \begin{pmatrix} 1 & 0 & 0 & \cdots \\ -1 & 1 & 0 & \cdots \\ 0 & -1 & 1 & \cdots \\ 0 & 0 & -1 & \cdots \\ \cdots & \cdots & \cdots & \ddots \end{pmatrix}$$

You'd expect the sum of all elements $\sum M_{ij}$ to be the same either way, whether you sum over rows or over columns first. But we have, if you'll excuse the lax notation,

$$\begin{aligned} \sum_{\text{over rows first}} M &= \sum_{\text{over columns}} \begin{pmatrix} 1 \\ 0 \\ 0 \\ \vdots \end{pmatrix} = 1, & \sum_{\text{over columns first}} M &= \sum_{\text{over rows}} (0 \ 0 \ 0 \ \cdots) = 0 \\ \therefore \left(\sum_{\text{over rows first}} M = 1 \right) &\neq \left(\sum_{\text{over columns first}} M = 0 \right) \end{aligned}$$

¹why is this even a scientific convention smh ts pmo

²the precise statement is with a concept called *cardinality*

So we have illustrated that you can't always interchange the order in which you sum infinitely many numbers, contrary to intuition. Real Analysis gives the full derivation for the precise conditions in which you can safely do so³. As we have seen, it teaches you how to properly deal with ∞ . So in a sense, Real Analysis can make you like *The Man Who Knew Infinity*⁴.

More than that, Real Analysis answers to the highest echelon that is possible with logic (provably so, in fact) the *why* behind essentially everything you've ever been taught in maths from kindergarten to high school, leave some small details about \mathbb{C} . You realise the true degree of freedom that maths gives you. And you learn complete objectivity. You don't have to take anyone's word for it, just prove it yourself. You learn to make a statement, and the proof of it, so damn precise that there is *zero* ambiguity or subjectivity in it; I say this without hyperbole. Doesn't matter if a teacher thinks your expression is lacking; you can be *absolutely sure* that your statement is correct, given you're subtle enough with what is meant by correct.

1.3 Why do Analysis? - Motivation 2 (very important)

Secondly, we will see how blindly following the whims of what our intuition deems reasonable can very easily lead to contradiction. The many times that this occurred historically, there was great philosophical anguish about what assumptions to keep and what to discard, substantially impeding on mathematical progress.

For example, take $\sqrt{2}$, a quantity that arises in geometry very naturally as the length of the hypotenuse in a triangle with both other side lengths 1. In ancient Greece, many held the worldview that everything in the real world, like this hypotenuse, can be quantified as a rational number in \mathbb{Q} . And they held this assumption to be so blatantly obvious that it underpinned their philosophy of everything from music to the universe itself. So when it was found that $\sqrt{2} \in \mathbb{Q}$ would imply a contradiction, even though the triangle is very much real, it led to a crisis in which many ancient Greeks had to reevaluate their entire philosophy.



Because we've been taught from a young age that irrational numbers exist and do indeed show up in the real world, we now find the concept to be no doubt very intuitive. But how can we be absolute *sure* that there isn't some other cancerous contradiction hiding somewhere in our mathematical foundations as well, just waiting to spread into everything we think we know, as it did for the Greeks? The truth is, we're *not* sure. And it's in fact been *proven*⁵ that we can never be sure. Not unless we rob ourselves of the ability to do primary school arithmetic, which would be very absurd. So as we shall see, we do the next best thing: minimise our set of assumptions to the bare minimum that is still feasible.

To be clear, the issue isn't with logic itself. As a rather lengthy aside, the question of whether logic is valid is philosophical in nature, and seemingly can't ever have a definitive resolution. For example, it could very well be the case that we and our fictitious universe only exist in the incoherent dream of a butterfly. If so, logic as we perceive it only *appears* valid to us, yet completely wrong to a scientist observing the dream from outside in the real world. But even though such fundamental questions about reality may be interesting, doubting logic every step of the way puts overly restrictive bounds on what mathematical progress is possible. So they're for the philosopher to ponder over. Instead, the mathematician prescribes a handful of intuitive operations on True/False values, which formalise pure logic. E.g.

(True & True) is True, but

(True & False) is False, as True and False are said to contradict each other.

Then they prescribe a few more rules on how to produce logical statements (with True/False value) from free variables (which can be mathematical objects), completing the formal link from maths to logic. E.g.

$$((\forall x, P(x)) \implies P(0)) \text{ is True}$$

³see Fubini

⁴cold, cold, cold

⁵see Gödel's Incompleteness Theorems

Finally, they combine all of this into a rulebook which they call the formal logic. Confident that the logic reflects reality, they hand it over to the philosopher to question the validity of. The mathematician is now happy to accept their logic as absolute and instead ponders questions about how the logic actually affects mathematical statements (theorems, lemmas, corollaries, etc.). And even if the philosopher someday finds that the logic is only valid in the dream of a butterfly and invalid outside of it, much of the work done by mathematicians could very well still be moulded to adhere to whatever rulebook is followed by the world outside the dream.

Now that we have set our logic in stone, how do we proceed? Logic can only tell us about the relationships between logical statements, but makes no comment about many individual statements involving free variables. E.g. we can show

$$(\text{I am a lazy person} \ \& \ \text{all lazy people are monkeys}) \implies (\text{I am a monkey})^6$$

as a whole is True directly from our logic, but we can't individually show that (I am a lazy person) or (all lazy people are monkeys) are True with logic alone because of the free variables (I), (lazy person/people) and (monkey(s)). And if either of those turn out to be False, then we have no information about whether (I am a monkey) is True or False. I could very well be a monkey despite not being lazy. Thus, all in all, we have an (\implies) statement that is True but we can't show definitely that its implication that (I am a monkey) is also True... which is a good thing because I'm pretty confident I'm not a monkey (but can I *show* definitively that I'm not? Once again, no).

This is a necessary tradeoff we made to establish barebone logic the mathematician will never have to doubt.

So how do we prove a statement P (called a conjecture until proof is established)? Instead of the fool's errand of looking for whether P is definitively True, we loosen our standards slightly and instead show the truth of an (\implies) statement as follows: if we accept a base few assumptions, which we call our *axioms*, then C is True. That is, our standard of proof for P is:

$$Axiom_1 \ \& \ Axiom_2 \ \& \ Axiom_3 \ \& \ \dots \implies C$$

As a concluding remark, we do not deny the value of intuition. In fact, intuition is absolutely essential for *developing* a conjecture and motivating why it might be plausible. Without it we are stuck running a fool's errand of perpetual guess and check, never able to come up with something original and insightful. But once we are confident that our conjecture is indeed plausible, we *need* to follow a gold standard of proof independent of intuition. It's the best way we know of to minimise the chance as much as feasible for a contradiction to sneak up on us from a single far-reaching result we prematurely took as granted long ago. That would shatter the foundations of mathematics, much like the contradictory assumption $\sqrt{2} \in \mathbb{Q}$ did for the Greeks. Of course, in perhaps the greatest failure of modern maths, we've shown logically that we still can't ever completely eliminate that possibility, so long as we demand our axioms be strong enough for maths to be of value. Nonetheless, mathematicians collectively agree on a handful of axioms called *ZF* (all of which we will encounter in the next 2 chapters) as the basis for our gold standard of proof, as it's the best that we can feasibly do, and it seems unlikely that it'll ever truly see a contradiction.

The gold standard we set is hence:

Definition D.1: The Gold Standard of Proof

A conjecture P is considered “proven” *if and only if* we can show the following with our logic:
 $ZF \implies P$

⁶given this is stated in a way that avoids mentioning sets

1.4 Logic Preliminaries

2 Construction of \mathbb{N}

2.1 Peano Axioms

We lay our assumed Peano Axioms that relate to $\mathbb{N} = \{0, 1, 2, \dots\}$, motivated by the construction of \mathbb{N} as an analogue for the physical intuition of “counting numbers”. In doing so, we implicitly rely on some set theory, namely some of what is termed the ZFC Axioms of sets, which will be explored in the next chapter⁷. The Peano and ZFC Axioms are our *only* assumptions in all of Analysis.

We shall see that this is sufficient for a very rich set of results about $\mathbb{N}, \mathbb{Z}, \mathbb{Q}, \mathbb{R}$ and some even more general statements⁸ to follow directly by formal logic from these foundational axioms, given relevant definitions. In fact, almost all of modern maths relies on no more than the 8-9 ZFC and 5 Peano Axioms. As I’m sure you’ll agree after seeing how intuitive the axioms really are, this is quite a remarkable development in modern maths. Motivated by these far-reaching implications of the Peano and ZFC Axioms, we set spelling out how a statement follows from these axioms as the gold standard for proof in modern maths.

We will see how blindly following the whims of what our intuition deems reasonable can very easily lead to contradiction. The many times that this occurred historically, there was great philosophical anguish about what assumptions to discard, greatly impeding on mathematical progress. For example, take $\sqrt{2}$, a quantity that arises in geometry very naturally as the length of a diagonal in a square with side length 1. When a proof arose that $\sqrt{2}$ could not be expressed as a fraction in simplest terms, it caused great distraught for the ancient Greeks.

We start with the bare minimum by demanding that some fixed object, namely 0, be well defined. 0 will be crucial to defining a few later axioms, as it gives us a sort of starting point.

Axiom P.1: Existence of 0

There exists an object 0 that is natural and can be used in later axioms.

$\exists 0 \in \mathbb{N} \quad \text{s.t.} \quad \text{P.3 and P.5 are met.}$

0

Now to populate \mathbb{N} , we introduce a new primitive operation (that will later be superseded by addition) called *succession*, with the property of closure. Informally, we hope that it behaves like our intuition for incrementing a number by 1, and thus forces the infinitely many “counting numbers” to now be well defined and in \mathbb{N} . Denote the succession of all $n \in \mathbb{N}$ as $n++$, and label $(1 := 0++)$, $(2 := 1++)$, $(3 := 2++)$, ...


⁷This is so that the formulation of the Axiom of Infinity can be in terms of \mathbb{N} and is thus much more intuitive. Rest assured, while the Peano Axioms refer to \mathbb{N} , they do not themselves make reference to the Axiom of Infinity, so there’s no circularity. The standard in the literature is to merge the 5 Peano Axioms with our formulation of the ZFC Axiom of Infinity to yield a singular, strong and concise formulation of Axiom of Infinity, so ZFC can stand on its own as the foundation of modern maths. This will be explored briefly in the next chapter as a further digression on pure vs impure set theory than that of Tao’s.

⁸About all structures isomorphic to the ones mentioned, and also much later on, all metric spaces

Axiom P.2: Closure of \mathbb{N} under $(++)$

All naturals have natural successions.

$$\forall n \in \mathbb{N} \quad n \in \mathbb{N} \implies (n++) \in \mathbb{N}$$




However, P.1 and P.2 alone are not strong enough to force \mathbb{N} to populate with our infinitely many elements. In particular, they do not prevent existence of an $n \in \mathbb{N}$ with $n++ = 0$. And yet, informally speaking, this leads to a modulo cycle in which only $n + 1$ elements exist in \mathbb{N} , and everything beyond that overflows back to 0, like in a computer. Modulo sets have many applications, but they are not what we desire as a full set of “counting numbers”, particularly if that n turns out to be 0 itself, leading to a trivial $\mathbb{N} = \{0\}$. So we explicitly force \mathbb{N} to not be such a modulo cycle by demanding that a natural’s succession never overflows back to 0.

Axiom P.3: \mathbb{N} is not a modulo cycle overflowing to 0

No natural’s succession can be 0.

$$\forall n \in \mathbb{N} \quad (n++) \neq 0$$




As it stands, the issue hasn’t been fully resolved yet. There is still the pathological case of an *offset* modulo cycle, in which an $n \in \mathbb{N}$ overflows back so that $n \mapsto m++$ even though we already have a previous $m \mapsto m++$. We now complete our requirements on $(++)$ by demanding no such dual mappings to the same succession i.e. we make $(++)$ *injective*.

Axiom P.4: \mathbb{N} has no otherwise offset modulo cycles either

Succession is injective (one-to-one). That is, distinct naturals have distinct successions.

$$\forall n, m \in \mathbb{N} \quad n \neq m \implies (n++) \neq (m++)$$




We now lay out by far the most useful axiom of \mathbb{N} to handle the sheer infinite size of \mathbb{N} recursively. It is induction.

Axiom P.5: (PMI) Principle of Mathematical Induction

If a proposition of naturals is met for 0 (Base Case), and it being satisfied for some natural implies it's met for the succession as well (Inductive Case), then as a domino effect, then the proposition is met for all naturals.

$$\forall P : \mathbb{N} \rightarrow \{\text{True}, \text{False}\} \quad (P(0) \ \& \ (\forall k \in \mathbb{N}) P(k) \implies P(k++)) \implies \forall n \in \mathbb{N} \quad P(n)$$



Abstractly speaking, all sets X equipped with an analogous operation $S : X \rightarrow X$ (referred to together as an algebraic structure (X, S)), that obeys the 5 Peano Axioms is said to be *isomorphic* with $(\mathbb{N}, (++)$), denoted $(X, S) \cong (\mathbb{N}, (++)$). Note that due to our later discussions only relying on logical implications from the axioms, which are met by (X, S) , the results we derive must apply to every such (X, S) , given we rewrite our definitions accordingly. As such, we do not strictly define a particular $(\mathbb{N}, (++)$), but rather refer abstractly to any such solution (X, S) to Peano. So in a sense, the Peano Axioms lay out a kind of citizenship test, in which (X, S) is considered a natural structure if and only if it passes, and it gets all the perks of being a natural citizen that we will now derive.

There is one final assumption that must be

2.2 Addition

2.3 Multiplication

3 ZFC Set Theory

3.2 Historical Axioms - Too Good to be True

3.1 ZFC Axioms

3.3 Functions

3.4 Images

3.5 Cartesian Products

3.6 Cardinality

4 Constructing \mathbb{Z} and \mathbb{Q}

4.1 Constructing \mathbb{Z} - Subtractive Closure of \mathbb{N}

4.2 Constructing \mathbb{Q} - Divisive Closure of \mathbb{Z}

4.3 Exponentiation of \mathbb{Q} to \mathbb{Z} Powers and the Modulus

4.4 Why \mathbb{Q} is Inadequate - Motivation for \mathbb{R}

5 Constructing \mathbb{R} - Cauchy Completion of \mathbb{Q} - The (Almost) Perfect Field

5.0 Aside - Dedekind Construction of \mathbb{R}

5.1 Cauchy Sequences

5.2 Defining Equivalence of Cauchy Sequences

5.3 Cauchy Construction of \mathbb{R}

5.4 Defining an Order on \mathbb{R}

5.5 The Least Upper Bound Property (LUBP) - Fixing the Problems of \mathbb{Q}

5.6 Exponentiation of \mathbb{R} to \mathbb{Q} Powers

6 Limits of Sequences

6.1 Convergence and Limit Laws

6.2 Defining $\mathbb{R}[-\infty, \infty]$

6.3 Suprema and Infima

6.4 Limsup, Liminf and Limit Points

6.5 Some Standard Limits

6.6 Subsequences

6.7 As Far as Exponentiation of \mathbb{R} Goes - Inadequacies of \mathbb{R}

7 Series

7.1 Finite Series

7.2 Infinite Series

7.3 Series with $[0, \infty)$

7.4 The Problem with Rearranging Series

7.5 Root and Ratio Tests

8 Handling ∞

8.1 $|\mathbb{N}|$ Cardinality

8.2 Summation on Infinite Sets

8.3 $|\mathbb{R}|$ Cardinality

8.4 Axiom of Choice - Full Discussion

8.5 Well-Ordering of $|R|$ Sets

9 Continuity

9.1 Subsets of \mathbb{R}

9.2 Algebra on \mathbb{R} -valued Functions

9.3 Limiting Values of Functions

9.4 Continuous Functions

9.5 Left and Right Limits

9.6 The Max Principle

9.7 The Intermediate Value Theorem (IVT)

9.8 Monotonicity

9.9 Uniform Continuity

9.10 Limits at ∞

10 Differentiation

11 Riemann Integral

12 Metric Spaces

13 Generalising Continuity to Metric Spaces

14 Uniform Convergence

15 Power Series

16 Fourier Series

17 Multivariable Differential Calc

18 Lebesgue Measure - oooh

19 Lebesgue Integration