

SmartCrop AI

Crop Disease Analysis Report

Generated: 27 April 2026 13:35 | Location: Khudai, Punjab

A. Prediction Summary

Field	Result
Status	DISEASE DETECTED
Crop	Tomato
Detected Disease	Late Blight (Tomato)
Confidence	92.5%
Risk Level	HIGH

B. Weather Conditions

Parameter	Value
Temperature	34.0 deg C
Humidity	78.0%
Wind Speed	12.0 km/h
Rainfall	2.5 mm

C. Weather Risk Analysis

Humidity is elevated (70-80%), which significantly increases the risk of fungal and bacterial disease spread. Apply preventive fungicides if not already done. High temperatures (≥ 32 deg C) accelerate pathogen reproduction cycles and stress crops, reducing natural disease resistance. Light rainfall prolongs leaf wetness duration, which can facilitate spore germination and infection.

D. Immediate Recommendation

Isolate affected plants immediately to prevent spread of Late Blight (Tomato). Remove infected leaves, begin chemical treatment within 24-48 hours, and recheck all nearby plants for early symptoms.

- Isolate and mark all visibly affected plants immediately.
- Document spread by photographing surrounding plants.
- Begin chemical or organic treatment within 24-48 hours.

- Inform neighbouring farmers if outbreak risk is High.

F. Prevention Strategies

- Maintain 30-45 cm plant spacing to ensure air circulation and reduce leaf wetness.
- Use drip or furrow irrigation instead of overhead watering.
- Rotate crops every season — avoid replanting the same family for 2-3 years.
- Source certified, disease-resistant seed varieties suited to your region.
- Remove and destroy crop debris after harvest to eliminate overwintering pathogens.
- Scout fields twice weekly during high-humidity and rainy periods.

SmartCrop AI | Crop Disease Detection System

Report generated on 27 April 2026 13:38

DISCLAIMER: This report is generated by an AI model and is intended as a decision-support tool only. Always consult a certified agronomist before applying chemical treatments.