

# COOPERATIVE GAME THEORY

RICHARD MUŽÍK

RICHARD@IMUZYK.CZ

APRIL 21, 2023

# INTRODUCTION

## Cooperative game

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

■  $S \subseteq N$  ... coalition

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

- $S \subseteq N$  ... coalition
- $v(S)$  ... value of coalition

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

- $\Gamma^n$  ... set of  $n$ -person cooperative games
- $S \subseteq N$  ... coalition
- $v(S)$  ... value of coalition

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

- $\Gamma^n$  ... set of  $n$ -person cooperative games
- $S \subseteq N$  ... coalition
- $v(S)$  ... value of coalition
- usually  $N = \{1, \dots, n\}$



## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

- $\Gamma^n$  ... set of  $n$ -person cooperative games
- $S \subseteq N$  ... coalition
- $v(S)$  ... value of coalition
- usually  $N = \{1, \dots, n\}$ 
  - ▶  $(S, v_S)$  is **subgame**  $(N, v)$ :
    - $v_S: 2^S \rightarrow \mathbb{R}$
    - $v_S(T) := v(T)$  pro  $T \subseteq S$

## COOPERATION - EXAMPLES OF MODELS: *MINIMAL SPANNING-TREE GAMES*

Goal: *Find the best connection of players to a source*

## COOPERATION - EXAMPLES OF MODELS: *MINIMAL SPANNING-TREE GAMES*

Goal: *Find the best connection of players to a source*

- $N = N' \cup \{o\}$  ... set of players + source

# COOPERATION - EXAMPLES OF MODELS:

## *MINIMAL SPANNING-TREE GAMES*

Goal: *Find the best connection of players to a source*

- $N = N' \cup \{o\}$  ... set of players + source
- $c_{ij}$  ... cost of connecting  $i, j$

## COOPERATION - EXAMPLES OF MODELS:

### *MINIMAL SPANNING-TREE GAMES*

Goal: *Find the best connection of players to a source*

- $N = N' \cup \{o\}$  ... set of players + source
- $c_{ij}$  ... cost of connecting  $i, j$
- solution: a network, where each  $i \in N$  is connected to  $o$  with minimal sum of costs

# GOAL OF THE MODEL OF COOPERATIVE GAMES

*Money first!*

# GOAL OF THE MODEL OF COOPERATIVE GAMES

*Money first!*

- **Payoff vector  $\mathbf{x} \in \mathbb{R}^n$** 
  - ▶  $x_i$  represents payoff of player  $i$

# GOAL OF THE MODEL OF COOPERATIVE GAMES

*Money first!*

- **Payoff vector**  $\mathbf{x} \in \mathbb{R}^n$ 
  - ▶  $x_i$  represents payoff of player  $i$
- Vector  $\mathbf{x} \in \mathbb{R}^n$  is **efficient**, if  $\sum_{i \in N} x_i = v(N)$



# GOAL OF THE MODEL OF COOPERATIVE GAMES

*Money first!*

- **Payoff vector**  $\mathbf{x} \in \mathbb{R}^n$ 
  - ▶  $x_i$  represents payoff of player  $i$
- Vector  $\mathbf{x} \in \mathbb{R}^n$  is **efficient**, if  $\sum_{i \in N} x_i = v(N)$ 
  - ▶ Usually, we distribute  $v(N)$ 
    1. *value of cooperation*  $v(N)$
    2. *shared costs*  $c(N)$

# GOAL OF THE MODEL OF COOPERATIVE GAMES

*Money first!*

- **Payoff vector**  $\mathbf{x} \in \mathbb{R}^n$ 
  - ▶  $x_i$  represents payoff of player  $i$
- Vector  $\mathbf{x} \in \mathbb{R}^n$  is **efficient**, if  $\sum_{i \in N} x_i = v(N)$ 
  - ▶ Usually, we distribute  $v(N)$ 
    1. *value of cooperation*  $v(N)$
    2. *shared costs*  $c(N)$
- Vector  $\mathbf{x} \in \mathbb{R}^n$  is **individually rational**, if  $x_i \geq v(i)$

# GOAL OF THE MODEL OF COOPERATIVE GAMES

*Money first!*

- **Payoff vector**  $\mathbf{x} \in \mathbb{R}^n$ 
  - ▶  $x_i$  represents payoff of player  $i$
- Vector  $\mathbf{x} \in \mathbb{R}^n$  is **efficient**, if  $\sum_{i \in N} x_i = v(N)$ 
  - ▶ Usually, we distribute  $v(N)$ 
    1. *value of cooperation*  $v(N)$
    2. *shared costs*  $c(N)$
- Vector  $\mathbf{x} \in \mathbb{R}^n$  is **individually rational**, if  $x_i \geq v(i)$ 
  - ▶ players prefer  $x_i$  over  $v(i)$

# GOAL OF THE MODEL OF COOPERATIVE GAMES

*Money first!*

- **Payoff vector**  $\mathbf{x} \in \mathbb{R}^n$ 
  - ▶  $x_i$  represents payoff of player  $i$
- Vector  $\mathbf{x} \in \mathbb{R}^n$  is **efficient**, if  $\sum_{i \in N} x_i = v(N)$ 
  - ▶ Usually, we distribute  $v(N)$ 
    1. *value of cooperation*  $v(N)$
    2. *shared costs*  $c(N)$
- Vector  $\mathbf{x} \in \mathbb{R}^n$  is **individually rational**, if  $x_i \geq v(i)$ 
  - ▶ players prefer  $x_i$  over  $v(i)$
- $\mathcal{I}^*(v) = \{\mathbf{x} \in \mathbb{R}^n \mid \mathbf{x}(N) = v(N)\}$  ... **preimputation**
  - ▶  $\mathbf{x}(S) := \sum_{i \in S} x_i$

# GOAL OF THE MODEL OF COOPERATIVE GAMES

*Money first!*

- **Payoff vector**  $\mathbf{x} \in \mathbb{R}^n$ 
  - ▶  $x_i$  represents payoff of player  $i$
- Vector  $\mathbf{x} \in \mathbb{R}^n$  is **efficient**, if  $\sum_{i \in N} x_i = v(N)$ 
  - ▶ Usually, we distribute  $v(N)$ 
    1. *value of cooperation*  $v(N)$
    2. *shared costs*  $c(N)$
- Vector  $\mathbf{x} \in \mathbb{R}^n$  is **individually rational**, if  $x_i \geq v(i)$ 
  - ▶ players prefer  $x_i$  over  $v(i)$
- $\mathcal{I}^*(v) = \{x \in \mathbb{R}^n \mid x(N) = v(N)\}$  ... **preimputation**
  - ▶  $x(S) := \sum_{i \in S} x_i$
- $\mathcal{I}(v) = \{x \in \mathcal{I}^*(v) \mid \forall i \in N : x_i \geq v(i)\}$  ... **imputation**

# **SOLUTION CONCEPTS**



- Set of payoff vectors satisfying further properties are **solution concepts**



- Set of payoff vectors satisfying further properties are **solution concepts**
- Can reflect payoff distribution, which is

- Set of payoff vectors satisfying further properties are **solution concepts**
- Can reflect payoff distribution, which is
  - ▶ ...*fair*...
  - ▶ ...*non-discriminatory*...
  - ▶ ...*stable (players will accept it)*...
  - ▶ ...

Formally:

Formally:

1. sets of payoff vectors

Formally:

1. sets of payoff vectors

►  $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$

Formally:

1. sets of payoff vectors
  - ▶  $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$
2. functions on games

Formally:

1. sets of payoff vectors

▶  $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$

2. functions on games

▶  $\Sigma: \Gamma^n \rightarrow 2^{\mathbb{R}^n}$

Formally:

1. sets of payoff vectors

▶  $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$

2. functions on games

▶  $\Sigma: \Gamma^n \rightarrow 2^{\mathbb{R}^n}$

We distinguish



Formally:

1. sets of payoff vectors
  - ▶  $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$
2. functions on games
  - ▶  $\Sigma: \Gamma^n \rightarrow 2^{\mathbb{R}^n}$

We distinguish

1. **single-point** solution concepts
2. **multi-point** solution concepts

Formally:

1. sets of payoff vectors
  - ▶  $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$
2. functions on games
  - ▶  $\Sigma: \Gamma^n \rightarrow 2^{\mathbb{R}^n}$

We distinguish

1. **single-point** solution concepts
  - ▶ as a set:  $\Sigma(v) = \{x\}$ 
    - we prefer:  $\Sigma(v) = x$
2. **multi-point** solution concepts

Formally:

1. sets of payoff vectors

▶  $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$

2. functions on games

▶  $\Sigma: \Gamma^n \rightarrow 2^{\mathbb{R}^n}$

We distinguish

1. **single-point** solution concepts

▶ as a set:  $\Sigma(v) = \{x\}$

■ we prefer:  $\Sigma(v) = x$

▶ as a function:  $\Sigma: \Gamma^n \rightarrow \mathbb{R}$

2. **multi-point** solution concepts

# MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

# MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

The core

# MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

# MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

■ assumption: *homo economicus*

# MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

■ assumption: *homo economicus*

- ▶ model of human as a player
- ▶ strictly rational and selfish
- ▶ follows his subjective goals



# MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

- assumption: *homo economicus*
  - ▶ model of human as a player
  - ▶ strictly rational and selfish
  - ▶ follows his subjective goals
- $v(N)$  ... value, which is distributed among players

# MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

- assumption: *homo economicus*
  - ▶ model of human as a player
  - ▶ strictly rational and selfish
  - ▶ follows his subjective goals
- $v(N)$  ... value, which is distributed among players
- $x(S) > v(S) \implies$  coalition  $S$  does not leave  $N$

# MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

- assumption: *homo economicus*
  - ▶ model of human as a player
  - ▶ strictly rational and selfish
  - ▶ follows his subjective goals
- $v(N)$  ... value, which is distributed among players
- $x(S) > v(S) \implies$  coalition  $S$  does not leave  $N$ 
  - ▶ would lead to  $(S, v_S)$
  - ▶  $v(S)$  ... distributed value

# MULTI-POINT SOLUTION CONCEPT: THE CORE

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

# MULTI-POINT SOLUTION CONCEPT: THE CORE

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

Reminder

Nash equilibrium

# MULTI-POINT SOLUTION CONCEPT: THE CORE

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

## Reminder

## Nash equilibrium

Strategy profile  $(s_1, \dots, s_n)$  is **Nash equilibrium**, if it holds for every player  $i$ ,

$$v_i(s_1, \dots, s_{i-1}, s_i, s_{i+1}, \dots, s_n) \geq v_i(s_1, \dots, s_{i-1}, t_i, s_{i+1}, \dots, s_n)$$

for every  $t_i \in S_i$ .

# MULTI-POINT SOLUTION CONCEPT: THE CORE

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

## Emptiness of the core

# MULTI-POINT SOLUTION CONCEPT: THE CORE

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

## Emptiness of the core

There are cooperative games  $(N, v)$  with empty core.



# MULTI-POINT SOLUTION CONCEPT: THE CORE

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

## Emptiness of the core

There are cooperative games  $(N, v)$  with empty core.

- Non-essential games

# MULTI-POINT SOLUTION CONCEPT: THE CORE

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

## Emptiness of the core

There are cooperative games  $(N, v)$  with empty core.

- Non-essential games

- $v(N) < \sum_{i \in N} v(i)$

# MULTI-POINT SOLUTION CONCEPT: THE CORE

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

## Emptiness of the core

There are cooperative games  $(N, v)$  with empty core.

- Non-essential games
  - ▶  $v(N) < \sum_{i \in N} v(i)$
- Let  $x \in \mathcal{C}(v)$

# MULTI-POINT SOLUTION CONCEPT: THE CORE

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

## Emptiness of the core

There are cooperative games  $(N, v)$  with empty core.

- Non-essential games

- ▶  $v(N) < \sum_{i \in N} v(i)$

- Let  $x \in \mathcal{C}(v)$

- ▶  $x(N) = v(N) < \sum_{i \in N} x(i) \leq x(N)$

# MULTI-POINT SOLUTION CONCEPT: THE CORE

## The core

For a cooperative game  $(N, v)$ , the **core**  $\mathcal{C}(v)$  is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

## Emptiness of the core

There are cooperative games  $(N, v)$  with empty core.

- Non-essential games
  - ▶  $v(N) < \sum_{i \in N} v(i)$
- Let  $x \in \mathcal{C}(v)$ 
  - ▶  $x(N) = v(N) < \sum_{i \in N} x(i) \leq x(N)$
- $x$  does not exist  $\implies \mathcal{C}(v) = \emptyset$

# MULTI-POINT SOLUTION CONCEPT: THE CORE

Question: *When is the core non-empty?*

## MULTI-POINT SOLUTION CONCEPT: THE CORE

Question: *When is the core non-empty?*

We can encode the core as linear program (P) and determine the dual program (D)

# MULTI-POINT SOLUTION CONCEPT: THE CORE

Question: *When is the core non-empty?*

We can encode the core as linear program (P) and determine the dual program (D)

$$\blacksquare (P) = \begin{cases} \min_{x \in \mathbb{R}^n} & x(N) \\ \text{subject to} & x(S) \geq v(S) \text{ for } S \subseteq N \end{cases}$$



# MULTI-POINT SOLUTION CONCEPT: THE CORE

Question: *When is the core non-empty?*

We can encode the core as linear program (P) and determine the dual program (D)

$$\blacksquare (P) = \begin{cases} \min_{x \in \mathbb{R}^n} & x(N) \\ \text{subject to} & x(S) \geq v(S) \text{ for } S \subseteq N \end{cases}$$

$$\blacksquare (D) = \begin{cases} \max_{y \in \mathbb{R}_+^{(2^n-1)}} & \sum_{S \subseteq N} y_S v(S) \\ \text{subject to} & \sum_{\emptyset \neq S \subseteq N} y_S \chi_S = \chi_N \end{cases}$$

# MULTI-POINT SOLUTION CONCEPT: THE CORE

Question: *When is the core non-empty?*

We can encode the core as linear program (P) and determine the dual program (D)

$$\begin{aligned} \blacksquare (P) &= \begin{cases} \min_{x \in \mathbb{R}^n} & x(N) \\ \text{subject to} & x(S) \geq v(S) \text{ for } S \subseteq N \end{cases} \\ \blacksquare (D) &= \begin{cases} \max_{y \in \mathbb{R}_+^{(2^n-1)}} & \sum_{S \subseteq N} y_S v(S) \\ \text{subject to} & \sum_{\emptyset \neq S \subseteq N} y_S \chi_S = \chi_N \end{cases} \end{aligned}$$

We derive the weak Bondareva-Shapley theorem

## Weak Bondareva-Shapley theorem

Cooperative game  $(N, v)$  has non-empty core if and only if

$$v(N) \geq \sum_{S \subseteq N} y_S v(S) \text{ for all feasible } y \in \mathbb{R}^{(2^n-1)}$$

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

■  $v(S \cup i) - v(S)$

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- $v(S \cup i) - v(S)$ 
  - *marginal contribution* (of player  $i$  in  $S \cup i$ )

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- $v(S \cup i) - v(S)$ 
  - ▶ *marginal contribution* (of player  $i$  in  $S \cup i$ )
- $\frac{s!(n-s-1)!}{n!}$



# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- $v(S \cup i) - v(S)$ 
  - ▶ *marginal contribution* (of player  $i$  in  $S \cup i$ )
- $\frac{s!(n-s-1)!}{n!}$ 
  - ▶ weights reflecting different sizes of coalitions

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- $v(S \cup i) - v(S)$ 
  - ▶ *marginal contribution* (of player  $i$  in  $S \cup i$ )
- $\frac{s!(n-s-1)!}{n!}$ 
  - ▶ weights reflecting different sizes of coalitions
- $\sum_{S \subseteq N \setminus i}$

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- $v(S \cup i) - v(S)$ 
  - ▶ *marginal contribution* (of player  $i$  in  $S \cup i$ )
- $\frac{s!(n-s-1)!}{n!}$ 
  - ▶ weights reflecting different sizes of coalitions
- $\sum_{S \subseteq N \setminus i}$ 
  - ▶ sum of all marginal contributions of  $i$

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:
  1. Form the grandcoalition  $N$ .

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:
  1. Form the grandcoalition  $N$ .
  2. Enter the the coalition individually and randomly.

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:
  1. Form the grandcoalition  $N$ .
  2. Enter the the coalition individually and randomly.
  3. When player  $i$  enters coalition  $S$ , he receives  $v(S \cup i) - v(S)$ .



# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:
  1. Form the grandcoalition  $N$ .
  2. Enter the the coalition individually and randomly.
  3. When player  $i$  enters coalition  $S$ , he receives  $v(S \cup i) - v(S)$ .
- $s!(n-s-1)!$  ... number of situations, in which  $i$  enters  $S$

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:
  1. Form the grandcoalition  $N$ .
  2. Enter the the coalition individually and randomly.
  3. When player  $i$  enters coalition  $S$ , he receives  $v(S \cup i) - v(S)$ .
- $s!(n-s-1)!$  ... number of situations, in which  $i$  enters  $S$
- $n!$  ... number of all possible ways to construct  $N$

# SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

## The Shapley value

For a cooperative game  $(N, v)$ , the **Shapley value**  $\phi(v)$  of player  $i$  is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:
  1. Form the grandcoalition  $N$ .
  2. Enter the the coalition individually and randomly.
  3. When player  $i$  enters coalition  $S$ , he receives  $v(S \cup i) - v(S)$ .
- $s!(n-s-1)!$  ... number of situations, in which  $i$  enters  $S$
- $n!$  ... number of all possible ways to construct  $N$
- $\phi_i(v)$  ... the average value of player  $i$ 's payment

# SHAPLEY VALUE

*It is possible to define it using its properties...*

# SHAPLEY VALUE

*It is possible to define it using its properties...*

## The Shapley value

The **Shapley value**  $\phi(v)$  is the only function  $f: \Gamma^n \rightarrow \mathbb{R}$  satisfying for all games  $(N, v), (N, w)$ :

# SHAPLEY VALUE

*It is possible to define it using its properties...*

## The Shapley value

The **Shapley value**  $\phi(v)$  is the only function  $f: \Gamma^n \rightarrow \mathbb{R}$  satisfying for all games  $(N, v), (N, w)$ :

1. (AXIOM OF EFFICIENCE)
2. (AXIOM OF SYMMETRY)
3. (AXIOM OF NULL PLAYER)
4. (AXIOM OF ADDITIVITY)

# SHAPLEY VALUE

*It is possible to define it using its properties...*

## The Shapley value

The **Shapley value**  $\phi(v)$  is the only function  $f: \Gamma^n \rightarrow \mathbb{R}$  satisfying for all games  $(N, v), (N, w)$ :

1. (AXIOM OF EFFICIENCE)

$$\blacktriangleright \sum_{i \in N} f_i(v) = v(N)$$

2. (AXIOM OF SYMMETRY)

3. (AXIOM OF NULL PLAYER)

4. (AXIOM OF ADDITIVITY)

# SHAPLEY VALUE

*It is possible to define it using its properties...*

## The Shapley value

The **Shapley value**  $\phi(v)$  is the only function  $f: \Gamma^n \rightarrow \mathbb{R}$  satisfying for all games  $(N, v), (N, w)$ :

1. (AXIOM OF EFFICIENCE)

$$\blacktriangleright \sum_{i \in N} f_i(v) = v(N)$$

2. (AXIOM OF SYMMETRY)

$$\blacktriangleright \forall i, j \in N (\forall S \subseteq N \setminus \{i, j\} : v(S \cup i) = v(S \cup j)) \implies f_i(v) = f_j(v)$$

3. (AXIOM OF NULL PLAYER)

4. (AXIOM OF ADDITIVITY)



# SHAPLEY VALUE

*It is possible to define it using its properties...*

## The Shapley value

The **Shapley value**  $\phi(v)$  is the only function  $f: \Gamma^n \rightarrow \mathbb{R}$  satisfying for all games  $(N, v), (N, w)$ :

1. (AXIOM OF EFFICIENCE)

$$\blacktriangleright \sum_{i \in N} f_i(v) = v(N)$$

2. (AXIOM OF SYMMETRY)

$$\blacktriangleright \forall i, j \in N (\forall S \subseteq N \setminus \{i, j\} : v(S \cup i) = v(S \cup j)) \implies f_i(v) = f_j(v)$$

3. (AXIOM OF NULL PLAYER)

$$\blacktriangleright \forall i \in N (\forall S \subseteq N : v(S) = v(S \cup i)) \implies f_i(v) = 0$$

4. (AXIOM OF ADDITIVITY)

# SHAPLEY VALUE

*It is possible to define it using its properties...*

## The Shapley value

The **Shapley value**  $\phi(v)$  is the only function  $f: \Gamma^n \rightarrow \mathbb{R}$  satisfying for all games  $(N, v), (N, w)$ :

1. (AXIOM OF EFFICIENCE)

$$\blacktriangleright \sum_{i \in N} f_i(v) = v(N)$$

2. (AXIOM OF SYMMETRY)

$$\blacktriangleright \forall i, j \in N (\forall S \subseteq N \setminus \{i, j\} : v(S \cup i) = v(S \cup j)) \implies f_i(v) = f_j(v)$$

3. (AXIOM OF NULL PLAYER)

$$\blacktriangleright \forall i \in N (\forall S \subseteq N : v(S) = v(S \cup i)) \implies f_i(v) = 0$$

4. (AXIOM OF ADDITIVITY)

$$\blacktriangleright v, w \in \Gamma^n : f(v + w) = f(v) + f(w)$$



# WEBER SET

1. We fix one construction of  $N$

1. We fix one construction of  $N$ 
  - ▶  $\sigma \in \Sigma_n$  ... represent it by permutation
  - ▶  $\sigma(i)$  ... the order, in which player  $i$  enters the coalition

# WEBER SET

1. We fix one construction of  $N$ 
  - ▶  $\sigma \in \Sigma_n$  ... represent it by permutation
  - ▶  $\sigma(i)$  ... the order, in which player  $i$  enters the coalition
2. We compute player's payments

# WEBER SET

1. We fix one construction of  $N$ 
  - ▶  $\sigma \in \Sigma_n$  ... represent it by permutation
  - ▶  $\sigma(i)$  ... the order, in which player  $i$  enters the coalition
2. We compute player's payments
  - ▶  $m_v^\sigma$  ... **marginal vector**
  - ▶  $(m_v^\sigma)_i = v(S_{\sigma(i)} \cup i) - v(S_{\sigma(i)})$ 
    - $S_{\sigma(i)} = \{j \in N \mid \sigma(j) < \sigma(i)\}$  ... predecessors of  $i$  under  $\sigma$

# WEBER SET

1. We fix one construction of  $N$ 
  - ▶  $\sigma \in \Sigma_n$  ... represent it by permutation
  - ▶  $\sigma(i)$  ... the order, in which player  $i$  enters the coalition
2. We compute player's payments
  - ▶  $m_v^\sigma$  ... **marginal vector**
  - ▶  $(m_v^\sigma)_i = v(S_{\sigma(i)} \cup i) - v(S_{\sigma(i)})$ 
    - $S_{\sigma(i)} = \{j \in N \mid \sigma(j) < \sigma(i)\}$  ... predecessors of  $i$  under  $\sigma$
3. We consider all combinations



# WEBER SET

1. We fix one construction of  $N$ 
  - ▶  $\sigma \in \Sigma_n$  ... represent it by permutation
  - ▶  $\sigma(i)$  ... the order, in which player  $i$  enters the coalition
2. We compute player's payments
  - ▶  $m_v^\sigma$  ... **marginal vector**
  - ▶  $(m_v^\sigma)_i = v(S_{\sigma(i)} \cup i) - v(S_{\sigma(i)})$ 
    - $S_{\sigma(i)} = \{j \in N \mid \sigma(j) < \sigma(i)\}$  ... predecessors of  $i$  under  $\sigma$
3. We consider all combinations
  - ▶  $\mathcal{W}(v) = \text{conv} \{m_v^\sigma \mid \sigma \in \Sigma_n\}$  ... **Webber set**

# WEBER SET

1. We fix one construction of  $N$ 
    - ▶  $\sigma \in \Sigma_n$  ... represent it by permutation
    - ▶  $\sigma(i)$  ... the order, in which player  $i$  enters the coalition
  2. We compute player's payments
    - ▶  $m_v^\sigma$  ... **marginal vector**
    - ▶  $(m_v^\sigma)_i = v(S_{\sigma(i)} \cup i) - v(S_{\sigma(i)})$ 
      - $S_{\sigma(i)} = \{j \in N \mid \sigma(j) < \sigma(i)\}$  ... predecessors of  $i$  under  $\sigma$
  3. We consider all combinations
    - ▶  $\mathcal{W}(v) = \text{conv} \{m_v^\sigma \mid \sigma \in \Sigma_n\}$  ... **Webber set**
- It holds:  $\phi(v) = \sum_{\sigma \in \Sigma_n} \frac{m_v^\sigma}{n!}$

# WEBER SET

1. We fix one construction of  $N$ 
  - ▶  $\sigma \in \Sigma_n$  ... represent it by permutation
  - ▶  $\sigma(i)$  ... the order, in which player  $i$  enters the coalition
2. We compute player's payments
  - ▶  $m_v^\sigma$  ... **marginal vector**
  - ▶  $(m_v^\sigma)_i = v(S_{\sigma(i)} \cup i) - v(S_{\sigma(i)})$ 
    - $S_{\sigma(i)} = \{j \in N \mid \sigma(j) < \sigma(i)\}$  ... predecessors of  $i$  under  $\sigma$
3. We consider all combinations
  - ▶  $\mathcal{W}(v) = \text{conv} \{m_v^\sigma \mid \sigma \in \Sigma_n\}$  ... **Webber set**
  - It holds:  $\phi(v) = \sum_{\sigma \in \Sigma_n} \frac{m_v^\sigma}{n!}$

## Relation between the Shapley value and the Webber set

For a cooperative game  $(N, v)$  it holds:

$$\phi(v) \in \mathcal{W}(v)$$

Moreover,  $\phi(v)$  is the center of gravity of  $\mathcal{W}(v)$ .

**Provable by induction:**

The Weber set contains the core

For every cooperative game  $(N, v)$ , it holds  $\mathcal{C}(v) \subseteq \mathcal{W}(v)$

# CLASSES OF GAMES



## ■ monotonic game

$$(S \subseteq T \subseteq N) (v(S) \leq v(T))$$

# CLASSES OF GAMES

## ■ monotonic game

$$(S \subseteq T \subseteq N) (v(S) \leq v(T))$$

## ■ superadditive game

$$(S, T \subseteq N, S \cap T = \emptyset) (v(S) + v(T) \leq v(S \cup T))$$



# CLASSES OF GAMES

## ■ monotonic game

$$(S \subseteq T \subseteq N) (v(S) \leq v(T))$$

## ■ superadditive game

$$(S, T \subseteq N, S \cap T = \emptyset) (v(S) + v(T) \leq v(S \cup T))$$

## ■ convex game

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

# CLASSES OF GAMES

## ■ monotonic game

$$(S \subseteq T \subseteq N) (v(S) \leq v(T))$$

## ■ superadditive game

$$(S, T \subseteq N, S \cap T = \emptyset) (v(S) + v(T) \leq v(S \cup T))$$

## ■ convex game

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

## ■ essential game

$$v(N) \geq \sum_{i \in N} v(i)$$

# CLASSES OF GAMES

## ■ monotonic game

$$(S \subseteq T \subseteq N) (v(S) \leq v(T))$$

## ■ superadditive game

$$(S, T \subseteq N, S \cap T = \emptyset) (v(S) + v(T) \leq v(S \cup T))$$

## ■ convex game

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

## ■ essential game

$$v(N) \geq \sum_{i \in N} v(i)$$

## ■ balanced game

$$\mathcal{C}(v) \neq \emptyset$$

Question: *What are the relations between them?*

Question: *What are the relations between them?*

Balanced and essential

Balanced cooperative games are essential.

Question: *What are the relations between them?*

Balanced and essential

Balanced cooperative games are essential.

- $(N, v)$  is essential

Question: *What are the relations between them?*

## Balanced and essential

Balanced cooperative games are essential.

■  $(N, v)$  is essential

►  $\mathcal{I}(v) = \{x \in \mathbb{R}^n \mid x(N) = v(N) \wedge \forall i \in N : x_i \geq v(i)\} \neq \emptyset$

Question: *What are the relations between them?*

## Balanced and essential

Balanced cooperative games are essential.

- $(N, v)$  is essential

- ▶  $\mathcal{I}(v) = \{x \in \mathbb{R}^n \mid x(N) = v(N) \wedge \forall i \in N : x_i \geq v(i)\} \neq \emptyset$

- $(N, v)$  is balanced



Question: *What are the relations between them?*

## Balanced and essential

Balanced cooperative games are essential.

- $(N, v)$  is essential

- ▶  $\mathcal{I}(v) = \{x \in \mathbb{R}^n \mid x(N) = v(N) \wedge \forall i \in N : x_i \geq v(i)\} \neq \emptyset$

- $(N, v)$  is balanced

- ▶  $\emptyset \neq \mathcal{C}(v) \subseteq \mathcal{I}(v)$

Question: *What are the relations between them*

Convex and superadditive

Convex cooperative games are superadditive.

Question: *What are the relations between them*

Convex and superadditive

Convex cooperative games are superadditive.

■  $(N, v)$  is convex

►  $(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$

Question: *What are the relations between them*

## Convex and superadditive

Convex cooperative games are superadditive.

- $(N, v)$  is convex
  - ▶  $(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$
- $(N, v)$  is superadditive
  - ▶  $(S, T \subseteq N, S \cap T = \emptyset) (v(S) + v(T) \leq v(S \cup T))$

Question: *What are the relations between them*

## Convex and superadditive

Convex cooperative games are superadditive.

- $(N, v)$  is convex
  - ▶  $(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$
- $(N, v)$  is superadditive
  - ▶  $(S, T \subseteq N, S \cap T = \emptyset) (v(S) + v(T) \leq v(S \cup T))$
- $S \cap T = \emptyset$ 
  - ▶  $(S \cap T) = \mathbf{o}$
  - ▶  $v(S) + v(T) \leq v(S \cup T)$

**TODO: There will be beautiful picture**

# CLASSES OF GAMES

## Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

# CLASSES OF GAMES

## Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

## Core of convex games



# CLASSES OF GAMES

## Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

## Core of convex games

For a convex cooperative game  $(N, v)$ , it holds  $\mathcal{C}(v) = \mathcal{W}(v)$

# CLASSES OF GAMES

## Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

## Core of convex games

For a convex cooperative game  $(N, v)$ , it holds  $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only  $m_v^{id} \in \mathcal{C}(v)$

# CLASSES OF GAMES

## Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

## Core of convex games

For a convex cooperative game  $(N, v)$ , it holds  $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only  $m_v^{id} \in \mathcal{C}(v)$

$$1. m_v^{id}(N) = \sum_{i \in N} v(\{1, \dots, i\}) - v(\{1, \dots, i-1\}) = V(N)$$

# CLASSES OF GAMES

## Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

## Core of convex games

For a convex cooperative game  $(N, v)$ , it holds  $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only  $m_v^{id} \in \mathcal{C}(v)$

1.  $m_v^{id}(N) = \sum_{i \in N} v(\{1, \dots, i\}) - v(\{1, \dots, i-1\}) = V(N)$
2.  $m_v^{id}(S) \geq v(S)$

# CLASSES OF GAMES

## Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

## Core of convex games

For a convex cooperative game  $(N, v)$ , it holds  $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only  $m_v^{id} \in \mathcal{C}(v)$

1.  $m_v^{id}(N) = \sum_{i \in N} v(\{1, \dots, i\}) - v(\{1, \dots, i-1\}) = v(N)$
2.  $m_v^{id}(S) \geq v(S)$ 
  - $S = \{s_1, s_2, \dots, s_k\}$ 
    - $s_1 < s_2 < \dots < s_k$

# CLASSES OF GAMES

## Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

## Core of convex games

For a convex cooperative game  $(N, v)$ , it holds  $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only  $m_v^{id} \in \mathcal{C}(v)$

1.  $m_v^{id}(N) = \sum_{i \in N} v(\{1, \dots, i\}) - v(\{1, \dots, i-1\}) = V(N)$
2.  $m_v^{id}(S) \geq v(S)$ 
  - ▶  $S = \{s_1, s_2, \dots, s_k\}$ 
    - $s_1 < s_2 < \dots < s_k$
  - ▶  $v(\{1, \dots, s_i\}) - v(\{1, \dots, s_{i-1}\}) \geq v(\{s_1, \dots, s_i\}) - v(\{s_1, \dots, s_{i-1}\})$ 
    - $\{s_1, s_2, \dots, s_l\} \subseteq \{1, \dots, s_l\}$  for  $l \leq k$

# CLASSES OF GAMES

## Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

## Core of convex games

For a convex cooperative game  $(N, v)$ , it holds  $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only  $m_v^{id} \in \mathcal{C}(v)$

1.  $m_v^{id}(N) = \sum_{i \in N} v(\{1, \dots, i\}) - v(\{1, \dots, i-1\}) = V(N)$
2.  $m_v^{id}(S) \geq v(S)$ 
  - ▶  $S = \{s_1, s_2, \dots, s_k\}$ 
    - $s_1 < s_2 < \dots < s_k$
  - ▶  $v(\{1, \dots, s_i\}) - v(\{1, \dots, s_{i-1}\}) \geq v(\{s_1, \dots, s_i\}) - v(\{s_1, \dots, s_{i-1}\})$ 
    - $\{s_1, s_2, \dots, s_l\} \subseteq \{1, \dots, s_l\}$  for  $l \leq k$
  - ▶  $m_v^{id}(S) = \sum_{s_i \in S} v(\{1, \dots, s_i\}) - v(\{1, \dots, s_{i-1}\})$

# CLASSES OF GAMES

## Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

## Core of convex games

For a convex cooperative game  $(N, v)$ , it holds  $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only  $m_v^{id} \in \mathcal{C}(v)$

1.  $m_v^{id}(N) = \sum_{i \in N} v(\{1, \dots, i\}) - v(\{1, \dots, i-1\}) = v(N)$

2.  $m_v^{id}(S) \geq v(S)$

▶  $S = \{s_1, s_2, \dots, s_k\}$

■  $s_1 < s_2 < \dots < s_k$

▶  $v(\{1, \dots, s_i\}) - v(\{1, \dots, s_{i-1}\}) \geq v(\{s_1, \dots, s_i\}) - v(\{s_1, \dots, s_{i-1}\})$

■  $\{s_1, s_2, \dots, s_l\} \subseteq \{1, \dots, s_l\}$  for  $l \leq k$

▶  $m_v^{id}(S) = \sum_{s_i \in S} v(\{1, \dots, s_i\}) - v(\{1, \dots, s_{i-1}\})$

▶  $m_v^{id} \geq \sum_{s_i \in S} v(\{s_1, \dots, s_i\}) - v(\{s_1, \dots, s_{i-1}\}) = v(S)$



## Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

Consequence:

## The Shapley value and convex games

## Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

Consequence:

## The Shapley value and convex games

For a convex cooperative game  $(N, v)$ , it holds:

1.  $\phi(v) \in \mathcal{C}(v)$
2.  $\phi(v)$  is the centre of gravity of  $\mathcal{C}(v)$

# **DISADVANTAGES OF THE MODEL AND MOTIVATION FOR NEXT LECTURES**

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

1. we might not be interested in forming  $N$

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

1. **we might not be interested in forming  $N$** 
  - we want to analyze coalition formation

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

1. **we might not be interested in forming  $N$** 
  - ▶ we want to analyze coalition formation
2. **not every coalition makes sense**

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

1. **we might not be interested in forming  $N$**

- ▶ we want to analyze coalition formation

2. **not every coalition makes sense**

- ▶ two players from completely different fields (why would they want to cooperate)



## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

1. **we might not be interested in forming  $N$** 
  - ▶ we want to analyze coalition formation
2. **not every coalition makes sense**
  - ▶ two players from completely different fields (why would they want to cooperate)
3.  **$2^n$  real values representing  $v$**

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

1. **we might not be interested in forming  $N$**

- ▶ we want to analyze coalition formation

2. **not every coalition makes sense**

- ▶ two players from completely different fields (why would they want to cooperate)

3.  **$2^n$  real values representing  $v$**

- ▶ expensive to get all information
- ▶ expensive to store all information

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

1. **we might not be interested in forming  $N$**
2. **not every coalition makes sense**
3.  **$2^n$  real values representing  $v$**

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

1. **we might not be interested in forming  $N$** 
  - games with coalition structure
2. **not every coalition makes sense**
3.  **$2^n$  real values representing  $v$**

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

1. **we might not be interested in forming  $N$** 
  - ▶ games with coalition structure
2. **not every coalition makes sense**
  - ▶ games on graphs (Martin Černý)
3.  **$2^n$  real values representing  $v$**

## Cooperative game

A **cooperative game** is an ordered pair  $(N, v)$ , where  $N$  is a set of players and  $v: 2^N \rightarrow \mathbb{R}$  is the characteristic function. Further,  $v(\emptyset) = 0$ .

1. **we might not be interested in forming  $N$** 
  - ▶ games with coalition structure
2. **not every coalition makes sense**
  - ▶ games on graphs (Martin Černý)
3.  **$2^n$  real values representing  $v$** 
  - ▶ incomplete games (David Sychrovský + Filip Úradník)
  - ▶ interval games (Martin Kunst)
  - ▶ stochastic games (David Ryzák)
  - ▶ models with compact characteristic function (Martin Černý)