

COOPERATIVE GAME THEORY

RICHARD MUŽÍK

RICHARD@IMUZYK.CZ

APRIL 21, 2023

INTRODUCTION

Cooperative game

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

■ $S \subseteq N$... coalition

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

- $S \subseteq N$... coalition
- $v(S)$... value of coalition

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

- Γ^n ... set of n -person cooperative games
- $S \subseteq N$... coalition
- $v(S)$... value of coalition

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

- Γ^n ... set of n -person cooperative games
- $S \subseteq N$... coalition
- $v(S)$... value of coalition
- usually $N = \{1, \dots, n\}$

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

- Γ^n ... set of n -person cooperative games
- $S \subseteq N$... coalition
- $v(S)$... value of coalition
- usually $N = \{1, \dots, n\}$
 - ▶ (S, v_S) is **subgame** (N, v) :
 - $v_S: 2^S \rightarrow \mathbb{R}$
 - $v_S(T) := v(T)$ pro $T \subseteq S$

COOPERATION - EXAMPLES OF MODELS: *MINIMAL SPANNING-TREE GAMES*

Goal: *Find the best connection of players to a source*

COOPERATION - EXAMPLES OF MODELS: *MINIMAL SPANNING-TREE GAMES*

Goal: *Find the best connection of players to a source*

- $N = N' \cup \{o\}$... set of players + source

COOPERATION - EXAMPLES OF MODELS:

MINIMAL SPANNING-TREE GAMES

Goal: *Find the best connection of players to a source*

- $N = N' \cup \{o\}$... set of players + source
- c_{ij} ... cost of connecting i, j

COOPERATION - EXAMPLES OF MODELS:

MINIMAL SPANNING-TREE GAMES

Goal: *Find the best connection of players to a source*

- $N = N' \cup \{o\}$... set of players + source
- c_{ij} ... cost of connecting i, j
- solution: a network, where each $i \in N$ is connected to o with minimal sum of costs

GOAL OF THE MODEL OF COOPERATIVE GAMES

Money first!

GOAL OF THE MODEL OF COOPERATIVE GAMES

Money first!

- **Payoff vector $\mathbf{x} \in \mathbb{R}^n$**
 - ▶ x_i represents payoff of player i

GOAL OF THE MODEL OF COOPERATIVE GAMES

Money first!

- **Payoff vector** $\mathbf{x} \in \mathbb{R}^n$
 - ▶ x_i represents payoff of player i
- Vector $\mathbf{x} \in \mathbb{R}^n$ is **efficient**, if $\sum_{i \in N} x_i = v(N)$

GOAL OF THE MODEL OF COOPERATIVE GAMES

Money first!

- **Payoff vector** $\mathbf{x} \in \mathbb{R}^n$
 - ▶ x_i represents payoff of player i
- Vector $\mathbf{x} \in \mathbb{R}^n$ is **efficient**, if $\sum_{i \in N} x_i = v(N)$
 - ▶ Usually, we distribute $v(N)$
 1. *value of cooperation* $v(N)$
 2. *shared costs* $c(N)$

GOAL OF THE MODEL OF COOPERATIVE GAMES

Money first!

- **Payoff vector** $\mathbf{x} \in \mathbb{R}^n$
 - ▶ x_i represents payoff of player i
- Vector $\mathbf{x} \in \mathbb{R}^n$ is **efficient**, if $\sum_{i \in N} x_i = v(N)$
 - ▶ Usually, we distribute $v(N)$
 1. *value of cooperation* $v(N)$
 2. *shared costs* $c(N)$
- Vector $\mathbf{x} \in \mathbb{R}^n$ is **individually rational**, if $x_i \geq v(i)$

GOAL OF THE MODEL OF COOPERATIVE GAMES

Money first!

- **Payoff vector** $\mathbf{x} \in \mathbb{R}^n$
 - ▶ x_i represents payoff of player i
- Vector $\mathbf{x} \in \mathbb{R}^n$ is **efficient**, if $\sum_{i \in N} x_i = v(N)$
 - ▶ Usually, we distribute $v(N)$
 1. *value of cooperation* $v(N)$
 2. *shared costs* $c(N)$
- Vector $\mathbf{x} \in \mathbb{R}^n$ is **individually rational**, if $x_i \geq v(i)$
 - ▶ players prefer x_i over $v(i)$

GOAL OF THE MODEL OF COOPERATIVE GAMES

Money first!

- **Payoff vector** $\mathbf{x} \in \mathbb{R}^n$
 - ▶ x_i represents payoff of player i
- Vector $\mathbf{x} \in \mathbb{R}^n$ is **efficient**, if $\sum_{i \in N} x_i = v(N)$
 - ▶ Usually, we distribute $v(N)$
 1. *value of cooperation* $v(N)$
 2. *shared costs* $c(N)$
- Vector $\mathbf{x} \in \mathbb{R}^n$ is **individually rational**, if $x_i \geq v(i)$
 - ▶ players prefer x_i over $v(i)$
- $\mathcal{I}^*(v) = \{\mathbf{x} \in \mathbb{R}^n \mid \mathbf{x}(N) = v(N)\}$... **preimputation**
 - ▶ $\mathbf{x}(S) := \sum_{i \in S} x_i$

GOAL OF THE MODEL OF COOPERATIVE GAMES

Money first!

- **Payoff vector** $\mathbf{x} \in \mathbb{R}^n$
 - ▶ x_i represents payoff of player i
- Vector $\mathbf{x} \in \mathbb{R}^n$ is **efficient**, if $\sum_{i \in N} x_i = v(N)$
 - ▶ Usually, we distribute $v(N)$
 1. *value of cooperation* $v(N)$
 2. *shared costs* $c(N)$
- Vector $\mathbf{x} \in \mathbb{R}^n$ is **individually rational**, if $x_i \geq v(i)$
 - ▶ players prefer x_i over $v(i)$
- $\mathcal{I}^*(v) = \{x \in \mathbb{R}^n \mid x(N) = v(N)\}$... **preimputation**
 - ▶ $x(S) := \sum_{i \in S} x_i$
- $\mathcal{I}(v) = \{x \in \mathcal{I}^*(v) \mid \forall i \in N : x_i \geq v(i)\}$... **imputation**

SOLUTION CONCEPTS

- Set of payoff vectors satisfying further properties are **solution concepts**

- Set of payoff vectors satisfying further properties are **solution concepts**
- Can reflect payoff distribution, which is

- Set of payoff vectors satisfying further properties are **solution concepts**
- Can reflect payoff distribution, which is
 - ▶ ...*fair*...
 - ▶ ...*non-discriminatory*...
 - ▶ ...*stable (players will accept it)*...
 - ▶ ...

Formally:

Formally:

1. sets of payoff vectors

Formally:

1. sets of payoff vectors

► $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$

Formally:

1. sets of payoff vectors
 - ▶ $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$
2. functions on games

Formally:

1. sets of payoff vectors

▶ $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$

2. functions on games

▶ $\Sigma: \Gamma^n \rightarrow 2^{\mathbb{R}^n}$

Formally:

1. sets of payoff vectors

▶ $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$

2. functions on games

▶ $\Sigma: \Gamma^n \rightarrow 2^{\mathbb{R}^n}$

We distinguish

Formally:

1. sets of payoff vectors
 - ▶ $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$
2. functions on games
 - ▶ $\Sigma: \Gamma^n \rightarrow 2^{\mathbb{R}^n}$

We distinguish

1. **single-point** solution concepts
2. **multi-point** solution concepts

Formally:

1. sets of payoff vectors
 - ▶ $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$
2. functions on games
 - ▶ $\Sigma: \Gamma^n \rightarrow 2^{\mathbb{R}^n}$

We distinguish

1. **single-point** solution concepts
 - ▶ as a set: $\Sigma(v) = \{x\}$
 - we prefer: $\Sigma(v) = x$
2. **multi-point** solution concepts

Formally:

1. sets of payoff vectors

▶ $\Sigma(v) = \{x \in \mathbb{R}^n \mid \dots\}$

2. functions on games

▶ $\Sigma: \Gamma^n \rightarrow 2^{\mathbb{R}^n}$

We distinguish

1. **single-point** solution concepts

▶ as a set: $\Sigma(v) = \{x\}$

■ we prefer: $\Sigma(v) = x$

▶ as a function: $\Sigma: \Gamma^n \rightarrow \mathbb{R}$

2. **multi-point** solution concepts

MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

The core

MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

■ assumption: *homo economicus*

MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

■ assumption: *homo economicus*

- ▶ model of human as a player
- ▶ strictly rational and selfish
- ▶ follows his subjective goals

MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

- assumption: *homo economicus*
 - ▶ model of human as a player
 - ▶ strictly rational and selfish
 - ▶ follows his subjective goals
- $v(N)$... value, which is distributed among players

MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

- assumption: *homo economicus*
 - ▶ model of human as a player
 - ▶ strictly rational and selfish
 - ▶ follows his subjective goals
- $v(N)$... value, which is distributed among players
- $x(S) > v(S) \implies$ coalition S does not leave N

MULTI-POINT SOLUTION CONCEPT: THE CORE

Idea: *Payoff distribution leads to cooperation...*

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

- assumption: *homo economicus*
 - ▶ model of human as a player
 - ▶ strictly rational and selfish
 - ▶ follows his subjective goals
- $v(N)$... value, which is distributed among players
- $x(S) > v(S) \implies$ coalition S does not leave N
 - ▶ would lead to (S, v_S)
 - ▶ $v(S)$... distributed value

MULTI-POINT SOLUTION CONCEPT: THE CORE

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

MULTI-POINT SOLUTION CONCEPT: THE CORE

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

Reminder

Nash equilibrium

MULTI-POINT SOLUTION CONCEPT: THE CORE

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

Reminder

Nash equilibrium

Strategy profile (s_1, \dots, s_n) is **Nash equilibrium**, if it holds for every player i ,

$$v_i(s_1, \dots, s_{i-1}, s_i, s_{i+1}, \dots, s_n) \geq v_i(s_1, \dots, s_{i-1}, t_i, s_{i+1}, \dots, s_n)$$

for every $t_i \in S_i$.

MULTI-POINT SOLUTION CONCEPT: THE CORE

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

Emptiness of the core

MULTI-POINT SOLUTION CONCEPT: THE CORE

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

Emptiness of the core

There are cooperative games (N, v) with empty core.

MULTI-POINT SOLUTION CONCEPT: THE CORE

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

Emptiness of the core

There are cooperative games (N, v) with empty core.

- Non-essential games

MULTI-POINT SOLUTION CONCEPT: THE CORE

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

Emptiness of the core

There are cooperative games (N, v) with empty core.

- Non-essential games

- $v(N) < \sum_{i \in N} v(i)$

MULTI-POINT SOLUTION CONCEPT: THE CORE

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

Emptiness of the core

There are cooperative games (N, v) with empty core.

- Non-essential games

- $v(N) < \sum_{i \in N} v(i)$

- Let $x \in \mathcal{C}(v)$

MULTI-POINT SOLUTION CONCEPT: THE CORE

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

Emptiness of the core

There are cooperative games (N, v) with empty core.

- Non-essential games

- ▶ $v(N) < \sum_{i \in N} v(i)$

- Let $x \in \mathcal{C}(v)$

- ▶ $x(N) = v(N) < \sum_{i \in N} x(i) \leq x(N)$

MULTI-POINT SOLUTION CONCEPT: THE CORE

The core

For a cooperative game (N, v) , the **core** $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

Emptiness of the core

There are cooperative games (N, v) with empty core.

- Non-essential games
 - ▶ $v(N) < \sum_{i \in N} v(i)$
- Let $x \in \mathcal{C}(v)$
 - ▶ $x(N) = v(N) < \sum_{i \in N} x(i) \leq x(N)$
- x does not exist $\implies \mathcal{C}(v) = \emptyset$

MULTI-POINT SOLUTION CONCEPT: THE CORE

Question: *When is the core non-empty?*

MULTI-POINT SOLUTION CONCEPT: THE CORE

Question: *When is the core non-empty?*

We can encode the core as linear program (P) and determine the dual program (D)

MULTI-POINT SOLUTION CONCEPT: THE CORE

Question: *When is the core non-empty?*

We can encode the core as linear program (P) and determine the dual program (D)

$$\blacksquare (P) = \begin{cases} \min_{x \in \mathbb{R}^n} & x(N) \\ \text{subject to} & x(S) \geq v(S) \text{ for } S \subseteq N \end{cases}$$

MULTI-POINT SOLUTION CONCEPT: THE CORE

Question: *When is the core non-empty?*

We can encode the core as linear program (P) and determine the dual program (D)

$$\blacksquare (P) = \begin{cases} \min_{x \in \mathbb{R}^n} & x(N) \\ \text{subject to} & x(S) \geq v(S) \text{ for } S \subseteq N \end{cases}$$

$$\blacksquare (D) = \begin{cases} \max_{y \in \mathbb{R}_+^{(2^n-1)}} & \sum_{S \subseteq N} y_S v(S) \\ \text{subject to} & \sum_{\emptyset \neq S \subseteq N} y_S \chi_S = \chi_N \end{cases}$$

MULTI-POINT SOLUTION CONCEPT: THE CORE

Question: *When is the core non-empty?*

We can encode the core as linear program (P) and determine the dual program (D)

$$\begin{aligned} \blacksquare (P) &= \begin{cases} \min_{x \in \mathbb{R}^n} & x(N) \\ \text{subject to} & x(S) \geq v(S) \text{ for } S \subseteq N \end{cases} \\ \blacksquare (D) &= \begin{cases} \max_{y \in \mathbb{R}_+^{(2^n-1)}} & \sum_{S \subseteq N} y_S v(S) \\ \text{subject to} & \sum_{\emptyset \neq S \subseteq N} y_S \chi_S = \chi_N \end{cases} \end{aligned}$$

We derive the weak Bondareva-Shapley theorem

Weak Bondareva-Shapley theorem

Cooperative game (N, v) has non-empty core if and only if

$$v(N) \geq \sum_{S \subseteq N} y_S v(S) \text{ for all feasible } y \in \mathbb{R}^{(2^n-1)}$$

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

■ $v(S \cup i) - v(S)$

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- $v(S \cup i) - v(S)$
 - *marginal contribution* (of player i in $S \cup i$)

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- $v(S \cup i) - v(S)$
 - ▶ *marginal contribution* (of player i in $S \cup i$)
- $\frac{s!(n-s-1)!}{n!}$

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- $v(S \cup i) - v(S)$
 - ▶ *marginal contribution* (of player i in $S \cup i$)
- $\frac{s!(n-s-1)!}{n!}$
 - ▶ weights reflecting different sizes of coalitions

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- $v(S \cup i) - v(S)$
 - ▶ *marginal contribution* (of player i in $S \cup i$)
- $\frac{s!(n-s-1)!}{n!}$
 - ▶ weights reflecting different sizes of coalitions
- $\sum_{S \subseteq N \setminus i}$

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- $v(S \cup i) - v(S)$
 - ▶ *marginal contribution* (of player i in $S \cup i$)
- $\frac{s!(n-s-1)!}{n!}$
 - ▶ weights reflecting different sizes of coalitions
- $\sum_{S \subseteq N \setminus i}$
 - ▶ sum of all marginal contributions of i

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:
 1. Form the grandcoalition N .

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:
 1. Form the grandcoalition N .
 2. Enter the the coalition individually and randomly.

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:
 1. Form the grandcoalition N .
 2. Enter the the coalition individually and randomly.
 3. When player i enters coalition S , he receives $v(S \cup i) - v(S)$.

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:
 1. Form the grandcoalition N .
 2. Enter the the coalition individually and randomly.
 3. When player i enters coalition S , he receives $v(S \cup i) - v(S)$.
- $s!(n-s-1)!$... number of situations, in which i enters S

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:
 1. Form the grandcoalition N .
 2. Enter the the coalition individually and randomly.
 3. When player i enters coalition S , he receives $v(S \cup i) - v(S)$.
- $s!(n-s-1)!$... number of situations, in which i enters S
- $n!$... number of all possible ways to construct N

SINGLE-POINT SOLUTION CONCEPT: THE SHAPLEY VALUE

Idea: *Divide the profit in a fair way...*

The Shapley value

For a cooperative game (N, v) , the **Shapley value** $\phi(v)$ of player i is

$$\phi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

- The players agree on the following procedure:
 1. Form the grandcoalition N .
 2. Enter the the coalition individually and randomly.
 3. When player i enters coalition S , he receives $v(S \cup i) - v(S)$.
- $s!(n-s-1)!$... number of situations, in which i enters S
- $n!$... number of all possible ways to construct N
- $\phi_i(v)$... the average value of player i 's payment

SHAPLEY VALUE

It is possible to define it using its properties...

SHAPLEY VALUE

It is possible to define it using its properties...

The Shapley value

The **Shapley value** $\phi(v)$ is the only function $f: \Gamma^n \rightarrow \mathbb{R}$ satisfying for all games $(N, v), (N, w)$:

SHAPLEY VALUE

It is possible to define it using its properties...

The Shapley value

The **Shapley value** $\phi(v)$ is the only function $f: \Gamma^n \rightarrow \mathbb{R}$ satisfying for all games $(N, v), (N, w)$:

1. (AXIOM OF EFFICIENCE)
2. (AXIOM OF SYMMETRY)
3. (AXIOM OF NULL PLAYER)
4. (AXIOM OF ADDITIVITY)

SHAPLEY VALUE

It is possible to define it using its properties...

The Shapley value

The **Shapley value** $\phi(v)$ is the only function $f: \Gamma^n \rightarrow \mathbb{R}$ satisfying for all games $(N, v), (N, w)$:

1. (AXIOM OF EFFICIENCE)

$$\blacktriangleright \sum_{i \in N} f_i(v) = v(N)$$

2. (AXIOM OF SYMMETRY)

3. (AXIOM OF NULL PLAYER)

4. (AXIOM OF ADDITIVITY)

SHAPLEY VALUE

It is possible to define it using its properties...

The Shapley value

The **Shapley value** $\phi(v)$ is the only function $f: \Gamma^n \rightarrow \mathbb{R}$ satisfying for all games $(N, v), (N, w)$:

1. (AXIOM OF EFFICIENCE)

$$\blacktriangleright \sum_{i \in N} f_i(v) = v(N)$$

2. (AXIOM OF SYMMETRY)

$$\blacktriangleright \forall i, j \in N (\forall S \subseteq N \setminus \{i, j\} : v(S \cup i) = v(S \cup j)) \implies f_i(v) = f_j(v)$$

3. (AXIOM OF NULL PLAYER)

4. (AXIOM OF ADDITIVITY)

SHAPLEY VALUE

It is possible to define it using its properties...

The Shapley value

The **Shapley value** $\phi(v)$ is the only function $f: \Gamma^n \rightarrow \mathbb{R}$ satisfying for all games $(N, v), (N, w)$:

1. (AXIOM OF EFFICIENCE)

$$\blacktriangleright \sum_{i \in N} f_i(v) = v(N)$$

2. (AXIOM OF SYMMETRY)

$$\blacktriangleright \forall i, j \in N (\forall S \subseteq N \setminus \{i, j\} : v(S \cup i) = v(S \cup j)) \implies f_i(v) = f_j(v)$$

3. (AXIOM OF NULL PLAYER)

$$\blacktriangleright \forall i \in N (\forall S \subseteq N : v(S) = v(S \cup i)) \implies f_i(v) = 0$$

4. (AXIOM OF ADDITIVITY)

SHAPLEY VALUE

It is possible to define it using its properties...

The Shapley value

The **Shapley value** $\phi(v)$ is the only function $f: \Gamma^n \rightarrow \mathbb{R}$ satisfying for all games $(N, v), (N, w)$:

1. (AXIOM OF EFFICIENCE)

$$\blacktriangleright \sum_{i \in N} f_i(v) = v(N)$$

2. (AXIOM OF SYMMETRY)

$$\blacktriangleright \forall i, j \in N (\forall S \subseteq N \setminus \{i, j\} : v(S \cup i) = v(S \cup j)) \implies f_i(v) = f_j(v)$$

3. (AXIOM OF NULL PLAYER)

$$\blacktriangleright \forall i \in N (\forall S \subseteq N : v(S) = v(S \cup i)) \implies f_i(v) = 0$$

4. (AXIOM OF ADDITIVITY)

$$\blacktriangleright v, w \in \Gamma^n : f(v + w) = f(v) + f(w)$$

WEBER SET

1. We fix one construction of N

1. We fix one construction of N
 - ▶ $\sigma \in \Sigma_n$... represent it by permutation
 - ▶ $\sigma(i)$... the order, in which player i enters the coalition

WEBER SET

1. We fix one construction of N
 - ▶ $\sigma \in \Sigma_n$... represent it by permutation
 - ▶ $\sigma(i)$... the order, in which player i enters the coalition
2. We compute player's payments

WEBER SET

1. We fix one construction of N
 - ▶ $\sigma \in \Sigma_n$... represent it by permutation
 - ▶ $\sigma(i)$... the order, in which player i enters the coalition
2. We compute player's payments
 - ▶ m_v^σ ... **marginal vector**
 - ▶ $(m_v^\sigma)_i = v(S_{\sigma(i)} \cup i) - v(S_{\sigma(i)})$
 - $S_{\sigma(i)} = \{j \in N \mid \sigma(j) < \sigma(i)\}$... predecessors of i under σ

WEBER SET

1. We fix one construction of N
 - ▶ $\sigma \in \Sigma_n$... represent it by permutation
 - ▶ $\sigma(i)$... the order, in which player i enters the coalition
2. We compute player's payments
 - ▶ m_v^σ ... **marginal vector**
 - ▶ $(m_v^\sigma)_i = v(S_{\sigma(i)} \cup i) - v(S_{\sigma(i)})$
 - $S_{\sigma(i)} = \{j \in N \mid \sigma(j) < \sigma(i)\}$... predecessors of i under σ
3. We consider all combinations

WEBER SET

1. We fix one construction of N
 - ▶ $\sigma \in \Sigma_n$... represent it by permutation
 - ▶ $\sigma(i)$... the order, in which player i enters the coalition
2. We compute player's payments
 - ▶ m_v^σ ... **marginal vector**
 - ▶ $(m_v^\sigma)_i = v(S_{\sigma(i)} \cup i) - v(S_{\sigma(i)})$
 - $S_{\sigma(i)} = \{j \in N \mid \sigma(j) < \sigma(i)\}$... predecessors of i under σ
3. We consider all combinations
 - ▶ $\mathcal{W}(v) = \text{conv} \{m_v^\sigma \mid \sigma \in \Sigma_n\}$... **Webber set**

WEBER SET

1. We fix one construction of N
 - ▶ $\sigma \in \Sigma_n$... represent it by permutation
 - ▶ $\sigma(i)$... the order, in which player i enters the coalition
 2. We compute player's payments
 - ▶ m_v^σ ... **marginal vector**
 - ▶ $(m_v^\sigma)_i = v(S_{\sigma(i)} \cup i) - v(S_{\sigma(i)})$
 - $S_{\sigma(i)} = \{j \in N \mid \sigma(j) < \sigma(i)\}$... predecessors of i under σ
 3. We consider all combinations
 - ▶ $\mathcal{W}(v) = \text{conv} \{m_v^\sigma \mid \sigma \in \Sigma_n\}$... **Webber set**
- It holds: $\phi(v) = \sum_{\sigma \in \Sigma_n} \frac{m_v^\sigma}{n!}$

WEBER SET

1. We fix one construction of N
 - ▶ $\sigma \in \Sigma_n$... represent it by permutation
 - ▶ $\sigma(i)$... the order, in which player i enters the coalition
2. We compute player's payments
 - ▶ m_v^σ ... **marginal vector**
 - ▶ $(m_v^\sigma)_i = v(S_{\sigma(i)} \cup i) - v(S_{\sigma(i)})$
 - $S_{\sigma(i)} = \{j \in N \mid \sigma(j) < \sigma(i)\}$... predecessors of i under σ
3. We consider all combinations
 - ▶ $\mathcal{W}(v) = \text{conv} \{m_v^\sigma \mid \sigma \in \Sigma_n\}$... **Webber set**
 - It holds: $\phi(v) = \sum_{\sigma \in \Sigma_n} \frac{m_v^\sigma}{n!}$

Relation between the Shapley value and the Webber set

For a cooperative game (N, v) it holds:

$$\phi(v) \in \mathcal{W}(v)$$

Moreover, $\phi(v)$ is the center of gravity of $\mathcal{W}(v)$.

Provable by induction:

The Weber set contains the core

For every cooperative game (N, v) , it holds $\mathcal{C}(v) \subseteq \mathcal{W}(v)$

CLASSES OF GAMES

■ monotonic game

$$(S \subseteq T \subseteq N) (v(S) \leq v(T))$$

CLASSES OF GAMES

■ monotonic game

$$(S \subseteq T \subseteq N) (v(S) \leq v(T))$$

■ superadditive game

$$(S, T \subseteq N, S \cap T = \emptyset) (v(S) + v(T) \leq v(S \cup T))$$

CLASSES OF GAMES

■ monotonic game

$$(S \subseteq T \subseteq N) (v(S) \leq v(T))$$

■ superadditive game

$$(S, T \subseteq N, S \cap T = \emptyset) (v(S) + v(T) \leq v(S \cup T))$$

■ convex game

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

CLASSES OF GAMES

■ monotonic game

$$(S \subseteq T \subseteq N) (v(S) \leq v(T))$$

■ superadditive game

$$(S, T \subseteq N, S \cap T = \emptyset) (v(S) + v(T) \leq v(S \cup T))$$

■ convex game

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

■ essential game

$$v(N) \geq \sum_{i \in N} v(i)$$

CLASSES OF GAMES

■ monotonic game

$$(S \subseteq T \subseteq N) (v(S) \leq v(T))$$

■ superadditive game

$$(S, T \subseteq N, S \cap T = \emptyset) (v(S) + v(T) \leq v(S \cup T))$$

■ convex game

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

■ essential game

$$v(N) \geq \sum_{i \in N} v(i)$$

■ balanced game

$$\mathcal{C}(v) \neq \emptyset$$

Question: *What are the relations between them?*

Question: *What are the relations between them?*

Balanced and essential

Balanced cooperative games are essential.

Question: *What are the relations between them?*

Balanced and essential

Balanced cooperative games are essential.

- (N, v) is essential

Question: *What are the relations between them?*

Balanced and essential

Balanced cooperative games are essential.

■ (N, v) is essential

► $\mathcal{I}(v) = \{x \in \mathbb{R}^n \mid x(N) = v(N) \wedge \forall i \in N : x_i \geq v(i)\} \neq \emptyset$

Question: *What are the relations between them?*

Balanced and essential

Balanced cooperative games are essential.

■ (N, v) is essential

► $\mathcal{I}(v) = \{x \in \mathbb{R}^n \mid x(N) = v(N) \wedge \forall i \in N : x_i \geq v(i)\} \neq \emptyset$

■ (N, v) is balanced

Question: *What are the relations between them?*

Balanced and essential

Balanced cooperative games are essential.

- (N, v) is essential

- ▶ $\mathcal{I}(v) = \{x \in \mathbb{R}^n \mid x(N) = v(N) \wedge \forall i \in N : x_i \geq v(i)\} \neq \emptyset$

- (N, v) is balanced

- ▶ $\emptyset \neq \mathcal{C}(v) \subseteq \mathcal{I}(v)$

Question: *What are the relations between them*

Convex and superadditive

Convex cooperative games are superadditive.

Question: *What are the relations between them*

Convex and superadditive

Convex cooperative games are superadditive.

■ (N, v) is convex

► $(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$

Question: *What are the relations between them*

Convex and superadditive

Convex cooperative games are superadditive.

- (N, v) is convex
 - ▶ $(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$
- (N, v) is superadditive
 - ▶ $(S, T \subseteq N, S \cap T = \emptyset) (v(S) + v(T) \leq v(S \cup T))$


Question: *What are the relations between them*

Convex and superadditive

Convex cooperative games are superadditive.

- (N, v) is convex
 - ▶ $(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$
- (N, v) is superadditive
 - ▶ $(S, T \subseteq N, S \cap T = \emptyset) (v(S) + v(T) \leq v(S \cup T))$
- $S \cap T = \emptyset$
 - ▶ $(S \cap T) = \mathbf{o}$
 - ▶ $v(S) + v(T) \leq v(S \cup T)$

CLASSES OF GAMES



CLASSES OF GAMES

Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

CLASSES OF GAMES

Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

Core of convex games

CLASSES OF GAMES

Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

Core of convex games

For a convex cooperative game (N, v) , it holds $\mathcal{C}(v) = \mathcal{W}(v)$

CLASSES OF GAMES

Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

Core of convex games

For a convex cooperative game (N, v) , it holds $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only $m_v^{id} \in \mathcal{C}(v)$

CLASSES OF GAMES

Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

Core of convex games

For a convex cooperative game (N, v) , it holds $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only $m_v^{id} \in \mathcal{C}(v)$

$$1. m_v^{id}(N) = \sum_{i \in N} v(\{1, \dots, i\}) - v(\{1, \dots, i-1\}) = V(N)$$

CLASSES OF GAMES

Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

Core of convex games

For a convex cooperative game (N, v) , it holds $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only $m_v^{id} \in \mathcal{C}(v)$

1. $m_v^{id}(N) = \sum_{i \in N} v(\{1, \dots, i\}) - v(\{1, \dots, i-1\}) = V(N)$
2. $m_v^{id}(S) \geq v(S)$

CLASSES OF GAMES

Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

Core of convex games

For a convex cooperative game (N, v) , it holds $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only $m_v^{id} \in \mathcal{C}(v)$

1. $m_v^{id}(N) = \sum_{i \in N} v(\{1, \dots, i\}) - v(\{1, \dots, i-1\}) = V(N)$
2. $m_v^{id}(S) \geq v(S)$
 - $S = \{s_1, s_2, \dots, s_k\}$
 - $s_1 < s_2 < \dots < s_k$

CLASSES OF GAMES

Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

Core of convex games

For a convex cooperative game (N, v) , it holds $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only $m_v^{id} \in \mathcal{C}(v)$

1. $m_v^{id}(N) = \sum_{i \in N} v(\{1, \dots, i\}) - v(\{1, \dots, i-1\}) = V(N)$
2. $m_v^{id}(S) \geq v(S)$
 - ▶ $S = \{s_1, s_2, \dots, s_k\}$
 - $s_1 < s_2 < \dots < s_k$
 - ▶ $v(\{1, \dots, s_i\}) - v(\{1, \dots, s_{i-1}\}) \geq v(\{s_1, \dots, s_i\}) - v(\{s_1, \dots, s_{i-1}\})$
 - $\{s_1, s_2, \dots, s_l\} \subseteq \{1, \dots, s_l\}$ for $l \leq k$

CLASSES OF GAMES

Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

Core of convex games

For a convex cooperative game (N, v) , it holds $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only $m_v^{id} \in \mathcal{C}(v)$

1. $m_v^{id}(N) = \sum_{i \in N} v(\{1, \dots, i\}) - v(\{1, \dots, i-1\}) = V(N)$
2. $m_v^{id}(S) \geq v(S)$
 - ▶ $S = \{s_1, s_2, \dots, s_k\}$
 - $s_1 < s_2 < \dots < s_k$
 - ▶ $v(\{1, \dots, s_i\}) - v(\{1, \dots, s_{i-1}\}) \geq v(\{s_1, \dots, s_i\}) - v(\{s_1, \dots, s_{i-1}\})$
 - $\{s_1, s_2, \dots, s_l\} \subseteq \{1, \dots, s_l\}$ for $l \leq k$
 - ▶ $m_v^{id}(S) = \sum_{s_i \in S} v(\{1, \dots, s_i\}) - v(\{1, \dots, s_{i-1}\})$

CLASSES OF GAMES

Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

Core of convex games

For a convex cooperative game (N, v) , it holds $\mathcal{C}(v) = \mathcal{W}(v)$

Proof: We show only $m_v^{id} \in \mathcal{C}(v)$

1. $m_v^{id}(N) = \sum_{i \in N} v(\{1, \dots, i\}) - v(\{1, \dots, i-1\}) = v(N)$
2. $m_v^{id}(S) \geq v(S)$
 - ▶ $S = \{s_1, s_2, \dots, s_k\}$
 - $s_1 < s_2 < \dots < s_k$
 - ▶ $v(\{1, \dots, s_i\}) - v(\{1, \dots, s_{i-1}\}) \geq v(\{s_1, \dots, s_i\}) - v(\{s_1, \dots, s_{i-1}\})$
 - $\{s_1, s_2, \dots, s_l\} \subseteq \{1, \dots, s_l\}$ for $l \leq k$
 - ▶ $m_v^{id}(S) = \sum_{s_i \in S} v(\{1, \dots, s_i\}) - v(\{1, \dots, s_{i-1}\})$
 - ▶ $m_v^{id} \geq \sum_{s_i \in S} v(\{s_1, \dots, s_i\}) - v(\{s_1, \dots, s_{i-1}\}) = v(S)$

Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

Consequence:

The Shapley value and convex games

Convex games

$$(S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$$

Consequence:

The Shapley value and convex games

For a convex cooperative game (N, v) , it holds:

1. $\phi(v) \in \mathcal{C}(v)$
2. $\phi(v)$ is the centre of gravity of $\mathcal{C}(v)$

DISADVANTAGES OF THE MODEL AND MOTIVATION FOR NEXT LECTURES

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

1. we might not be interested in forming N

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

1. **we might not be interested in forming N**
 - ▶ we want to analyze coalition formation

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

1. **we might not be interested in forming N**
 - ▶ we want to analyze coalition formation
2. **not every coalition makes sense**

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

1. **we might not be interested in forming N**

- ▶ we want to analyze coalition formation

2. **not every coalition makes sense**

- ▶ two players from completely different fields (why would they want to cooperate)

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

1. **we might not be interested in forming N**
 - ▶ we want to analyze coalition formation
2. **not every coalition makes sense**
 - ▶ two players from completely different fields (why would they want to cooperate)
3. **2^n real values representing v**

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

1. **we might not be interested in forming N**

- ▶ we want to analyze coalition formation

2. **not every coalition makes sense**

- ▶ two players from completely different fields (why would they want to cooperate)

3. **2^n real values representing v**

- ▶ expensive to get all information
- ▶ expensive to store all information

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

1. **we might not be interested in forming N**
2. **not every coalition makes sense**
3. **2^n real values representing v**

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

1. **we might not be interested in forming N**
 - games with coalition structure
2. **not every coalition makes sense**
3. **2^n real values representing v**

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

1. **we might not be interested in forming N**
 - ▶ games with coalition structure
2. **not every coalition makes sense**
 - ▶ games on graphs (Martin Černý)
3. **2^n real values representing v**

Cooperative game

A **cooperative game** is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

1. **we might not be interested in forming N**
 - ▶ games with coalition structure
2. **not every coalition makes sense**
 - ▶ games on graphs (Martin Černý)
3. **2^n real values representing v**
 - ▶ incomplete games (David Sychrovský + Filip Úradník)
 - ▶ interval games (Martin Kunst)
 - ▶ stochastic games (David Ryzák)
 - ▶ models with compact characteristic function (Martin Černý)