

Richard Mužík

richard@imuzik.cz

Introduction to Cooperative Game Theory *as part of series* Cooperative game theory

Introduction

In this lecture you will find brief introduction to cooperative games.

Definition 1 (Cooperative game) A cooperative game is an ordered pair (N, v) , where N is a set of players and $v: 2^N \rightarrow \mathbb{R}$ is the characteristic function. Further, $v(\emptyset) = 0$.

Definition 2 (Payoff vector) Payoff vector is $x \in \mathbb{R}^n$, where x_i represents payoff of player i . It is efficient, if $\sum_{i \in N} x_i = v(N)$. It is individually rational, if $x_i \geq v(i)$.

Definition 3 (Core) For a cooperative game (N, v) , the core $\mathcal{C}(v)$ is

$$\mathcal{C}(v) = \{x \in \mathcal{I}^*(v) \mid x(S) \geq v(S), \forall S \subseteq N\}.$$

Observation 4 (Emptiness of the core) There are cooperative games (N, v) with empty core.

Theorem 5 (Weak Bondareva-Shapley) Cooperative game (N, v) has non-empty core if and only if

$$v(N) \geq \sum_{S \subseteq N} y_S v(S) \text{ for all feasible } y \in \mathbb{R}^{(2^n-1)}$$

Definition 6 For a cooperative game (N, v) , the Shapley value $\varphi(v)$ of player i is

$$\varphi_i(v) = \sum_{S \subseteq N \setminus i} \frac{s!(n-s-1)!}{n!} (v(S \cup i) - v(S))$$

Definition 7 (Marginal vector) For cooperative game (N, v) and permutation $\sigma \in \Sigma_n$ is m_v^σ marginal vector, where $(m_v^\sigma)_i = v(S_{\sigma(i)} \cup i) - v(S_{\sigma(i)})$ and $S_{\sigma(i)} = \{j \in N \mid \sigma(j) < \sigma(i)\}$

Definition 8 (Weber set) For cooperative game (N, v) the Weber set is

$$\mathcal{W}(v) = \text{conv} \{m_v^\sigma \mid \sigma \in \Sigma_n\}$$

Lemma 9 (Shapley value and Weber set) For a cooperative game (N, v) it holds:

$$\varphi(v) \in \mathcal{W}(v)$$

Moreover, $\varphi(v)$ is the center of gravity of $\mathcal{W}(v)$.

Theorem 10 (Weber set and core) For every cooperative game (N, v) , it holds $\mathcal{C}(v) \subseteq \mathcal{W}(v)$

Definition 11 (Classes) The cooperative game (N, v) is said to be

- *monotonic game* $\iff (S \subseteq T \subseteq N) (v(S) \leq v(T))$
- *superadditive game* $\iff (S, T \subseteq N, S \cap T = \emptyset) (v(S) + v(T) \leq v(S \cup T))$
- *convex game* $\iff (S, T \subseteq N) (v(S) + v(T) \leq v(S \cap T) + v(S \cup T))$
- *essential game* $\iff v(N) \geq \sum_{i \in N} v(i)$
- *balanced game* $\iff \mathcal{C}(v) \neq \emptyset$

Theorem 12 (Balanced and essential) *Balanced cooperative games are essential.*

Theorem 13 (Convex and superadditive) *Convex cooperative games are superadditive.*

Theorem 14 (Core of convex games) *For a convex cooperative game (N, v) , it holds $\mathcal{C}(v) = \mathcal{W}(v)$*

Corollary 15 (Shapley value and convex games) *For a convex cooperative game (N, v) , it holds:*

1. $\varphi(v) \in \mathcal{C}(v)$
2. $\varphi(v)$ is the centre of gravity of $\mathcal{C}(v)$

Bibliography

- [1] Hans Peters *Game theory: A multi-leveled approach*. Springer Texts in Business and Economics, 2nd edition, 2015.