Maxwell-Boltzmann distribution.

$$|n(\Xi)_{cl} = \sum_{j} \exp(-\beta(E_{j} - M))$$

$$\langle n_{j} \rangle_{cl} = \exp(-\beta(E_{j} - M))$$

$$|n(\Xi)_{cl} = \sum_{j} \exp(-\beta E_{j})$$

$$|n(\Xi)_{cl} = \sum_{j} \exp(-\beta E_{j})$$

$$|n(\Xi)_{cl} = \sum_{j} \exp(-\beta E_{j})$$

$$\langle n_{j} \rangle_{cl} = \sum_{j} \exp(-\beta E_{j})$$

$$|n(\Xi)_{cl} = \sum_{j}$$

in the thermodynamical limit

$$\frac{\langle n_i \rangle_{cl}}{\langle n_i \rangle_{cl}} \longrightarrow p_0(r) dr = \frac{V}{(2\pi)^3} \exp\left(-\frac{\beta h^2 \kappa^2}{2m}\right) d^3 \vec{k}$$

$$\frac{V}{(2\pi)^3} \int exp\left(-\frac{\beta h^2 \kappa^2}{2m}\right) d^3 \vec{k}$$

Helmholtz

Taking
$$\phi_{cl}$$
, we get

$$\begin{aligned}
& = -\left(\frac{\partial \phi}{\partial M}\right)_{T,N} \\
& + \left(\frac{\partial \phi}{\partial M}\right)_{T,N$$