



Universidad Michocana de
San Nicolás de Hidalgo

Posgrado Conjunto en Ciencias Matemáticas
UNAM-UMSNH

Stochastic Topology in the Curve Complex of a Surface

A thesis submitted in partial fulfillment for the
degree of **Master in Sciences**

by:

Ricardo Esteban Chávez Cáliz

ricardo.ch.cz@gmail.com

1150505J

Supervisor:

Dr. Noé Bárcenas Torres

barcenatmatmor.unam.mx

MORELIA, MICH., October 2020





Universidad Nacional Autónoma de México

Posgrado Conjunto en Ciencias Matemáticas
UNAM-UMSNH

Stochastic Topology in the Curve Complex of a Surface

A thesis submitted in partial fulfillment for the
degree of **Master in Sciences**

by:

Ricardo Esteban Chávez Cáliz

ricardo.ch.cz@gmail.com

1150505J

Supervisor:

Dr. Noé Bárcenas Torres

barcenatmatmor.unam.mx

MORELIA, MICH., October 2020

Fahre fort, übe nicht allein die Kunst, sondern dringe auch in ihr Inneres; sie verdient es, denn nur die Kunst und die Wissenschaft erhöhen den Menschen bis zur Gottheit.

Do not merely practice your art, but penetrate into its interior; it deserves that, because only art and science exalt man to divinity.

Ludwig van Beethoven

Acknowledgments

This work is the result of multiple efforts that pushed collaboration between different areas, people, and institutions. I am fortunate enough to be part of this community, to which it is possible to thank only some parts of it.

First, this thesis would not have been possible without the support and patience of my principal supervisor, Dr. Noé Bárcenas Torres, from whom I have received some of the best tips for Math and Life.

I thank Dr. Jesus Hernandez Hernandez who introduced me to the world of rigid expansions and whose enthusiasm for collaboration planted the seed for this work. His follow-up and expertise allowed us to concrete the results.

I want to thank Dr. Octavio Arizmendi Echegaray for his valuable help in all the probability calculations and insights for random graphs.

I want to express my appreciation to the professors in CIMAT from whom I learned Statistics, Probability, and Scientific Computing.

Thanks to my committee members for all their helpful observations that refined the final work.

Financial support was provided by the project "Aplicación de ensamble de Baum-Connes: topología algebraica y geometría no conmutativa" with key IA100315. Also, I thank CONACYT foundational Research Grant 250747 and PAPIIT Grant IA 100119.

Last but not least, I want to thank all my family and friends for their support in this process, without them I am nothing.

Contents

Acknowledgments

iv

1	The curve graph of a surface	3
1.1	Mapping class group of a surface	4
1.1.1	Nielsen–Thurston classification	5
1.2	Curve graph	6
1.2.1	Simple closed curves	6
1.2.2	The curve graph	7
1.3	Properties of the curve graph	7
1.3.1	Cardinality of the number of vertices	7
1.3.2	Connectivity	8
1.3.3	Locally infinite	8
1.3.4	Clique number	9
1.3.5	Infinite Diameter	9
1.4	Rigidity in graphs	10
2	Rigidity in random graphs	13
2.1	Models for random graphs	14
2.1.1	The Radó graph	14
2.1.2	Erdős–Rényi model	14
2.2	Rigid expansions	15
2.3	Radó graph as a model for the curve graph	19
2.4	Erdős–Rényi as a model for the curve graph	21
2.4.1	Connectivity	21
2.4.2	Locally infinite	21
2.4.3	Clique number	23
2.4.4	Diameter	24
2.5	Conclusions	25
3	Computational experimentation	27
3.1	Simulating Erdős–Rényi random graphs	27
3.2	Rigid expansions algorithms	28
3.3	Time execution comparison	31
3.4	Conclusions	32

A The Generate algorithm	35
---------------------------------	-----------

Bibliography	39
---------------------	-----------

Resumen

Proponemos el estudio de expansiones rígidas en gráficas desde un contexto estocástico, usando la gráfica de Radó y el modelo de Erdős-Rényi para gráficas aleatorias. Revisamos la motivación para estudiar este fenómeno como un concepto concebido en la gráfica de curvas de una superficie y la viabilidad de hacerlo usando estos modelos. Proveemos de simulaciones computacionales con las optimizaciones correspondientes

Palabras clave: Probabilidad, topología, superficies, gráficas aleatorias, expansiones rígidas.

Abstract

We propose the study of rigid expansions in graphs in a stochastic context using the Radó graph and the Erdős-Rényi model. We review the motivation to study this phenomenon as a concept originally conceived in the curve graph of a surface and then the feasibility to do it through probabilistic models. We also provide computational simulations with the corresponding optimizations.

Key words: Probability, topology, surfaces, random graphs, rigid expansions.

Introduction

The curve graph $\Gamma(S)$ associated to a surface S appears naturally in the study of $Mod(S)$, the mapping class group of S , which is a central subject in contemporary mathematical research. We are interested in a rigidity concept of this graph; in general, the idea behind rigidity phenomena is to describe morphisms among objects using their structure.

The folkloric version of rigidity in the $Mod(S)$ context is that if we consider X and Y , under suitable conditions, then every homomorphism $Mod(X) \rightarrow Mod(Y)$ will be induced by manipulation of the underlying surfaces.

Ivanov sketched in [20, Ivanov 97] the proof that every automorphism of $C(S)$, the flag complex of $\Gamma(S)$, is induced by a self-homeomorphism of S . Due to its simplicity and resemblance to the other rigidity results, this argument is the favorite in the literature.

A research line, lead by Aramayona and Leininger, introduced the idea of *rigid sets* as subsets of vertices *that allows to extend a local notion of rigidity to a global one*. To find large rigid sets, in [2, Aramayona, Leininger 16] there's a proof of the existence of an increasing sequence of finite rigid sets, that exhaust the curve graph. For this, they introduced a method called **rigid expansions**.

Rigidity in graphs is, regardless of its interpretation in the curve graph, an interesting phenomenon by itself. Due to the discrete nature of rigid expansions, it is reasonable to seek for a probabilistic approach; our goal is to address this particular path.

We want to answer the rather vague question: *How **common** is rigidity in graphs*, specifically by answering *how rigid expansions **usually** behave*. Also, we review the feasibility of *studying the curve complex of a surface from a probabilistic point of view*.

Probabilistic models give formal meaning to words like "common" or "usually". We study the rigidity phenomenon in this context and analyze the conditions under which these models fit the known properties of the curve graph.

In Chapter 1, we motivate the study of the curve graph and review its most essential properties. Then, we introduce rigidity within the context of Graph theory.

In the Chapter 2, we propose the study of rigidity from the stochastic point of view through the Radó graph and the Erdős-Rényi model. In the aim to study the curve graph of a surface with a simple model, we justify that the genus of the surface cannot be finite. Thus, we end up with an asymptotic probabilistic analog to Bering and Gaster's result. This result asserts that the Radó graph embeds into the curve graph $C(S)$ of a surface S if and only if S has infinite genus.

Finally, we present a computational implementation of the algorithm to handle rigid expansions. With the corresponding optimizations that the method requires, we can take a closer look at the rigidity phenomena.

Chapter 1

The curve graph of a surface

The study of surfaces in a strictly topological viewpoint has led us to forget significant information about them. A way to revert this is to attach a group to it, the **mapping class group** of the surface. It is denoted by $Mod(S)$ and encodes the *symmetries* of the surface. This group is defined as the set of isotopy classes of orientation-preserving homeomorphisms of S . In the first section of this chapter, we give the formal definition of this group and establish the very important role of this concept in Mathematics.

The **curve complex** of the surface, denoted by $C(S)$ appears naturally in the study of $Mod(S)$. It is a simplicial complex that encodes intersection patterns of simple closed curves in S . We focus part of the discussion on the relationship between the algebraic structure of $Mod(S)$ and the combinatorial topology of S .

Many of the progress in understanding $Mod(S)$ has been possible by a well-known analogy among two very important classes of groups: arithmetic groups and mapping class groups. In this parallelism panorama, the desire of an equivalent result to the Margulis Superrigidity for mapping class groups arises.

In the last section of this chapter, we settle the bases to understand rigidity within a Graph theory context. An approach called *rigid expansions*, see [2, Aramayona, Leininger 16] and [19, Hernandez 19], allows us to build up subgraphs preserving the rigidity property and is compatible with stochastic tools.

Many results and definitions in this chapter were extracted from [16, Farb]. They are quite popular and equivalents can easily be found in the literature, however, they are written here to establish nomenclature. Familiarity with basic concepts is assumed.

1.1 Mapping class group of a surface

We have the following fundamental, well-known result about surfaces.

Theorem 1.1.1 (Classification of surfaces). *Any closed, connected, orientable surface is homeomorphic to the connected sum of a 2-dimensional sphere with $g \geq 0$ tori. Any compact, connected, orientable surface is obtained from a closed surface by removing $b \geq 0$ open disks with disjoint closures. Even more, the set of homeomorphism types of compact surfaces is in bijective correspondence with the set $\{(g, b) : g, b \geq 0\}$.*

We are so familiar with this result that we usually forget what it is saying. It seems like, in the eyes of a topologist, surfaces are just boring, but this is because we are forgetting all the geometric information about them. $Mod(S)$ helps to recover this data, the magic happens when this group acts on the **Teichmüller space** of S , which is the space of hyperbolic metrics on S up to isotopy. A central result is that this action is properly discontinuous and the quotient space $M(S) = Teich(S)/Mod(S)$ is the **moduli space of Riemannian surfaces homeomorphic to S** . The space $M(S)$ is an essential object in mathematics and the group $Mod(S)$ encodes most of the topological features of $M(S)$.

$Mod(S)$, $Teich(S)$, and $M(S)$ can be found in several of different contexts in mathematics: hyperbolic geometry, algebraic geometry, combinatorial group theory, geometric group theory, symplectic geometry, 3-manifold theory, dynamics, and so on. The algebraic structure of $Mod(S)$, the geometry of $Teich(S)$, and the topology of $M(S)$ are just the strands that are used to weave the rich tapestry of the nature of the surface.

Before we continue, let us establish some nomenclature. The g in 1.1.1 is called the *genus* of the surface and the b is the number of *boundary components*. One way to obtain a non-compact surface from a compact one is to remove m points from the interior of it; in this case, we say that the resulting surface has m punctures. From now on, unless otherwise specified, we will be thinking of compact, connected, oriented surfaces that are possibly punctured (in this case they cease to be compact). Therefore, we can specify the surfaces by the triplet (g, b, m) . We will denote by $S_{g,m}$ a surface of genus g with m punctures and empty boundary; such a surface is homeomorphic to the interior of a compact surface with m boundary components. Also, for a closed surface of genus g , we will abbreviate $S_{g,0}$ as S_g and ∂S denotes the (possibly disconnected) boundary of S .

There are numerous definitions for the mapping class group of a surface. We will be working with the following:

Definition 1.1.1.1. Let S be a surface, the **mapping class group** of S , denoted by $Mod(S)$, is the following quotient:

$$Mod(S) = Homeo^+(S)/Homeo_0(S),$$

where $Homeo^+(S)$ is the group of orientation-preserving homeomorphisms of S , that are the identity on the boundary. $Homeo_0(S)$ is the subgroup formed by homeomorphisms of S that are isotopic to the identity.

We could consider diffeomorphisms instead of homeomorphisms, or homotopy classes instead of isotopy classes; this results in isomorphic groups, see [16, Farb, p. 41] for details in why we can do this. Summarizing, we can find the following variations in the definition of $Mod(S)$:

$$\begin{aligned} Mod(S) &= \pi_0(Homeo^+(S, \partial S)) \\ &\approx Homeo^+(S, \partial S)/homotopy \\ &\approx \pi_0(Diff^+(S, \partial S)), \end{aligned}$$

where $Diff^+(S, \partial S)$ is the group of orientation-preserving diffeomorphisms of S that are the identity on the boundary. It can be taken to be either smooth homotopy relative to the boundary or smooth isotopy relative to the boundary.

Thanks to Thurston's classification theorem there is a characterization of the homeomorphisms of a compact orientable surface. This classification is useful to describe the curve graph which will be analyzed in the next section.

1.1.1 Nielsen–Thurston classification

Given a homeomorphism $f : S \rightarrow S$, there is a map g isotopic to f such that at least one of the following statements holds:

- g is periodic, i.e. some power of g is the identity;
- g preserves some finite union of disjoint simple closed curves on S (in this case, g is called reducible); or
- g is pseudo-Anosov.

The definition of a **pseudo-Anosov map** relies on the notion of a measured foliation, a geometric structure on S . It consists of a singular foliation and a measure in the transverse direction (i.e. that is constant in transverse arcs). For the full definition of pseudo-Anosov elements and the proof of this theorem, we can refer to [16, Farb, ch. 13].

The study of mapping class groups is a wide and challenging area. It is outside of the interests of this thesis to review the details and repercussions of this vast field. Yet, there are several known properties of $Mod(S)$ that it would be nice to have in mind for further work:

- Finitely generated and presented.
- It has a subgroup of finite index which doesn't have torsion.
- $Mod(S_{g,m}) \cong Out(\pi_1(S_{g,m}))$, for the closed case.
- $H_1(Mod(S_{g,m}), \mathbb{Z}) = 1$ when $(g \geq 3, m = 0)$.

1.2 Curve graph

1.2.1 Simple closed curves

Definition 1.2.0.1. A *closed curve* in a surface S is a continuous map $\mathbb{S}^1 \rightarrow S$ and it's called **simple** if the map is an embedding. We will usually identify a closed curve with its image in S . A closed curve is called **essential** if it is not homotopic to a point, a puncture, or a boundary component.

Among the adjectives that a curve can acquire we have the following:

- α is **separating**, if $S - \alpha$ has two components, otherwise it is called **non-separating**.
- It is called **essential** if no component of $S - \alpha$ is a disk.
- It is **non-peripheral** if no component of $S - \alpha$ is an annulus.

We are interested in **essential** and **non-peripheral** curves, and all curves will be assumed in this sense unless otherwise specified.

The idea behind the construction of the curve graph is to stratify the set of homotopy classes of curves on a surface. For this to make sense we define the **geometric intersection number** between free homotopy classes a and b of simple closed curves in a surface S . This is defined to be the minimal number of intersection points between a representative curve in the class a and a representative curve in the class b :

$$i(a, b) = \min\{|\alpha \cap \beta| : \alpha \in a, \beta \in b\}.$$

It is convenient to adopt a slight abuse of notation by writing $i(\alpha, \beta)$ for the intersection number between the homotopy classes of simple closed curves α and β . It is useful to think that this number can be computed by finding representatives α and β that achieve the minimal intersection in their homotopy classes so that $i(a, b) = |\alpha \cap \beta|$ (when this is the case, we say that α and β are in minimal position). Although the geometric intersection number is a useful and intuitive invariant it is not always easy to compute, whenever this is the case we can appeal to the algebraic intersection number. For further discussion on this see [16, Farb].

1.2.2 The curve graph

Definition 1.2.0.2. *The **curve graph** $\Gamma(S)$ of a surface S is constructed with the following data:*

- **Vertices.** *There is a vertex in $\Gamma(S)$ for every isotopy class of essential, non-peripheral, simple, closed curves in S .*
- **Edges.** *There is an edge between the corresponding vertices of isotopy classes a and b whenever $i(a, b) = 0$.*

Definition 1.2.0.3. *The **curve complex of the surface** $C(S)$ is defined to be the flag complex of the curve graph just defined.*

1.3 Properties of the curve graph

The goal of this section is to state known properties of the curve graph, we use this to establish the appropriate parameters in a probabilistic model. Notice that the construction of the curve complex is completely determined by the curve graph, hence the probabilistic models can work in the same sense. Let's keep in mind the following exceptional cases; they are responsible for the conditions stated in the hypothesis of the following theorems for g and n . For $S^2, S_{0,1}, S_{0,2}, S_{0,3}$ the curve graph is empty and for $T^2, S_{1,1}$ and $S_{0,4}$ is a countable disjoint union of points.

1.3.1 Cardinality of the number of vertices

Theorem 1.3.1. *If $g \geq 1$ or $n \geq 4$ then the set of vertices in $\Gamma(S_{g,n})$ is countably infinite.*

It is well known that for T^2 there is an explicit identification for the isotopy classes of essential curves with the rational numbers. In this case, there aren't disjoint curves, the following figure can help to convince us of this fact.




FIGURE 1.1: T^2 with representatives of typical elements of curves.

This identification can be thought of as the induction basis. The induction step over g comes from splitting the surface, by induction hypotheses none of the resulting surfaces can have a infinitely many classes of curves. Since the surfaces are second countable, the number of classes of curves cannot be uncountable.

1.3.2 Connectivity

Theorem 1.3.2. *If $3g + n \geq 5$, then $\Gamma(S_{g,n})$ is connected.*

To proof this theorem we can show that, for any two isotopy classes a and b of simple closed curves in $S_{g,n}$ there exists a sequence of isotopy classes

$$a = c_1, \dots, c_k = b,$$

where $i(c_i, c_{i+1}) = 0$. This can be done proceeding by induction over $i(a, b)$. The full proof of this theorem can be found in [16, Farb, p. 93].

1.3.3 Locally infinite

Theorem 1.3.3. *If $3g + n \geq 5$, then $\Gamma(S_{g,n})$ is locally infinite.*

The idea behind the proof is that given any $\alpha \in \Gamma(S)$ we can construct a family of isotopy classes of curves that are disjoint to α . The following picture gives us an intuitive idea on how to do this whenever we have large enough genus.




FIGURE 1.2: S_3 with typical representative curves which exemplify the idea behind locally infinity property.

For the complete argument, let α be any simple closed curve on S , the surface $S - \alpha$, obtained by cutting S open along α , contains at least one connected component of Euler characteristic at most -2 (guaranteed by the $3g + n \geq 5$ condition). Such component contains infinitely many distinct homotopy classes of simple closed curves disjoint from α .

1.3.4 Clique number

A **clique** in a graph G is a complete subgraph of G . The clique number $cl(G)$ of a graph G is the maximum order of a clique of G .

Theorem 1.3.4. *If $3g + n \geq 5$, then the clique number of $\Gamma(S_{g,n})$ is $3g - 3 + n$.*

$3g - 3 + n$ is the number of curves in a pants decomposition of S , i.e. a maximal collection of disjoint, not freely homotopic, essential, simple closed curves, which decompose S into $2g - 2 + n$ open subsurfaces homeomorphic to a thrice punctured sphere. For a full proof of this well-known fact refer to [18, Hatcher, Thurston 80].




FIGURE 1.3: Exemplification of a pants decomposition of a surface.

1.3.5 Infinite Diameter

Theorem 1.3.5. *If $3g + n \geq 5$ then $\text{diam}(\Gamma(S)) = \infty$.*

The proof for this theorem relies on the fact that for any pseudo-Anosov element $h \in \text{Mod}(S)$, any $\gamma \in V(\Gamma(S))$ and any $k \in \mathbb{Z}$

$$d_C(h^k(\gamma), \gamma) \geq c|k|,$$

This provides the infinite diameter property. The proof of this theorem is highly non-trivial, refer to [25, Masur, Minsky 99] for a complete review.

The curve graph and the curve complex are fundamental tools in the study of the surfaces. There several known properties of them that it would be nice to have in mind to improve probabilistic models for further work.

1. $C(S)$ is hyperbolic.
2. In the infinite genus case $\text{diam}(\Gamma(S)) = 2$.
3. There's an isomorphism between $\text{Mod}^*(S)$ and $\text{Aut}(C(S))$, except when $(g, m) \in \{(1, 2), (1, 1), (2, 0), (0, 4)\}$.

1.4 Rigidity in graphs

This section intends to track down the motivation of rigid expansions and give the required definitions.

As mentioned in the introduction of the chapter, rigidity appears in the mapping class group context in light of its comparison with arithmetic groups. In [4, Aramayona, Souto 16] we can find a survey on the search of an analog for the Margulis Superrigidity theorem. In this article, they provide three different perspectives: a Lie theoretical, a geometric, and a folkloric one.

The Lie theoretic version states that every homomorphism $\text{Mod}(X) \rightarrow \text{Mod}(Y)$ is induced by a homomorphism between the associated groups of diffeomorphisms with compact support disjoint from the boundary $\text{Diff}_c(X) \rightarrow \text{Diff}_c(Y)$.

A direct formulation of geometric superrigidity cannot hold when the moduli space is endowed with any reasonable metric. However, there are ways to turn this around, saying that every (irreducible) homomorphism between mapping class groups induces a holomorphic map between the corresponding moduli spaces.

The folkloric version of Mostow and Margulis superrigidity claims that the only homomorphisms between lattices are the “*obvious ones*”, in the $\text{Mod}(S)$ context this will

mean that if we consider X and Y , under suitable conditions, then every homomorphism $Mod(X) \rightarrow Mod(Y)$ will be induced by manipulation of the underlying surfaces.

A result due to Ivanov [20, Ivanov 97], Korkmaz [23, Korkmaz 99] and Luo [24, Luo 00], asserts that, excluding few well-understood cases, the curve complexes are simplicially rigid. This means that the group $Aut(C(S))$ of simplicial automorphisms of $C(S)$ is isomorphic to the extended mapping class group. This result is sometimes interpreted as a proof that **the automorphisms of the curve complex are all geometric**.

In the aim to generalize this result to broader types of simplicial self-maps, in [2, Aramayona, Leininger 12] was introduced the concept of **rigid sets** in the curve complex. Let S be a surface different from $S_{1,2}$, $Y \subset C(S)$ is called **rigid** if for every locally injective simplicial map $\Phi : Y \rightarrow C(S)$ there exists $h \in Mod^*(S)$ with $h|_Y = \Phi$, unique up to the pointwise stabilizer of Y in $Mod^*(S)$.

Later in [3, Aramayona, Leininger 16] a method for enlarging rigid subgraphs is presented and the proof that for almost all surfaces of finite topological type, there exists an **increasing sequence of finite rigid sets** that exhaust the curve graph of which has trivial pointwise stabilizer in $Mod^*(S_{g,n})$.

In [19, Hernández 19] there is a proof of a similar result to Aramayona and Leininger's. The method, called **rigid expansions**, allowed to obtain new results concerning edge-preserving maps.

Losing the big picture that rigidity represents, we can land this vast journey in the following graph-theoretic definitions.

Definition 1.4.0.1. Let Γ be a simplicial graph and $H < \Gamma$ a vertex-induced subgraph. A function $f : H \rightarrow \Gamma$ is **locally injective** if $f|_{star(v)}$ is injective for all $v \in V(H)$.

Note 1.4.1. Remember, $star(v)$ is the vertex-induced subgraph with vertices $\{v\} \cup N(v)$ (v plus its neighborhood).

Definition 1.4.1.1. $H < \Gamma$ is **rigid** if every locally injective function defined in H can be extended to an automorphism of Γ .

A vertex $v \in V$ in a graph is uniquely determined by $A \subset V(G)$, denoted $v = \langle A \rangle$, if v is the unique common neighbor of every element of A , i.e.

$$\{v\} = \bigcap_{w \in A} \text{link}(w).$$

Definition 1.4.1.2. *The first rigid expansion of $Y \subset \Gamma$, denoted by Y^1 , is the vertex-induced subgraph whose vertices are*

$$V(Y) \cup \{v \in V(\Gamma) : \exists A \subset V(Y) \text{ where } v = \langle A \rangle\}.$$

We also define $Y^0 = Y$ and, inductively, $Y^k = (Y^{k-1})^1$.

Recalling that in Proposition 3.5 in [2, Aramayona, Leininger 16], Aramayona and Leininger prove that if $Y \subset C(S)$ is a rigid set, then so is Y^r for all $r \geq 0$. So this method in fact preserves the desired property.

It would be nice to have conditions which determine whether a subgraph is rigid or not. So far we don't know non-trivial, necessary conditions to check rigidity, i.e. other than connectivity there's not much else.

With these definitions, we can proceed to settle a probabilistic model so that we can analyze the rigidity concept in graphs from a stochastic point of view. With the reviewed properties of the curve graph, we can determine the feasibility of studying the curve graph through simple models.

Chapter 2

Rigidity in random graphs

The use of the probabilistic method in discrete mathematics has become a prominent idea in the area in recent times. It provides existence proofs where objects have certain desirable properties and the construction of explicit examples is challenging. This has been just the beginning of the use of probabilistic tools within a deterministic context.

Complex topological spaces arise quite naturally in a lot of scientific contexts. Probability theory implements different approaches to model those spaces; even in complex configurations, it can be possible by doing approximations, to study topological invariants. In this sense, stochastic topology can be thought of as a tool for topology in the same sense as statistical mechanics is used to studying a macroscopic physical system when classical mechanics finds these systems too complicated to solve.

Stochastic topology finds its early motivation in applied problems. Nevertheless, in recent articles, it has been used to provide deeper insight into theoretical questions. For example, with probabilistic analogs of very classical topology conjectures, like Whitehead's Asphericity Conjecture [13, Costa, Faber 15].

Probability theory can help us understand the ubiquity of certain mathematical phenomena. For example, *many* simplicial complexes and posets which arise from combinatorial constructions are homotopy equivalent to a wedge of spheres, or that hyperbolicity is *common* in random groups. With Probability theory, we can give formal meaning to these expressions.

In this chapter, we review rigid expansions in simple probabilistic models. Afterward, we analyze the feasibility of modeling the curve graph using these proposals.

Familiarity with basic concepts in probability theory such as probability spaces, random variables, and basic theorems are assumed.

2.1 Models for random graphs

2.1.1 The Radó graph

Let $0 < p < 1$ be fixed, $\mathcal{G}(\mathbb{N}, p)$ is the probability space which consists of all graphs with vertex set \mathbb{N} , whose edges are chosen independently with probability p . In other words, a random graph $G \in \mathcal{G}(\mathbb{N}, p)$ is a collection $(X_{ij}) = \{X_{ij} : 1 \leq i < j\}$ of independent *Bernoulli*(p) r.v., where a pair ij is an edge of G if and only if $X_{ij} = 1$.

Erdős and Rényi proved in [15, Erdős, Rényi 63], that every countably infinite random graph is isomorphic to the **Radó graph**. A construction of this graph can be done using binary numbers; identify the vertices of the graph with the natural numbers and then every edge appears between vertices x and y in the graph (assuming $x < y$) whenever the x -th bit of the binary representation of y is nonzero. This means, for example, that all odd-numbered vertices will be neighbors of vertex 0, and that the larger neighbors of vertex 1 are all vertices with numbers congruent to 2 or 3 mod 4.




FIGURE 2.1: Binary construction of the Radó graph.

2.1.2 Erdős-Rényi model

The Erdős-Rényi model for random graphs is the finite version of the Radó graph. In this model, the parameter p is usually taken as a function of n . This provides, unlike the past model, a variety of graphs when n tends to infinity.

Definition 2.1.0.1. Denote by $\mathcal{G}(n, p)$ to the probability space formed by all the graphs of n vertices and probability measure

$$\mathbb{P}(G \in \mathcal{G}(n, p)) = p^k (1 - p)^{\binom{n}{2} - k},$$

where k is the number of edges in G , the σ -algebra is given by the power set.

Note: There is a variation of the model, where we rather choose randomly exactly m edges among the $\binom{n}{2}$ possible.

We can also think this model like $\binom{n}{2}$ i.i.d. $Bernoulli(p)$ that represent the edges. From this, we can immediately get some properties of the degree of a vertex v .

- The probability that a given vertex v has degree k is given by

$$b(k; n-1, p) = \binom{n-1}{k} \cdot p^k \cdot (p-1)^{n-k-1}.$$

- The expected degree is $(n-1) \cdot p$.
- The variance of this degree is $(n-1) \cdot (1-p) \cdot p$.

The degree distribution can be helpful to do optimizations in the rigid expansions algorithms. This is outlined in the next chapter.

2.2 Rigid expansions

We will focus on the rigidity calculations for the finite case. Then, we will analyze what happens when n tends to infinite. For this, we need to compute the probability that the following events occur.

1. Let us call E_m to the event when $v = \langle A_m \rangle$, where v is a vertex and A_m a subset of vertices of size m .
2. A_k generates a rigid expansion.
3. A_k generates a rigid expansion with s new elements.

For the calculations concerning the first event, take a look at the following figure.




FIGURE 2.2: Probability of uniquely determined vertices.

If $\langle A_m \rangle = v$, there is a edge between v and every vertex in A , and none of the remaining $n - m - 1$ vertices is also connected to every vertex in A , i.e.

$$\mathbb{P}(E_m) := \mathbb{P}(\langle A_m \rangle = v) = p^m(1 - p^m)^{n-m-1}.$$

Using the `networkx` library in `python` we reproduce the following experiment:

Uniquely determined vertex experiment. Let n, p, m be fixed.

1. Generate an Erdős-Rényi graph $G \in \mathcal{G}(n, p)$ with labeled vertices.
2. Excluding the n -th vertex, take a random subset of vertices of size m .
3. Verify if this subset uniquely determine the n -th vertex.

In the next chapter, we explain how to generate random graphs for the first step of the experiment. To simplify the process we took, without losing generality, the last vertex as a particular element of the experiment.

Fixing different values for n and p is possible to compute the empiric probability for each possible value of m .




FIGURE 2.3: Theoretical and empirical probabilities of uniquely determined a vertex. For different values of n and p varying among all the possible values of m .

For these estimations, we calculated the empiric probability by repeating this experiment 500 times. Then, we counted the number of times when the n -th vertex was uniquely determined by the random set.

Notice that certain values of m are more *effective* than others, in the sense that, depending on the parameters of the model, it is more likely that a subset of a certain size uniquely determines a vertex. This can be used, as described in the next chapter, to optimize the simulations.

In this table appears the supremum of absolute differences between hypothesized and empirical probability $\forall m \in \{1, 2, \dots, n\}$ for the different values of n and p .

$p \backslash n$	8	15	20
0.1	$9.01E - 03$	$1.20E - 02$	$8.09E - 03$
0.1	$9.01E - 03$	$1.20E - 02$	$8.09E - 03$
0.1	$9.01E - 03$	$1.20E - 02$	$8.09E - 03$

TABLE 2.1: Supremum of absolute differences between hypothesized and empirical probabilities.

For the second event, if A_k does not generate a rigid expansion is because none of the subsets of A_k uniquely determined a vertex outside of it. We have:

$$\mathbb{P}(A_k \text{ generates a rigid expansion}) = 1 - \prod_{m=1}^k (\rho_{m,k})^{\binom{k}{m}},$$

where $\rho_{m,k} = \left(1 - \mathbb{P}(E_m)\right)^{n-k}$.

Just as before, we reproduce the following experiment:

Rigid expansion experiment. Let n, p, k be fixed.

1. Generate an Erdős-Rényi graph $G \in \mathcal{G}(n, p)$.
2. Take a random subset of vertices of size k .
3. Verify if this set generates a rigid expansion.

Notice that the third step is a critical point of the experiment; we must verify among **all the possible subsets** of A_k . In the next chapter, we explain the optimizations that needed to be done.

Again, we repeated this experiment 500 times and calculated the empiric probability that a random set generates a rigid expansion.




FIGURE 2.4: Theoretical and empirical probabilities of expanding A_k . For different values of n and p varying among all the possible values of k .

In the following table appears the supremum of absolute differences between hypothesized and empirical probabilities $\forall k \in \{1, 2, \dots, n\}$ for the different values of n and p .

$p \backslash n$	8	15	20
0.1	$9.01E-03$	$1.20E-02$	$8.09E-03$
0.1	$9.01E-03$	$1.20E-02$	$8.09E-03$
0.1	$9.01E-03$	$1.20E-02$	$8.09E-03$

TABLE 2.2: Supremum of absolute differences between hypothesized and empirical probabilities.

The calculations for the last question are helpful if we want to approximate the sequence of rigid expansions of A_k by a Markov chain. Consider $\{0, 1, \dots, n\}$ as the states space of the Markov chain with transition matrix given by:

$$a_{k,k+s} = \mathbb{P}(A_k \text{ generates a rigid expansion by } s \text{ elements}).$$

Notice that the deterministic process stops once an iteration fails to add new vertices. In our stochastic approximation, a new $G \in \mathcal{G}(n, p)$ is considered for each step, hence, it is allowed to "have extra tries to expand".

This probability calculations are more difficult to obtain. To start understanding this phenomenon we can simulate with our computational tools the following experiment:

Increase size by a rigid expansions experiment

Let n, p, k be fixed.

1. Generate an Erdős-Rényi graph $G \in \mathcal{G}(n, p)$.
2. Take A_k a random set of k vertices.
3. Produce the first rigid expansion from the graph spanned by A_k .
4. Return the size of the expanded subgraph.

This experiment yields a random variable that depends on n, p , and k . Fixing n and p we obtained a sample of size 50 for every possible value of k . Using the resulting histogram as an empirical density function we obtain the following figure. It graphically describes the nature of the transition matrix of a sequence of rigid expansions.




FIGURE 2.5: Empirical transition probability matrix.

2.3 Radó graph as a model for the curve graph

In Chapter 1 we outlined the properties of the curve graph associated with a surface, thus the proposed models should at least guarantee the following properties:

1. Countably infinite number of vertices.
2. Connectedness.

3. Locally infinite.
4. Clique number $3g - 3 + m$.
5. Infinite diameter.

The Radó graph satisfies that every finite or countably infinite graph is an induced subgraph of it [11, Cameron 97]. The clique number property of $\Gamma(S)$ implies that, if S is a surface of finite genus, is not possible that the Radó graph is embedded into $\Gamma(S)$. Even more, a result by Bering and Gaster [7, Bering, Gaster 17] states that the converse is also valid.

Theorem 2.3.1. *The random graph embeds into the curve graph $C(S)$ of a surface S if and only if S has infinite genus.*

Therefore, if we want to study the curve graph of a **surface of finite genus** using the Radó graph, we have to think of it as a subgraph of it. A simple approach to do it is to take a random subset of vertices of a graph $G \in \mathcal{G}(\mathbb{N}, p)$ and then consider the vertex induced subgraph. It turns out that for a.e. $G \in \mathcal{G}(\mathbb{N}, p)$ the sequence $cl(G_n)$ is almost entirely determined.

To study cliques in the Radó graph, let us denote E_r de expected number of cliques in a subgraph of size n , i.e $E_r := E(n, r) = \binom{n}{r} p^{\binom{r}{2}}$.

Let $0 < \epsilon < \frac{1}{2}$. Given a natural number $r > 2$ let n_r be the maximal natural number for which $E(n_r, r) \leq r^{-(1+\epsilon)}$, and let n'_r be the minimal natural number for which $E(n'_r, r) \geq r^{1+\epsilon}$.

Theorem 2.3.2. *For a.e. $G \in \mathcal{G}(\mathbb{N}, p)$ there is a constant $m_0 = m_0(G)$ such that if $n \geq m_0$ and $n'_r \leq n \leq n_{r+1}$, then $cl(G) = r$.*

The theorem states that if r is fixed and finite, the number of vertices must be finite as well, the proof can be found in [8, Bollobás p. 284]. Therefore, **it is not possible to obtain a curve graph by a uniform selection of vertices.**

Notice that the **clique number property is not generic** at all, unlike the others listed above, the clique number is the only property which actually depends on the genus of the surface. Therefore, is not a surprise that this property is highly restrictive in the plan of setting a generic model.

Prohibited configurations appear often in the literature, for example in [1, Alcazar 15] they want to ensure that a random graph does not have cycles, implying that the clique number is 2. A discrete MCMC algorithm was used to sample uniformly random

trees of size n , with the **Generate algorithm**. It produces a maximal tree of any not directed graph with n vertices uniformly among all the possible ones. In the appendix, we summarize the results of this method, read [9, Broder 89] for the full analysis.

There is a chance of finding an analog of the Generate algorithm to ensure a fixed clique number, but the scope of this work is to study rigid expansions in a simple probabilistic model. In this spirit, it remains to examine the plausibility of the Erdős-Rényi model and do an asymptotic analysis.

2.4 Erdős-Rényi as a model for the curve graph

2.4.1 Connectivity

Theorem 2.4.1. *Let $\omega(n)$ be a function that tends to infinity arbitrarily slow as n tends to infinity.*

- If $p \geq \frac{\log(n) + \omega(n)}{n}$ then

$$\lim_{n \rightarrow \infty} \mathbb{P}(G \in \mathcal{G}(n, p) \text{ is connected}) = 1.$$

- If $p \leq \frac{\log(n) - \omega(n)}{n}$ then

$$\lim_{n \rightarrow \infty} \mathbb{P}(G \in \mathcal{G}(n, p) \text{ is disconnected}) = 1.$$

Here $\omega(n)$ represents the control over the convergence, in other words, the uncertainty. The theorem is just saying that in order to reduce $\omega(n)$ we must increase the size of the graph.

This theorem can be proved by first showing that for a large n almost all graphs consist of a single connected component and k isolated points, the theorem follows from a counting argument. The complete proof can be found in [14, Erdős-Rényi, p. 59].

2.4.2 Locally infinite

To fulfill this property we need to take $p(n)$ in a way that the expected degree grows along with the vertices, i.e $np \rightarrow \infty$. Notice that the conditions for the connectivity threshold are more than enough to ensure this. The full argument is stated in the following theorem:

Theorem 2.4.2. Let $G \in \mathcal{G}(n, p)$ and $D_n \sim b(k; n-1, p)$ the random variable that describes the degree of a vertex in G . Take $p(n) = \frac{\epsilon}{n^a}$ with fixed $\epsilon, a > 0$ then:

- If $a \geq 1$ then $\{D_n\} \xrightarrow{p} c$, where c is a finite constant.
- If $a < 1$ then $\lim_{n \rightarrow \infty} \mathbb{P}(D_n \text{ is finite}) = 0$.

Proof. When $a = 1$ this is the Binomial's Poisson approximation.

For $a > 1$ take $k = 0$ so we have:

$$\lim_{n \rightarrow \infty} \mathbb{P}(D_n = 0) = \lim_{n \rightarrow \infty} \left(1 - \frac{\epsilon}{n^a}\right)^n = \lim_{n \rightarrow \infty} \exp\left(\ln\left(1 - \frac{\epsilon}{n^a}\right)^n\right) = \lim_{n \rightarrow \infty} \exp\left(n \cdot \ln\left(1 - \frac{\epsilon}{n^a}\right)\right).$$

If $f(n) = n \cdot \ln\left(1 - \frac{\epsilon}{n^a}\right)$, then $\lim_{n \rightarrow \infty} f(n) = \lim_{n \rightarrow \infty} \frac{\ln\left(1 - \frac{\epsilon}{n^a}\right)}{\frac{1}{n}}$. Using L'Hôpital's rule for limits we obtain:

$$\lim_{n \rightarrow \infty} f(n) = \lim_{n \rightarrow \infty} \frac{\frac{1}{(1 - \frac{\epsilon}{n^a})} \cdot (-\epsilon a n^{-a-1})}{-1 \cdot n^{-2}} = \lim_{n \rightarrow \infty} \frac{\frac{n^a}{n^a - \epsilon} \cdot (-\epsilon a n^{-a-1}) \cdot n^2}{-1} = \lim_{n \rightarrow \infty} -\frac{\epsilon a n}{n^a - \epsilon} = \lim_{n \rightarrow \infty} -\frac{\epsilon a}{a n^{a-1}}.$$

From here, if $a > 1$ then $\lim_{n \rightarrow \infty} f(n) = 0$, if $a < 1$ then $\lim_{n \rightarrow \infty} f(n) = -\infty$. So we have

$$\lim_{n \rightarrow \infty} \mathbb{P}(D_n = 0) = \begin{cases} \lim_{n \rightarrow \infty} e^{f(n)} = 1, & \text{if } a > 1 \\ \lim_{n \rightarrow \infty} e^{f(n)} = 0, & \text{if } a < 1 \end{cases}.$$

This concludes the first part of the theorem.

For $a < 1$ and $k > 0$:

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbb{P}(D_n = k) &= \lim_{n \rightarrow \infty} \binom{n}{k} \cdot \left(\frac{\epsilon}{n^a}\right)^k \cdot \left(1 - \frac{\epsilon}{n^a}\right)^{n-k} \\ &= \lim_{n \rightarrow \infty} C_k \cdot n^k \cdot \frac{\left(\frac{\epsilon}{n^a}\right)^k}{\left(\frac{n^a - \epsilon}{n^a}\right)^k} \cdot \left(1 - \frac{\epsilon}{n^a}\right)^n \\ &= C_k \lim_{n \rightarrow \infty} n^k \cdot \left(\frac{1}{n^a - \epsilon}\right)^k \cdot \left(1 - \frac{\epsilon}{n^a}\right)^n \\ &= C_k \lim_{n \rightarrow \infty} n^s \cdot \left(1 - \frac{\epsilon}{n^a}\right)^n \end{aligned}$$

where $s = k(1 - a) > 0$, hence $\lim_{n \rightarrow \infty} \mathbb{P}(D_n = k) = 0, \forall k > 0 \quad \square$

So, when $a < 1$ we obtain the locally infinite property, and this condition is always satisfied in the connectivity threshold.

Consider $X_k = X_k(G)$, the random variable that describes the number of vertices of degree k in a graph G . The following theorem gives a complete description of the degree distribution.

Theorem 2.4.3. *Let $\epsilon > 0$ be fixed, $\epsilon n^{-3/2} \leq p = p(n) \leq 1 - \epsilon n^{-3/2}$, let $k = k(n)$ be a natural number and set $\lambda_k = \lambda_k(n) = \mathbb{E}(X_k) = n \cdot b(k; n-1, p)$. Then the following assertions hold.*

- If $\lim_{n \rightarrow \infty} \lambda_k(n) = 0$, then $\lim_{n \rightarrow \infty} \mathbb{P}(X_k = 0) = 1$.
- If $\lim_{n \rightarrow \infty} \lambda_k(n) = \infty$, then $\lim_{n \rightarrow \infty} \mathbb{P}(X_k > t) = 1$ for every fixed t .
- If $0 < \lim_{n \rightarrow \infty} \lambda_k(n) < \infty$, then X_k asymptotically behaves like a Poisson r.v. with mean λ_k :

$$P(X_k = r) \sim e^{-\lambda_k} \cdot \lambda_k^r / r!$$

for every fixed r .

The $\epsilon n^{-3/2} \leq p = p(n) \leq 1 - \epsilon n^{-3/2}$ hypothesis is to rule out when we consider a loose upper bound on the expected degree of X_k , if $pn^2 \rightarrow \infty$ then almost every $G \in \mathcal{G}(n, p)$ consist of independent edges and isolated vertices.

The first assertion comes directly from Markov's inequality and implies that if k is a finite fixed number and $\lim \lambda_k(n) = 0$ then a.a.s. there are no vertices of degree k . In the second case, there are an infinite number of vertices with degree k . In the third case, we can describe explicitly the degree distribution. The complete arguments for proving this theorem can be found in [8, Bollobás, p. 61].

2.4.3 Clique number

As outlined for the Radó graph model, the clique number is a highly restrictive property, for the Erdős Rényi model it is not different. Let X_r be the random variable that counts the number of r -cliques in a graph G , we are looking for a threshold were:

$$\lim_{n \rightarrow \infty} \mathbb{E}(X_r) > 0 \text{ and } \lim_{n \rightarrow \infty} \mathbb{E}(X_{r+1}) = 0.$$

But this is not possible if r is a fixed finite number. The closest we can get is stated in the following result:

Theorem 2.4.4. *Let $r = r(n) = O(n^{1/3})$ and let $p = p(n)$, $0 < p < 1$, be such that*

$$\binom{n}{r} p^{\binom{r}{2}} \rightarrow \infty \text{ and } \binom{n}{r+1} p^{\binom{r+1}{2}} \rightarrow 0,$$

then a.e $G \in \mathcal{G}(n, p)$ has clique number r .

This theorem can be proven using the calculations for $\mathbb{E}(X_r)$ and a first moment argument. A full proof can be found in [8, Bollobás, p. 290].

Given that $r(n)$ must grow along with n and $r = 3g - 3 + m$, this implies that the curve graph corresponds to a surface with infinite genus or with an infinite number of punctures.

2.4.4 Diameter

The diameter of a graph G , denoted by $\text{diam}(G)$, is the maximal distance between pairs of vertices of G .

If we want to model a surface of finite genus we must ensure infinite diameter, there are a number of theorems that describe the conditions under which this can be achieved [8, Bollobás, p. 259].

Following the past result, we now must guarantee that the diameter is equal to 2. The idea is to have an analog of Bering and Gaster result (Theorem 2.3.1).

Theorem 2.4.5. *Let d be a fixed integer, if*

$$\frac{(pn)^{d-1}}{n} \rightarrow 0 \text{ and } \frac{(pn)^d}{n} \rightarrow \infty$$

then, a.a.s $G \in \mathcal{G}(n, p)$ has diameter d

The proof of this theorem is due to Klee and Larman and can be found in [22, Klee, Larman 81]. For $d = 2$ this means we have that $p(n) \rightarrow 0$ and $p^2 n \rightarrow \infty$, i.e $p(n) = \frac{f(n)}{n^{1/2}}$ where $f(n) \in o(n^{1/2})$ and $f(n) \rightarrow \infty$.

Using the expression for $p(n)$ in 2.4.2, we must have $a < \frac{1}{2}$ to ensure diameter 2.

To conclude, the following result states that Erdős-Rényi graphs, that asymptotically approach the Radó graph, naturally satisfies the condition for the diameter. This gives us some insight into the interconnection between the object of study and the models when an asymptotic context is taken; the diameter condition of the curve graph appears naturally when modeling a surface of infinite genus.

Theorem 2.4.6. *If p is taken fixed $G(n, p)$ has diameter 2 with high probability.*

Proof. Let X_n be the random variable that counts the number of vertex pairs in a graph in $\mathcal{G}(n, p)$ with no common neighbors. By Markov's inequality we have that

$$\begin{aligned}
\mathbb{P}(X_n \geq 1) \leq \mathbb{E}(X_n) &= \binom{n}{2} \cdot \mathbb{P}(\text{Two vertices don't have common neighbors}) \\
&= \binom{n}{2} (1 - p^2)^{n-2},
\end{aligned}$$

where $\lim_{n \rightarrow \infty} \binom{n}{2} (1 - p^2)^{n-2} = 0$. \square

2.5 Conclusions

Theorems 2.4.1, 2.4.2, and 2.4.5 give the thresholds where each of the properties of the curve graph of infinite genus surface are satisfied. Theorem 2.4.4 describes the asymptotic behavior of the clique number.

In summary, if $\epsilon, a > 0$ are fixed real numbers and $\omega(n)$ a function that tends to infinity arbitrary slow, we have:

1. $p \geq \frac{\log(n) + \omega(n)}{n} \implies$ a.a.s $G \in \mathcal{G}(np)$ is connected.
2. $p = \frac{\epsilon}{n^a}$ with $a < 1 \implies$ a.a.s $G \in \mathcal{G}(np)$ is locally infinite.
3. $p = \frac{f(n)}{n^{1/2}}$ with $f(n) \in o(n^{1/2})$ and $f(n) \rightarrow \infty$, particularly if $p(n) = \frac{\epsilon}{n^a}$ $a < \frac{1}{2} \implies$ a.a.s $G \in \mathcal{G}(np)$ has diameter 2.
4. $r = r(n) = O(n^{1/3})$ with $\binom{n}{r} p^{\binom{r}{2}} \rightarrow \infty$ and $\binom{n}{r+1} p^{\binom{r+1}{2}} \rightarrow 0$ implies a.a.s $G \in \mathcal{G}(n, p)$ has clique number r .

For the simplified form of $p(n) = \frac{\epsilon}{n^a}$ we can plot the following thresholds:




FIGURE 2.6: Thresholds for the properties of the curve complex.

Some particular properties of the random graphs, such as a fixed clique number, will be linked to the asymptotic behavior of the vertices. Meanwhile, some conditions, like diameter 2, are strong enough that other properties come along.

The clique number property is highly restrictive when setting a generic model, thus different techniques must be implemented. The problem with this approach is that we will end up with a model so complicated that it will not be suitable for studying further phenomena.

There is a lot of progress in the study of random clique complexes that can help understand better the curve complex for infinite genus surfaces, for example in [21, Khale,09] we find the following result:

Theorem 2.5.1. *If $p = n^a$, with $a < -1/k$ or $a > -1/2k + 1$, then the k -th homology group of $X(G(n, p))$ is almost always vanishing, and if $-1/k < a < -1/(k + 1)$, then it is almost always non vanishing.*

Although interpretation might not be direct, the thresholds are within those defined for $C(S)$.

Studying a concept from a stochastic perspective could be helpful to understand its particularities. Also, the challenges to explain a phenomenon in a specific framework, although it can be defined in a general manner.

From the Gaster and Bering result, we can also conclude that some objects, with apparently very particular constructions, end up being very generic.

Using the Erdős-Rényi model we can bring insightful results of the asymptotic behavior of certain properties that in the Radó graph are given immediately.

Understanding the probabilistic nature of anything, in which computational operations are involved, can be helpful when optimizing procedures. This is explained in the next chapter.

Chapter 3

Computational experimentation

Nowadays Scientific Computing is one of the most important tools that we have for Stochastic means. It becomes crucial when the problem cannot be solved by traditional experimentation or theoretical means. There are many reasons why this might happen, for example whenever experimentation may be dangerous, too expensive, or time-consuming.

In this chapter, we use the outlined probabilistic material to accomplish an efficient implementation of the rigidity phenomenon in graphs. We describe our results with their technical difficulties and the actions taken to endure them.

All the computational experimentation was developed in `python`. We used `NetworkX` library to create and modify graphs.



3.1 Simulating Erdős-Rényi random graphs

There is a direct algorithm to obtain a graph in $G(n, p)$, it simulates a *Bernoulli*(p) r.v. for each of the $\frac{n(n-1)}{2}$ possible edges. Thus, it runs in $O(n^2)$ time.

It is possible to execute faster algorithms for small values of p . It runs in $O(n + m)$ time, where m is the expected number of edges, which equals to $\frac{pn(n-1)}{2}$. This is the one that we use for generating all the graphs in our executions. Details in performance and accuracy can be found in [6, Batagelj, Brandes 05].

Visual aid is helpful while writing the code for the experimentation. In Figure 3.1 appears a set of graphs obtained with the built-in algorithms for Erdős-Rényi graphs, fixing $n = 10$ and varying the parameter p .

Figure 3.2 shows the execution times when varying n .

FIGURE 3.1: Erdős-Rényi random graphs with n fixed and varying p .FIGURE 3.2: Execution times varying n . Normal and log scales.

3.2 Rigid expansions algorithms

A priori, the algorithm to determine a rigid expansion is supposed to be executed in a large amount of time. As the definition lets us see, it depends on the size of G , and exponentially on the size of A ; it must check among all possible subsets of A , that is 2^k verifications, where $|A| = k$. Thus, it is important to do some optimizations and evaluate when they have more impact on the expected execution time according to the parameters taken.

The following is the straightforward algorithm for rigid expansions.

Rigid Expansions Algorithm

Input: Random graph G (dictionary),
 set of vertices A (array).
 Output: Set of vertices obtained after expanding A (array).

1. Initialize N as empty (the set of new vertices).
2. For every B , subset of A :
 If $\bigcap_{b \in B} N(b) = v$ and $v \notin A \cup N$:
 Add v to N .
3. If N is not empty:
 Replace A by $A \cup N$ and return to step 1.
 Otherwise:
 Return A .

To optimize memory in step two the iterations were indexed by **generators**.

For time-execution optimizations, we implemented the following:

1. **Consideration of isolated vertices and leaves.** None of the isolated vertices in G have any influence in rigid expansions, so they should not be considered. Also, whenever A contains a leaf, it is convenient to ignore them; the unique neighbor of a leaf, which we will call *petioles* should be automatically added in the first expansion and then it does not contribute to uniquely determine new vertices. This means that the input should be replaced with:

$$A' = A \cup \{u : \exists x, N(x) = \{u\}\} - \{v : \deg(v) \leq 1\},$$

and add them again by the end of the expansions. This will be particularly helpful for small values of p .

2. **Relative size of A .** In Step 2, if A is big enough, it is faster to check if a vertex outside of A can be uniquely determined by a subset of A . This can reduce dramatically the execution time when p is small; it reduces the size of revisions by

taking only the *effective* part of A , this is convenient to do whenever

$$k \cdot \log(2) > \log(n - k) + (kp) \cdot \log(2),$$

where k is the size of A .

3. **Restriction to effective subsets.** Calculations in chapter 2 showed that some subsets are more likely to generate rigid expansions than others. This depends on the parameters of the space and the size of the subsets. If we restrict verifications to these effective subsets we can reduce the number of verifications.

With these optimizations we obtain the following algorithm:

Optimized Rigid Expansions Algorithm

Input: Random graph G (dictionary),
 set of vertices A (array),
 $n(\text{int})$ and $p(\text{float})$.

Output: Set of vertices obtained after expanding A (array).

1. Remove isolated vertices and replace A with A' .
2. Calculate the range of effective subsets.
 If $k \cdot \log(2) > \log(n - k) + (kp) \cdot \log(2)$:
 For every $v \in V - A$:
 Take $C = A \cap N(v)$ and for every B , effective subset of C :
 If $\bigcap_{b \in B} N(b) = v$ and $v \notin A \cup N$:
 Add v to N .
 Otherwise:
 For every B , effective subset of A :
 If $\bigcap_{b \in B} N(b) = v$ and $v \notin A \cup N$:
 Add v to N .
3. If N is not empty:
 Replace A with $A \cup N$, initialize N as empty and return to step 3.
 Otherwise:
 Return $A \cup \{v : \deg(v) \leq 1\}$.

3.3 Time execution comparison

The task of finding the first rigid expansion of $A \subset V$ of size k in $G \in \mathcal{G}(n, p)$ depends on n, p, k . To keep track of the enhancements implemented we measured the execution time, varying the parameters.

For each collection of n, p , and k we calculated the mean execution time for 30 different rigid expansions, with and without optimizations. We took k in some proportion of n , explicitly: $1/4$, $1/2$ and $3/4$.

Having a larger n impacts heavily when executing the non-optimized algorithm and even in certain thresholds the enhancement algorithms still take too much time. Also, considering the nature of rigid expansions and that we must execute multiple tests for each collection of parameters, n is fixed to be small, 20 in this case.

Results are presented in Figure 3.3.




FIGURE 3.3: Mean execution time varying n, p and k for 30 rigid expansions. Measured in ms.

Notice that these enhancements have an important impact on reducing the execution time. The results presented in Chapter 2 were obtained with these optimizations, so we can also conclude that accuracy is not compromised.

We expect to see an exponential behavior for the execution time. To have a better comparison we can use the logarithmic scale.

Results in log scale are presented in Figure 3.4




FIGURE 3.4: Mean execution time varying n, p and k for 30 rigid expansions. Measured in ms. log scale.

Notice that optimizations have more impact for lower values of p , this sounds reasonable given that the first-proposed optimizations are explicitly helpful for sparse graphs.

Increasing k has a big impact on performance that corresponds with still having to search among an exponentially bigger number of subsets. Even using the second and third optimization we are still obtaining the same exponential behavior.

Further optimizations can be implemented, such as excluding the parts of G that are not connected with vertices of A , nevertheless, this will only have an impact in sparse graphs as well.

3.4 Conclusions

We could not have done experiments on larger graphs without probabilistic optimizations. The understanding of the combinatorial nature of rigid expansions was also key.

In general, the use of probability theory in computational experiments has demonstrated to be powerful and efficient. Algorithms based on random sampling provide state-of-the-art techniques due to their great degree of flexibility and reliability.

To name a few:

- The PageRank algorithm was the first method used by Google to order search results [26, Page 99]. It outputs a probability distribution used to represent the likelihood that a person randomly clicking on links will arrive at any particular page.
- In motion planning, the use of Rapidly-exploring random trees (RRTs) are one of the most successful algorithms [1, Alcazar 15]. Problems in motion planning consist of finding a collision-free path that connects an initial configuration of geometric bodies to a final goal configuration. A RRT is a rooted tree that grows from a starting configuration by using random samples from the search space.

But it also works in the opposite direction, the use of computational tools can bring value for theoretical means. For theoretical means it has allowed, for example, to verify whether the established conditions in a probabilistic model are sharp enough ([5, Aronshtam 13]).

Bottom line, the use of computational tools can be very helpful to understand a topic even in the most theoretical contexts.

Appendix A

The Generate algorithm

In this section we describe the algorithm due to [9, Broder 89]. Given a not directed graph G with n vertices, it produces a maximal tree of G sampled uniformly among all the possible ones. For almost every graph the expected executed time of the algorithm is $O(n \cdot \log(n))$ for each tree and $O(n^3)$ in the worst cases.

One of the first algorithms published for this problem has execution time $O(n^5)$. It is based on the fact that the total number of directed trees in a graph can be explicitly calculated through a determinant of $n \times n$ size. The algorithm considers the edges of the graph labeled from 1 to m , each maximal tree is labeled by the set of its edges. This induces a lexicographic order in the set of trees and the same tree can be found by calculating at most m determinants. Further improvements by [12, Colbourn 89] reduce the number of calculations, thus reducing the execution time to $O(n^3)$ or $O(L(n))$, where $L(n)$ is the execution time of multiplying matrices of size $n \times n$, but the new algorithms turn out to be far more complicated.

For a stochastic approach, consider a particle that moves among vertices in a graph. At each step it moves, choosing uniformly random, from the current vertex to a neighbor of it. This stochastic process is a Markov chain called **Random Walk**.

Generate AlgorithmInput: Graph G (dictionary),Output: Maximal tree T (dictionary)

1. Choose a random vertex s of G (uniformly).
2. Simulate a simple random walk in G . It stops when every vertex gets visited.
3. For each i in $V - s$ collect the edge (j, i) , the first entrance corresponds to the vertex where the particle was before it visited for the first time the vertex i . Let T be the collection of such edges.
4. Return T .

T is a maximal tree because it contains $|V| - 1$ edges; it has an edge for every vertex in G except for s , and by construction it does not contains cycles.

The **Generate** algorithm is based on a simulation of Markov chains in the space of interest. In this case, the Markov chain has a stationary distribution $\pi_i = d_i / \sum_{j \in V} d_j$ where d_i is the degree of the vertex i . The pounted digraph associated to this chain $G_M = (V, E')$, is obtained by replacing each edge $\{i, j\} \in A$ by two directed edges; (i, j) with weight $1/d_i$ and (j, i) with weight $1/d_j$. The justification that the algorithm actually provides a method to sample with uniform distribution is summarized in the next three results, their proofs can be found in [9, Broder 89].

Let $\mathcal{T}_i(G_M)$ be the family of maximal directed trees of G_M with root i , when the root is not under consideration it will be denoted simply by $\mathcal{T}(G_M)$.

Theorem A.0.1. *Let M be a irreducible Markov chain in n states with stationary distribution π_1, \dots, π_n . Let G_M be the weighted digraph associated to M . Then*

$$\pi_i = \frac{\sum_{T \in \mathcal{T}_i(G_M)} \omega(T)}{\sum_{T \in \mathcal{T}(G_M)} \omega(T)}$$

where $\omega(T) = \prod_{a \in A(T)} \omega(a)$, this means that the weight of the a directed tree is defined as the product of the weight of the edges of the tree.

We define the (*forward tree*) at time t , F_t as follows: Let I_t be the set of states visited before time $t + 1$. For every $i \in I_t$, let $p(i, t)$ be the first time that the state i was visited.

The root of the tree F_t is $\{(X_{p(i,t)}, X_{p(i,t)-1}) | i \in I_t - X_0\}$, where $(X_t)_{t \in \mathbb{N}}$ corresponds to the Markov chain given by the random walk. In other words F_t is constructed by overlapping the first entrance at each state with inverted orientation. Clearly F_t is a directed tree with root where each edge points from the leaves to the root.

Let C be the *covering time*, i.e. the first time that all the states were visited. Clearly for $t \geq C$ the tree F_t is a directed maximal tree and $F_t = F_C$. Note that with the past definition, the random walk $\{X_t\}$ in the vertices of G_M induces a Markov chain $\{F_t\}$ in the space of all directed trees of G_M , it is called forward trees chain.

For this chain, every non-maximal tree is a transitive state and every maximal tree is an absorbent state. Even more, the next theorem establishes the distribution of F_C .

Theorem A.0.2. *With the same notation and conditions of the past theorem. Let F_C be the forward tree in time C . Then, for any maximal directed tree with root T of G_M we have*

$$\mathbb{P}(F_C = T) = \frac{\prod_{(i,j) \in T} P_{i,j}}{\sum_{T' \in \mathcal{T}(G_M)} \prod_{(i,j) \in T'} P_{i,j}}.$$

Corollary A.0.2.1. *(Proof of the **Generate** algorithm) Let M be a simple random walk in a connected non-directed graph $G = (V, E)$ starting from a vertex s and G_M the directed graph associated to M and covering time C for G . Starting from the stationary distribution, we have that F_C , without considering direction, is a maximal tree of G with random uniform distribution among all the maximal possible trees of G .*

This algorithm can be implemented using **Python**. Fixing G as the complete graph K_n it was possible, using the generate algorithm, to sample uniformly from the set of maximal trees with n vertices. In figure A.1 appears a set of trees obtained with this method, which are drawn using the function **draw_random** of the **NetworX** library.

The expected execution time of the algorithm per tree is equal to $\mathbb{E}(C_s)$. It is known that for the connected graph $\mathbb{E}(C_v) = O(n^3)$, nevertheless in [10, Broder, Andrei 89] there is a proof that if the transition matrix of a random walk have the second greater eigenvalue bounded away from 1, then the expected covering time is only $O(n \cdot \log(n))$. Almost every graph in the Erdős-Rényi model satisfy this condition when $p > \frac{c \log(n)}{n}$, in particular when $p = \frac{1}{2}$ and for almost every d -regular graphs [17, Friedman 89]. In figure A.2 shows the results of the algorithm's execution time.




FIGURE A.1: Maximal tree chosen randomly with uniform distribution among all the possible ones in complete graph (10) of vertices.




FIGURE A.2: Execution time of the algorithm in seconds varying the size of the tree. It appears the normal and the logarithmic scale.

Bibliography

- [1] V. Alcázar, S. Fernández, D. Borrajo, and M. Veloso. Using random sampling trees for automated planning. *AI Commun.*, 28(4):665–681, 2015.
- [2] J. Aramayona and C. J. Leininger. Finite rigid sets in curve complexes. *J Topology Anal*, 5(2), 2012.
- [3] J. Aramayona and C. J. Leininger. Exhausting curve complexes by finite rigid sets. *Pacific J. Math.*, 282(2):257–283, 2016.
- [4] J. Aramayona and J. Souto. Rigidity phenomena in the mapping class group. In *Handbook of Teichmüller theory. Vol. VI*, volume 27 of *IRMA Lect. Math. Theor. Phys.*, pages 131–165. Eur. Math. Soc., Zürich, 2016.
- [5] L. Aronshtam, N. Linial, T. Łuczak, and R. Meshulam. Collapsibility and vanishing of top homology in random simplicial complexes. *Discrete Comput. Geom.*, 49(2):317–334, 2013.
- [6] V. Batagelj and U. Brandes. Efficient generation of large random networks. *Phys. Rev. E*, 71:036113, Mar 2005.
- [7] E. A. Bering, IV and J. Gaster. The random graph embeds in the curve graph of any infinite genus surface. *New York J. Math.*, 23:59–66, 2017.
- [8] B. Bollobás. *Random graphs*, volume 73 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 2001.
- [9] A. Broder. Generating random spanning trees. In *Proceedings of the 30th Annual Symposium on Foundations of Computer Science*, SFCS '89, pages 442–447, Washington, DC, USA, 1989. IEEE Computer Society.
- [10] A. Z. Broder and A. R. Karlin. Bounds on the cover time. *Journal of Theoretical Probability*, 2(1):101–120, Jan 1989.
- [11] P. J. Cameron. The random graph. In *The mathematics of Paul Erdős, II*, volume 14 of *Algorithms Combin.*, pages 333–351. Springer, Berlin, 1997.

- [12] C. J. Colbourn, R. P. J. Day, and L. D. Nel. Unranking and ranking spanning trees of a graph. *J. Algorithms*, 10(2):271–286, 1989.
- [13] A. Costa and M. Farber. Large random simplicial complexes, II; the fundamental group. *J. Topol. Anal.*, 9(3):441–483, 2017.
- [14] P. Erdős and A. Rényi. On random graphs. I. *Publ. Math. Debrecen*, 6:290–297, 1959.
- [15] P. Erdős and A. Rényi. Asymmetric graphs. *Acta Math. Acad. Sci. Hungar.*, 14:295–315, 1963.
- [16] B. Farb and D. Margalit. *A primer on mapping class groups*, volume 49 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 2012.
- [17] J. Friedman, J. Kahn, and E. Szemerédi. On the second eigenvalue of random regular graphs. In *Proceedings of the Twenty-first Annual ACM Symposium on Theory of Computing*, STOC '89, pages 587–598, New York, NY, USA, 1989. ACM.
- [18] A. Hatcher and W. Thurston. A presentation for the mapping class group of a closed orientable surface. *Topology*, 19(3):221–237, 1980.
- [19] J. Hernández Hernández. Exhaustion of the curve graph via rigid expansions. *Glasg. Math. J.*, 61(1):195–230, 2019.
- [20] N. V. Ivanov. Automorphism of complexes of curves and of Teichmüller spaces. *Internat. Math. Res. Notices*, (14):651–666, 1997.
- [21] M. Kahle. Topology of random clique complexes. *Discrete Mathematics*, 309(6):1658–1671, 2009.
- [22] V. Klee and D. Larman. Diameters of random graphs. *Canadian J. Math.*, 33(3):618–640, 1981.
- [23] M. Korkmaz. Automorphisms of complexes of curves on punctured spheres and on punctured tori. *Topology Appl.*, 95(2):85–111, 1999.
- [24] F. Luo. Automorphisms of the complex of curves. *Topology*, 39(2):283–298, 2000.
- [25] H. A. Masur and Y. N. Minsky. Geometry of the complex of curves. I. Hyperbolicity. *Invent. Math.*, 138(1):103–149, 1999.
- [26] L. Page, S. Brin, R. Motwani, and T. Winograd. The pagerank citation ranking: Bringing order to the web. Technical Report 1999-66, Stanford InfoLab, November 1999. Previous number = SIDL-WP-1999-0120.