

Title

Riccardo Buscicchio^{1,2,4a} and John Doe⁴

¹ Dipartimento di Fisica “G. Occhialini”, Università degli Studi di Milano-Bicocca, Piazza della Scienza 3, 20126 Milano, Italy

² INFN, Sezione di Milano-Bicocca, Piazza della Scienza 3, 20126 Milano, Italy

³ Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Straße 1, 85741 Garching, Germany

⁴ Institute for Gravitational Wave Astronomy & School of Physics and Astronomy, University of Birmingham, Birmingham, B15 2TT, UK

Received: date / Revised version: date

Abstract. An abstract

PACS. 04.80.Nn, 95.55.Ym Gravitational wave detectors – 95.85.Sz Gravitational wave astronomical observations

1 Introduction

Intro with a range using SI units $10^{-4} - 10^{-1}$ Hz. A citation Ref. [1].

The paper is organized as follows. In 1, we do such and such. Another type of citation [2]. Some dimensionful quantities 3 kpc. Some other 13.5 Gyr. Another quantity with scale factor $8.2 \times 10^{10} M_{\odot}$. A pure number 26×10^6 .

$$\dot{f} = \frac{96\pi^{8/3}}{5c^5} (G\mathcal{M}_c)^{5/3} f^{11/3}, \quad (1)$$

A percentage 1 % and some dimensionful ranges $f > 2 - 3$ mHz or $f < 2 - 3$ mHz.

2 Methodology




Fig. 1. A single column figure

^a riccardo.buscicchio@unimib.it

References

1. P. Amaro-Seoane, J. Andrews, M. Arca Sedda, A. Askar, Q. Baghi, R. Balasov, I. Bartos, S.S. Bavera, J. Bellovary, C.P.L. Berry et al., Living Reviews in Relativity **26**, 2 (2023), 2203.06016
2. S. Babak, C. Caprini, D.G. Figueroa, N. Karnesis, P. Marcoccia, G. Nardini, M. Pieroni, A. Ricciardone, A. Sesana, J. Torrado, J. Cosmology Astropart. Phys. **2023**, 034 (2023), 2304.06368