

A 2-SYSTOLIC INEQUALITY ON COMPACT KÄHLER SURFACES WITH POSITIVE SCALAR CURVATURE

ZEHAO SHA

ABSTRACT. In this note, we prove a 2-systolic inequality on compact positive scalar curvature Kähler surfaces admitting a nonconstant holomorphic map to a positive-genus compact Riemann surface.

1. INTRODUCTION

Systolic geometry studies the relationship between the minimal size of non-trivial homological cycles in a Riemannian manifold and its global geometric properties. Given a closed Riemannian manifold (M, h) , the k -systole is defined as

$$\text{sys}_k(M, h) := \inf \{ \text{Vol}_g(Z) \mid Z \subset M \text{ embedded}, [Z] \neq 0 \in H_k(M; \mathbb{Z}) \}.$$

Consider a Riemannian manifold (M, h) with positive scalar curvature (PSC for abbreviation), Schoen–Yau [SY79] confirmed that any area-minimizing surface in M is homeomorphic to either S^2 or \mathbb{RP}^2 . The \mathbb{RP}^2 case was studied in [BBN10], which proved the sharp upper bound of the area for an embedded area-minimizing projective plane,

$$\text{Area}_h(\Sigma) \cdot \min_M S_M \leq 12\pi$$

with equality if $M \simeq \mathbb{RP}^3$. In the S^2 case, Bray–Brendle–Neves [BBN10] established the following important rigidity result concerning the π_2 -systole and the minimum of the scalar curvature $\min S_M$ of a PSC 3-manifold (M, h) :

Theorem 1.1 (Bray–Brendle–Neves, [BBN10]). Let (M^3, h) be a closed, orientable Riemannian 3-manifold with positive scalar curvature. Then the following inequality holds:

$$\text{sys}_{\pi_2}(M, h) \cdot \min_M S_M \leq 8\pi. \tag{1.1}$$

Moreover, equality holds if and only if M^3 is isometrically covered by $S^2 \times S^1$ with the round metric on S^2 , product with the flat metric on S^1 .

The sharp bound in (1.1) was recently refined by Xu [Xu25], who proved that it is not only rigid but also exhibits a strict quantitative gap away from the model case. Further developments on the interplay between the 2-systole and positive scalar curvature include [Zhu20, Ric20, Ori25].

In [Ste22], Stern introduced the following inequality for a non-constant S^1 -valued harmonic map u on a 3-manifold (M, h) using the level set method,

$$2\pi \int_{\theta \in S^1} \chi(\Sigma_\theta) \geq \frac{1}{2} \int_{\theta \in S^1} \int_{\Sigma_\theta} (|du|^{-2} |\text{Hess}(u)|^2 + S_M) \quad (1.2)$$

and generalized the Bray–Brendle–Neves’ systolic inequality for the homological 2-systole.

Motivated by Stern’s approach, we adapt the level set method to compact Kähler surfaces fibred over a Riemann surface with positive genus, and obtain a Bray–Brendle–Neves type inequality for the homological 2-systole. In particular, we have the following:

Theorem 1.2. Let (X, ω) be a compact PSC Kähler surface admitting a non-constant holomorphic map $f : X \rightarrow C$ to a compact Riemann surface C with genus $g(C) \geq 1$. Then, we have

$$\min_X S_X \cdot \text{sys}_2(X, \omega) \leq 8\pi. \quad (1.3)$$

Moreover, the equality holds if and only if X is isometrically covered by $\mathbb{CP}^1 \times E$ equipped with the product of the standard Fubini–Study metric on \mathbb{CP}^1 , and a flat metric on E satisfying $\text{Vol}(E) \geq \pi$, where E is an elliptic curve.

A direct consequence of the above result is:

Corollary 1.3. Let (X, ω) be a compact PSC Kähler surface admitting a non-constant holomorphic map $f : X \rightarrow C$ to a compact Riemann surface C with genus $g(S) \geq 2$. Then, we have

$$\min_X S_X \cdot \text{sys}_2(X, \omega) < 8\pi. \quad (1.4)$$

It is worth-noting that a compact Kähler surface X admits a PSC metric if and only if X is obtained from \mathbb{P}^2 or $\mathbb{P}(E)$ by a finite sequence of blow ups, where E is a rank 2 holomorphic vector bundle over a compact Riemann surface: For minimal compact Kähler surfaces (which are not the blow-up of other Kähler surfaces), Yau [Yau74] showed that the existence of a Kähler PSC metric is equivalent to X is ruled or \mathbb{P}^2 (see also [LeB95]). LeBrun conjectured that this statement is still valid after allowing the blowup. The remaining gap—whether blowing up preserves the *sign* of the scalar curvature was settled recently by Brown, who proved that Kähler blow-ups preserve the sign of scalar curvature and completed the classification [Bro24, Thm B].

Acknowledgment. The majority of this work was conducted while the author was a PhD student at Institut Fourier. The author is grateful for the support and resources provided by Institut Fourier during this period. The author also extends his gratitude to his advisor, Professor Gérard Besson, for introducing him to the field of PSC manifolds.

2. THE KÄHLER SURFACE FIBRED OVER A RIEMANN SURFACE WITH POSITIVE GENUS

Let $f : (X^n, \omega) \rightarrow C$ be a non-constant holomorphic map from a compact Kähler manifold to a compact Riemann surface C with genus $g \geq 1$. Thanks to the uniformization theorem, we can always choose a metric ω_0 on C with non-positive Gauss curvature.

Taking $z \in C$ be any point, then we have $f^{-1}(z) := D_z$ is a Cartier divisor in X with complex codimension 1. Indeed, D_z defines a line bundle $\mathcal{O}(D_z)$ with the first Chern class $c_1(\mathcal{O}(D_z))$ represented by $f^*\omega_0$ after normalization. In the subsequent part, we only consider the smooth part of D_z , and denote it by D_z directly.

Recall the adjunction formula for a smooth divisor D , the canonical bundle K_D satisfies

$$K_D = (K_X \otimes \mathcal{O}(D))|_D. \quad (2.1)$$

Since D is the fiber of a holomorphic map over C , the normal bundle $\mathcal{N}_D \cong \mathcal{O}(D)|_D$ is trivial. By taking the first Chern class of the adjunction formula, we obtain

$$c_1(D) = c_1(X)|_D,$$

which implies

$$\text{Ric}_D(\omega) = \text{Ric}_X(\omega)|_D. \quad (2.2)$$

Denote ν by the unit normal vector field of D of type $(1, 0)$, we obtain the traced Gauss equation

$$S_D(\omega) = \text{tr}_\omega \text{Ric}_D(\omega) = \text{tr}_\omega \text{Ric}_X(\omega)|_D = S_X(\omega) - \text{Ric}_X(\omega)(\nu, \bar{\nu}).$$

Moreover, since $\nu = \nabla^{1,0}f/|\nabla^{1,0}f|$, we obtain

$$\text{Ric}_X(\omega) (\nabla^{1,0}f, \nabla^{0,1}f) = |\nabla^{1,0}f|^2 (S_X(\omega) - S_D(\omega)). \quad (2.3)$$

Recall the Bochner formula for holomorphic maps, and the co-area formula:

Lemma 2.1 (Bochner formula). Let $f : (X, \omega) \rightarrow (N, \tilde{\omega})$ be holomorphic, then

$$\Delta|\partial f|^2 = |\nabla\partial f|^2 + \langle \text{Ric}(\omega), f^*\tilde{\omega} \rangle - \text{tr}_\omega^2(f^* \text{Rm}(\tilde{\omega})). \quad (2.4)$$

where $\text{Ric}(\omega)$ is the Ricci form of X , and $\text{Rm}(\tilde{\omega})$ is the curvature form of N .

Lemma 2.2 (Co-area formula). Let (X^n, ω) be a compact Kähler manifold and let (C, ω_0) be a compact Riemann surface such that

$$\int_C \omega_0 = \frac{1}{n},$$

after normalization. Then for any $g \in C^\infty(X)$ and $z \in E$ regular value of f , we have

$$\int_X g \omega^n = \int_C \left(\int_{f^{-1}(z)} \frac{g}{|\partial f|^2} \omega^{n-1} \right) \omega_0. \quad (2.5)$$

When $(N, \tilde{\omega}) = (C, \omega_0)$, we have

$$\Delta|\partial f|^2 \geq |\nabla\partial f|^2 + \text{Ric}_X(\omega) (\nabla^{1,0}f, \nabla^{0,1}f),$$

with equality holds if $\text{Rm}(\omega_0) \equiv 0$. Combining with the traced Gauss equation (2.3), we obtain

$$\Delta|\partial f|^2 \geq |\nabla\partial f|^2 + |\nabla^{1,0}f|^2 (S_X(\omega) - S_D(\omega)). \quad (2.6)$$

Combining with the co-area formula and (2.6), we can then see the following identity:

Lemma 2.3. Let (X^n, ω) be a compact Kähler manifold and let (C, ω_0) be a compact Riemann surface with genus $g \geq 1$ endowed with a constant curvature metric ω_0 . Suppose that $f : X \rightarrow C$ is a non-trivial holomorphic map. Then for any $\phi \in C^\infty(X)$, we have

$$\int_C \left[\int_{D_z} \phi^2 \left(\frac{|\nabla \partial f|^2}{|\partial f|^2} + S_X - S_{D_z} \right) \omega^{n-1} \right] \omega_0 \leq -n \int_X \sqrt{-1} \partial(\phi^2) \wedge \bar{\partial} |\partial f|^2 \wedge \omega^{n-1}. \quad (2.7)$$

Moreover, the equality holds if and only if (S, ω_0) is an elliptic curve with a flat metric.

Proof. Let $C = A \cup B$, where A contains the set for all critical values of f . Then, multiplying any smooth function ϕ^2 on the both sides and integrating over $f^{-1}(B)$, we have

$$\int_{f^{-1}(B)} \phi^2 \left[|\nabla \partial f|^2 + |\nabla^{1,0} f|^2 (S_X(\omega) - S_{D_z}(\omega)) \right] \omega^n \leq \int_{f^{-1}(B)} \phi^2 (\Delta |\partial f|^2) \omega^n. \quad (2.8)$$

Indeed, the right-hand side of (2.8) gives

$$\int_{f^{-1}(B)} \phi^2 (\Delta |\partial f|^2) \omega^n = n \int_{f^{-1}(B)} \phi^2 \sqrt{-1} \partial \bar{\partial} |\partial f|^2 \wedge \omega^{n-1}$$

We then apply the co-area formula to the left-hand side of (2.8), which yields

$$\begin{aligned} & \int_{f^{-1}(B)} \phi^2 \left[|\nabla \partial f|^2 + |\nabla^{1,0} f|^2 (S_X - S_{D_z}) \right] \omega^n \\ &= n \int_B \left[\int_{D_z} \left(\frac{|\nabla \partial f|^2}{|\partial f|^2} + S_X - S_D \right) \omega^{n-1} \right] \omega_0 \end{aligned}$$

By Sard's theorem, we can take the measure of A arbitrarily small, and this gives

$$n \int_X \phi^2 \sqrt{-1} \partial \bar{\partial} |\partial f|^2 \wedge \omega^{n-1} = -n \int_X \sqrt{-1} \partial(\phi^2) \wedge \bar{\partial} |\partial f|^2 \wedge \omega^{n-1}.$$

Consequently, we have

$$\int_C \left[\int_{D_z} \phi^2 \left(\frac{|\nabla \partial f|^2}{|\partial f|^2} + S_X - S_{D_z} \right) \omega^{n-1} \right] \omega_0 \leq -n \int_X \sqrt{-1} \partial(\phi^2) \wedge \bar{\partial} |\partial f|^2 \wedge \omega^{n-1},$$

which implies the desired equality. \square

3. THE 2-SYSTOLE IN KÄHLER SURFACE

This section is devoted to the study of the (homological) 2-systole in Kähler surfaces. We begin by recalling the fundamental definition of the k -systole in Riemannian geometry. Let (M, h) be a closed Riemannian manifold of dimension $n \geq k$. The k -systole $\text{sys}_k(M, h)$ is defined as the infimum of the volumes of all integral k -cycles representing nontrivial homology classes:

$$\text{sys}_k(M, h) := \inf \{ \text{Vol}_h(Z) \mid Z \subset M \text{ embedded}, [Z] \neq 0 \in H_k(M; \mathbb{Z}) \}.$$

In the context of Kähler geometry, additional structure enriches this concept. Let (X, ω) be a compact Kähler surface. The 2-systole is the least area among nonseparating real surfaces in X .

Definition 3.1 (2-systole in Kähler surfaces). For a compact Kähler surface (X, ω) , the 2-systole can be defined by

$$\text{sys}_2(X, \omega) = \inf \{ \text{Vol}_\omega(Z) \mid Z \subset X \text{ embedded}, [Z] \neq 0 \in H_2(X; \mathbb{Z}) \}.$$

The following result gives a Bray-Brendle-Neves type inequality [BBN10] for 2-systole for compact PSC Kähler surfaces over a Riemann surface with genus $g \geq 1$.

Theorem 3.2. Let (X, ω) be a compact Kähler surface admitting a non-constant holomorphic map $f : X \rightarrow C$ to a complex curve C with genus $g(C) \geq 1$. Then, we have

$$\min_X S_X \cdot \text{sys}_2(X, \omega) \leq 8\pi. \quad (3.1)$$

Moreover, the equality holds if and only if X is covered by $\mathbb{CP}^1 \times E$ equipped with the product of the Fubini–Study metric on \mathbb{CP}^1 with $\text{Vol}(\mathbb{CP}^1) = \pi$, and a flat metric on E so that the $\text{sys}_2(X, \omega)$ is achieved by the \mathbb{CP}^1 -fiber, where E is an elliptic curve.

Proof. By taking $\phi = 1$ in (2.7), we obtain

$$\int_C \left[\int_{D_z} \left(\frac{|\nabla \partial f|^2}{|\partial f|^2} + S_X \right) \omega \right] \omega_0 \leq \int_C \left(\int_{D_z} S_{D_z} \cdot \omega \right) \omega_0.$$

It follows from the Gauss-Bonnet formula,

$$\begin{aligned} 4\pi \int_C \chi(D_z) \omega_0 &= \int_C \left(\int_{D_z} S_{D_z} \cdot \omega \right) \omega_0 \\ &\geq \int_C \left(\int_{D_z} S_X \cdot \omega \right) \omega_0 \\ &\geq \min_X S_X \cdot \int_C \text{Vol}_\omega(D_z) \omega_0. \end{aligned}$$

Denote $N(z)$ by the number of the homological non-zero irreducible components of D_z , we then have

$$\text{Vol}_\omega(D_z) \geq N(z) \cdot \text{sys}_2(X, \omega).$$

Meanwhile, we have

$$\chi(D_z) \leq 2N(z).$$

Thus, we have

$$\begin{aligned} 8\pi \int_C N(z) \omega_0 &\geq 4\pi \int_C \chi(D_z) \omega_0 \\ &\geq \min_X S_X \cdot \int_C \text{Vol}_\omega(D_z) \omega_0 \\ &\geq \min_X S_X \cdot \text{sys}_2(X, \omega) \int_C N(z) \omega_0, \end{aligned}$$

which gives the desired result. The equality holds in case C admits a flat metric and ∇f is parallel along D_z , with each irreducible component of D_z is \mathbb{CP}^1 . \square

We finally see a simple but interesting example:

Example 3.3. Let $X = \mathbb{P}^1 \times C$ be a compact complex surface, where C is a compact Riemann surface of genus $g \geq 2$. Equip X with the product Kähler metric

$$\omega = \omega_{\text{FS}} \oplus \omega_C,$$

where on \mathbb{P}^1 we take the Fubini–Study metric normalized by

$$\text{Vol}_{\omega_{\text{FS}}}(\mathbb{P}^1) = \pi, \quad S_{\mathbb{P}^1} = 8,$$

and on C we choose a constant scalar curvature metric with

$$S_C = -8 + \varepsilon \quad \text{for some } \varepsilon \in (0, 8).$$

Then the product scalar curvature $S_X = S_{\mathbb{P}^1} + S_C = 8 + (-8 + \varepsilon) = \varepsilon$ is constant, hence $\min_X S_X = \varepsilon$ and X has positive scalar curvature.

Next, compare the areas of the two basic complex curves:

- For the \mathbb{P}^1 -fiber $F = \mathbb{P}^1 \times \{p\}$, calibration by ω gives $\text{Vol}_\omega(F) = \pi$.
- For the C -fiber $C_p = \{q\} \times C$, Gauss–Bonnet formula yields

$$\int_C K_C \omega = 2\pi\chi(C) = 2\pi(2 - 2g) = -4\pi(g - 1).$$

Then we obtain

$$\text{Vol}_\omega(C_p) = \frac{8\pi(g - 1)}{8 - \varepsilon}.$$

For $g \geq 2$ and $\varepsilon \in (0, 8)$ one has $\text{Area}_\omega(C_p) > \pi$, so the 2-systole is realized by the \mathbb{P}^1 -fiber:

$$\text{sys}_2(X, \omega) = \min \{ \text{Area}_\omega(F), \text{Area}_\omega(C_p) \} = \pi.$$

Consequently,

$$\min_X S_X \cdot \text{sys}_2(X, \omega) = \varepsilon \cdot \pi < 8\pi.$$

In particular, this product is independent of the genus g , and it can be made arbitrarily close to 8π by letting $\varepsilon \uparrow 8$.

4. DISCUSSION

We have introduced the BBN-type 2-systolic inequality for PSC Kähler surfaces admitting a non-constant holomorphic map to a base curve with positive genus. The remaining natural case is the *rational base* $C \simeq \mathbb{P}^1$.

Question 4.1 (Rational base). Let (X, ω) be a compact PSC Kähler surface admitting a non-constant holomorphic map $f : X \rightarrow \mathbb{P}^1$. Then

$$\min_X S_X \cdot \text{sys}_2(X, \omega) \leq 16\pi,$$

with equality realized by the product $(\mathbb{P}^1 \times \mathbb{P}^1, \omega_{\text{FS}} \oplus \omega_{\text{FS}})$ normalized by $\text{Vol}_{\omega_{\text{FS}}}(\mathbb{P}^1) = \pi$ on each factor.

More ambitiously, we expect the same universal bound without assuming a fibration:

Question 4.2 (Universal bound). Every compact PSC Kähler surface satisfies

$$\min_X S_X \cdot \text{sys}_2(X, \omega) \leq 16\pi.$$

This broader form is motivated by surfaces such as \mathbb{P}^2 , which do not admit nonconstant holomorphic maps onto \mathbb{P}^1 (nor onto positive-genus curves). Explicit computations in standard models remain $\leq 16\pi$. At present, we are not aware of any example exceeding 16π .

REFERENCES

- [BBEN10] H. Bray, S. Brendle, M. Eichmair, and A. Neves, *Area-minimizing projective planes in three-manifolds*, Comm. Pure Appl. Math. **63** (2010), no. 9, 1237–1247.
- [BBN10] H. Bray, S. Brendle, and A. Neves, *Rigidity of area-minimizing two-spheres in three-manifolds*, Comm. Anal. Geom. **18** (2010), no. 4, 821–830.
- [Bro24] G. M. Brown, *The sign of scalar curvature on Kähler blowups*, arXiv:2405.12189 (2024).
- [LeB95] C. LeBrun, *On the scalar curvature of complex surfaces*, Geom. Funct. Anal. **5** (1995), no. 3, 619–628.
- [Ori25] Shunichiro Orikasa, *Systolic inequality and scalar curvature*, arXiv:2509.17376 (2025).
- [Ric20] T. Richard, *On the 2-systole of stretched enough positive scalar curvature metrics on $S^2 \times S^2$* , SIGMA. Symmetry Integrability Geom. Methods Appl. **16** (2020).
- [Ste22] D. Stern, *Scalar curvature and harmonic maps to S^1* , J. Differential Geom. **122** (2022), no. 2, 259–269.
- [SY79] R. Schoen and S.-T. Yau, *Existence of incompressible minimal surfaces and the topology of three-dimensional manifolds with non-negative scalar curvature*, Ann. Math. **110** (1979), no. 1, 127–142.
- [Xu25] K. Xu, *A topological gap theorem for the π_2 -systole of positive scalar curvature 3-manifolds*, Duke Math. J. **174** (2025), no. 8, 1647–1664.
- [Yau74] S.-T. Yau, *On the curvature of compact Hermitian manifolds*, Invent. Math. **25** (1974), no. 3, 213–239.
- [Zhu20] Jintian Zhu, *Rigidity of area-minimizing 2-spheres in n -manifolds with positive scalar curvature*, Proc. Amer. Math. Soc. **148** (2020), no. 8, 3479–3489.

INSTITUTE FOR MATHEMATICS AND FUNDAMENTAL PHYSICS, 1899 SHILIAN SOUTH ROAD, 230088 HEFEI, CHINA

Email address: zhsha0251@gmail.com