

Mathematical Inequalities   Statistics (academic discipline)

# What is a real-world application of Chebyshev's inequality?



**There's more on Quora...**

Pick new people and topics to follow and see the best answers on Quora.

Update Your Interests

## Related Questions

- What is the relevance of the Chebyshev's inequality in real life application?
- What is an intuitive explanation of Chebyshev's inequality?
- Mathematics: What are some intuitive proofs of Markov's inequality and Chebyshev's inequality?
- What are some real-world applications of MCMC?
- How tight of a bound is Chebyshev's inequality?
- Where are inequalities used in real life?
- Why is the Cauchy-Schwarz inequality considered to be one of the most important inequalities in mathematics? What are its various applications...
- Do I use Chebyshev's inequality for this question?
- What are the real world applications/uses of econometrics?
- What are main applications of Jensen's Inequality?

+ Ask New Question

More Related Questions

## Question Stats

13 Public Followers  
4,915 Views  
Last Asked Jan 4, 2016  
Edits

Suppose you are an **insurance** company. You receive **claims** of random sizes at random times from your customers. Based on the claims you have received so far, you want to get an idea about how large the claims are likely to be in the future, so you can set aside enough reserves to cover the claims of the upcoming calendar year, say.

To keep things simple, you assume that the claims are independent and identically distributed. So each claim is assumed to be a random draw from a single, unknown distribution.

Without knowing anything about this underlying distribution, Chebyshev's inequality immediately tells you that you can be at least 90% sure that future claims will be within three standard deviations away from the mean. You can get pretty good estimates of the mean and the standard deviation from the claims you have observed so far. Once you have those estimates, you immediately have a 90% prediction interval for future claims, courtesy of Chebyshev's inequality, *without assuming anything about the underlying distribution* (except finite mean and variance, of course).

You can get much tighter prediction intervals for future claims if you are willing to assume a parametric structure for the claims. If you assume, for example, that the claims are drawn from a Gamma distribution, then you can simply estimate the parameters of the Gamma distribution from your observations and compute a tighter 90% prediction interval than the relatively wide one provided by Chebyshev's inequality. The risk with this approach is, of course, that the underlying distribution may not be Gamma at all. The real distribution may be log-normal, say, or Pareto, both of which have heavier probability tails than the Gamma distribution, in which case your Gamma assumption would lead you to dangerously underestimate the sizes of future claims. Chebyshev's inequality gives you a much more reliable, albeit wider, prediction interval.

2.6k Views · View Upvoters

Your feedback is private.



Upvote · 7



Share



Add a comment...

Comment

Recommended All

Sponsored by Aha!

**What is a product roadmap?**

Build brilliant roadmaps in minutes. Trusted by over 200,000 users worldwide.

Start a free 30-day trial.

Free trial at aha.io



Ido Levin, studied Mathematics

Answered Jan 5, 2016



Thanks for the A2A (Although there is already a good old answer by [Sami Umut Can](#) ).

**Chebyshev's inequality** bounds the probability that a random variable will deviates from its expectation:

$\forall \alpha P(|X - \mu| \geq \alpha \sigma) \leq \frac{1}{\alpha^2}$  where  $\mu$  and  $\sigma$  are the expected value and standard deviation of  $X$ , respectively.

Now, lets say you take an exam and you know that the grades are normalized to  $\mu = 70$  and  $\sigma = 10$ . You want to know what's the probability you will fail (you assume that the grades are independent and that the distribution is symmetric, which is a rough assumption). Then

$$P_{\text{fail}} = P(X \leq 50) = 0.5 P(|X - \mu| \geq 2\sigma) < 0.125.$$

1.7k Views



Upvote



Share



Add a comment...

Comment

Recommended All



Luciano Robino, works at Centro Atomico Bariloche

Answered Oct 22, 2014



As [Sami Umut Can](#) stated. The idea of Chebyshev's inequality is to get as much information from you data when you have no clue how to describe a random variable that you have measured. It constitutes the minuium amount of information you can get from any random variable distribution

1.3k Views



Upvote



Share



Add a comment...

Comment

Recommended All

Find out now with a free home estimate online. Sell to Opendoor in days and skip the hassle of listing.

Get quote at [opendoor.com](https://opendoor.com)

...

Top Stories from Your Feed