

A
Position / length
Time from patch onset
0
0.773
100
1,000
13,011
B
Scaled active species
Position / length
0
0.773
30.8
251
1.3e+04
C
Orientation
Cue duration 0.773
Total establishment 116
Time from patch onset
D
Fixed width 3
Relative width 20%
Constant physical slope
Cue / reference peak
L = 15
L = 30
Position / length
E
Fixed width 3
Relative width 20%
Constant physical slope
Wave-pinning
Goryachev
deBelly
Holmes 3
Spring
+1.00 n=19
+1.00 n=16
+1.00 n=20
+1.00 n=20
+0.61 n=20
+0.63 n=20
+1.00 n=21
+1.00 n=21
+1.00 n=21
+1.00 n=21
+1.00 n=21
+1.00 n=21
+0.41 n=12
+0.29 n=12
+0.55 n=12
+0.48 n=12
+0.43 n=12
+0.38 n=12
+1.00 n=10
+1.00 n=7
+1.00 n=23
+1.00 n=11
+0.05 n=23
+0.41 n=23
+1.00 n=21
+1.00 n=21
+1.00 n=21
+1.00 n=21
-0.87 n=21
-0.87 n=21
Orientation
Strength
Spearman correlation

Figure 1. Enlargement changes the cue duration required for lasting polarity replacement

A, Recorded Wave-pinning preparation cue at length 15: a finite left patch, withdrawn at 0.773 native time units. White to black is zero to peak input; coloured vertical lines mark the profile samples in B.

B, Native active profiles from the same preparation, scaled using the common minimum and span of the displayed snapshots. Colours identify recorded times.


C, Orientation from the same preparation. Grey marks basal verification after establishment arrival at about 116; the cue has already ended. The matched time scale in A and C separates input duration from autonomous development.

D, Three rightward finite challenges restart the identical qualifying nominal-amplitude, relative-20% basal checkpoint at each model and length: fixed physical support 3 (Spring 0.2 native length units), relative support 20%, and a linear ramp with constant physical slope. Patch peaks remain fixed within each model; a fixed-slope ramp has a larger peak in a longer domain. Goryachev patches use the independently selected, length-independent multiplier K=64 (peak 6.4), which makes this a response-permissive comparison rather than a calibrated equal biochemical dose across models.

E, Spearman rank correlation between length and the shortest tested passing pulse duration, separately for orientation replacement and strength replacement, across Wave-pinning, Goryachev, corrected deBelly, Holmes 3 and Spring. Each cell reports the correlation and included successful-length count. All audited scalar successes enter, including those failing the separate native-field tail criterion. At least four distinct successful lengths and nonconstant durations are required; censored and preparation-ineligible cases receive no invented time. The companion duration curves retain all lengths and field-status markers.

After each positive pulse, input returns to the native basal value for 10,000 native time units. Orientation requires all 201 final-500 samples to have scalar orientation above 0.05 and contrast at least 0.05, with orientation range below 0.005. Strength additionally requires every tail orientation to be within 1% of the absolute initial leftward scalar orientation; it does not compare individual pole heights or amounts. The separate native-field tail flag requires pointwise range and projected 500-unit drift within 0.1% of frozen field scales. The reported duration excludes preparation and recovery. It is the shortest passing tested duration, with sampled fail/pass boundaries refined to 1%; narrower unobserved intervals are not excluded.

deBelly uses corrected endpoint commands. Spring retains native length and time units. Cross-model amplitudes and clocks are not calibrated biochemical doses or cellular times.


Goryachev
Wave-pinning
Holmes
deBelly
Spring
A
Amount / reference
Length / reference
B
Extent / reference
Length / reference
C
Peak / reference
Length / reference
D
Goryachev
Wave-pinning
Holmes
deBelly
Spring
Initiation / L19
Native time
Length L
E
w = 0.95
w = 1.9
w = 3.8
w = 7.6
w = 15.2
Area 0.5×
Area 1×
Area 2×
Replacement time (native units)
Length L
F
Pulse-2 gain / headroom
Gap-end / pulse 1
Open circles: below activity gate at x = 0. Size increases from light (L19) to dark (L39).

G
Goryachev
lost
Wave-pinning
kept
Holmes
kept
deBelly
lost then commits
Spring
lost then commits
Challenger share
Time / pulse
L39
H
WGD−
WGD+
Initiation-limited
Goryachev
Saturating activation
Wave-pinning / Holmes
Slow non-activity state
deBelly / Spring
Biosensor
(schematic)
Time (schematic units)
I
Replacement time
(schematic)
narrow
broad
Width at fixed dose

Figure 2. Replacement resistance in enlarged cells has distinct size and pulse-memory signatures

Enlarged cells resist replacement through distinct initiation, cue-width and pulse-memory signatures.

A, Incumbent excess amount normalized to the model reference, across the same 97 retained states.

B, Incumbent extent normalized to the model reference, for those same states.

C, Incumbent peak normalized to the model reference, for those same states.

D, Initiation is the time from cue onset until maximum in-cue activity minus the profile minimum exceeds 10% of the model's L19 incumbent excess peak. Area 1, width 3.8; times are normalized to L19. All six deBelly upper bounds are 0.5 native units (downward arrows); first-update sampling leaves its size dependence unresolved.

E, Replacement time (first eligible challenger share ≥ 0.75) versus length, with a line per width and rows for areas 0.5, 1 and 2. Upward triangles/arrows are lower bounds: the cue never replaced within the horizon (6,000 Wave-pinning; 10,000 Goryachev/Holmes; 2,000 deBelly; 100 Spring, native units). Broad cues rescue Holmes across all lengths; narrow-cue censoring does not establish a growing penalty. Spring native widths equal the displayed widths divided by 38.

F, Gap-1 activity kept versus pulse-2 gain as a fraction of headroom (Eq. 1). Four equal pulses, total ON 1.5 times the model-length commitment calibration; gaps 0–16 pulse durations. Holmes L31–39 uses the L27 calibration. Other models are grey; lightness indicates L. Open circles at x = 0 code activity below the existing visibility gate. At the left wall, deBelly/Spring gain on pulse 2 despite no visible activity; Goryachev never does. The deBelly L19/gap16 first-gap endpoint was checked against native fields.

G, L39 gap4 challenger-share histories, labelled kept / lost / lost then commits; shading marks cue ON. deBelly includes full recovery; others show one post-train pulse duration. Continuous raw shares include below-gate values. deBelly's later timing and fields remain numerically sensitive (supplement).

H, Prospective biosensor histories for the same pulse train in WGD− and enlarged WGD+ cells. Initiation-limited resetting (Goryachev), saturating activation with gap retention (Wave-pinning/Holmes), and progress retained after activity loss (deBelly/Spring) give distinct dimensionless schematic histories.

I, Prospective equal-dose width responses for those classes: weak width rescue, strong width rescue, and weaker width rescue. Solid grey: WGD−; dashed black: WGD+. These are qualitative assay predictions.

$$x = \frac{s_{\mathrm{gap}\,1}}{s_1}, \qquad y = \frac{s_2 - s_1}{1 - s_1} \tag{1}$$


s_1 and s_2 are challenger shares at pulse ends 1 and 2; s_gap1 is the first-gap-end share. An observed endpoint below the existing visibility gate is coded as zero visible share in F; its raw

Figure 2. Replacement resistance in enlarged cells has distinct size and pulse-memory signatures · continued


value remains in the CSV.

A

Grey: cue support; diamonds: saddle; | heading




B



B: light → dark = C 30, 37.8, 47.6; Spring 2.5, 3.15, 3.97.
Open triangles: time lower bounds. Dotted lines: first saddle. Grey: heading undefined.

C


Lines: native map; circles: new pulse tests; open squares: existing edge pulse. Dotted verticals: half-contrast edges.




D

E

F



G



H




Figure 3. Enlargement slows both turning routes and selectively suppresses edge susceptibility

A. Wave-pinning activity profiles for a cue straddling the incumbent edge (i) or separated from it (ii), C=30. Rows show the initial state, half-turn, 90% turn and cue-free relaxation. Grey marks cue support, diamonds mark saddles and the black tick marks activity heading.

B. Enlargement slows half-turning through both incumbent sliding and replacement by a new region. Half-turn time is plotted against cue-edge gap for C=30, 37.8, 47.6 (Spring 2.5, 3.15, 3.97), light to dark. Vertical lines mark the first saddle at half-turn; grey marks undefined heading at one or more sizes. Open upward triangles with arrows denote time lower bounds.

C. Signed terminal displacement per dose for a Gaussian pulse centred at each arc position, at f=0.001: SD 1.5, duration 50 (Spring 0.15, 5). Light and dark are C20/C60 (Spring 1/3). Lines are the native linearized response to this pulse shape; circles are direct pulse tests and open squares mark the edge test. The edge pulse confirms the map locally; additional tests cover the domain centre, halfway to the edge, edge ±1 and ±3 pulse SD, quarter-basal arc and antipode. Maximum absolute gain error is 1.69% of map amplitude. Largest/smallest terminal edge-gain folds are Wave-pinning 0.701, Holmes 0.914, Spring 0.989, Goryachev 0.088. Enlargement suppresses susceptibility strongly in Wave-pinning and Goryachev, while Holmes and Spring change little. The spatial response supplies the drift prediction in G.

D. Mean-state activity on the same centred physical arc coordinate as C and E. Dotted lines locate half-contrast edges. Spring uses material positions multiplied by mean circumference.

E. Nearly all predicted drift (99.6–100%) is generated within two input correlation lengths of the two edges, suggesting lateral-edge fluctuations as the cellular readout. Curves show signed per-position contributions divided by total predicted diffusion; negative values remain. Rounded shares per edge/plateau interior/basal arc (%) are: Wave-pinning C20: 50.0/<0.1/<0.1; Wave-pinning C60: 49.9/0.1/<0.1; Holmes C20: 50.0/<0.1/<0.1; Holmes C60: 50.0/<0.1/<0.1; Spring C1: 50.0/0/0; Spring C3: 49.8/0.2/0.2; Goryachev C20: 50.0/0/<0.1; Goryachev C60: 50.0/0/<0.1. Interior and basal zones exclude edge zones; exact signed fractions are supplied with the source data.

F. Wave-pinning C20 and C60: the pulsed edge (solid) and opposite edge (dashed) move together and retain a permanent shift after withdrawal. Grey marks pulse-on; displacement is divided by pulse dose. Other time courses and integrated responses are supplementary.

G. A weak input anywhere shifts the domain by g(x) times dose; noise is many such inputs; drift follows from g and the input covariance; nothing is fitted. Shared-translation predicted versus measured diffusion is divided by each quantity's own smallest-size value at the same input (domain drift is also shown in Figure 4G). Grey is equality. Colours encode size; shapes encode f=0.0001, 0.0003, 0.001, 0.003, 0.005, 0.01 (circle, square, upward triangle, downward triangle, diamond, filled plus). Horizontal intervals are measured 95% bootstrap intervals. Wave-pinning's terminal gain fold gives $0.70^2 \approx 0.49$; the full spatial covariance predicts 0.490, versus 0.473 measured at f=0.001. The square relation is approximate because spatial shape and covariance also change. Goryachev fails quantitatively and retains its own scale. Absolute diffusion and the independent-edge candidate are supplementary.

Figure 3. Enlargement slows both turning routes and selectively suppresses edge susceptibility · continued

H. The drift prediction implies a local size exponent, compared with measured heading-persistence α from Figure 4E. Predictions use the exact mean input of every observation; no f=0.001 extrapolation is used. The drift subset requires f≤0.01 and multiple-domain fraction≤0.05 throughout the fixed size neighbourhood. Native stationary-state and pulse-linearity gates must also pass. Grey is equality; vertical intervals are observed 95% bootstrap intervals; colours and shapes match G. 53 qualified pairs include all 15 Wave-pinning observations at f=0.003, 0.005 and 0.01. Its predicted α spans 2.48–2.88, above the size-independent-drift expectation of 2. Within Wave-pinning, this comparison tests the excess α above 2 more than the within-model trend. Goryachev has no qualified pair. Every excluded condition and reason remains in the source data.

$$G(x_0) = \sum_j g_j h(x_j - x_0) \tag{3.1}$$

The main map is the standard Gaussian-pulse response per dose, not raw cell-input g.


$$c_j = g_j \sum_k g_k \int_0^\infty \overline{K_u(j,k;t,t+v)}^{\,t}\, \mathrm{d}v, \qquad D_s = \sum_j c_j \tag{3.2}$$

Signed per-position contributions retain the exact native input covariance.


$$\alpha_{\mathrm{pred}} = 2 - \frac{\mathrm{d}\ln D_s}{\mathrm{d}\ln \bar{C}} \tag{3.3}$$

Same fixed adjacent-size neighborhoods as Figure 4E; exact matched-input predictions and explicit drift-regime eligibility.


A


Drifting peak (same physical D_s)
Jumping peak (same rate)
Jumping peak (rate ∝ perimeter)
Time (×10³)
C = 20
C = 60


B


Drifting peak (same physical D_s)
Jumping peak (same rate)
Jumping peak (rate ∝ perimeter)
$\alpha = 2$
$\alpha = 0$
$\alpha = -1$
Heading $C(\ell)$
Lag (×10³)
C = 20
C = 60


C


Wave-pinning, f = 0.03
Holmes, f = 0.02
Spring, f = 0.005
Goryachev, f = 0.01
Time (×10³)
C = 20
C = 60
C = 1
C = 3
0–9.2
0–83


D


Wave-pinning
Holmes
Spring
Goryachev
Heading $C(\ell)$
Lag (×10³)
C = 20
C = 60
C = 1
C = 3


E


Wave-pinning
Holmes
Spring
Goryachev
C: light → dark
Local exponent α
Input fraction f


F


Wave-pinning
Holmes
Spring
Goryachev
$f_{1/2}/f_{1/2}(C_{20})$
C (Spring ×20)


G


Wave-pinning
Holmes
Spring
Goryachev
$D_s/D_s(C_{min})$
Input fraction f

H


Wave-pinning
Holmes
Spring
Goryachev
k_{pred}/k_{obs}
Input fraction f

I


Wave-pinning
Holmes
Spring
Goryachev
Remainder share R
Input fraction f

J



Wave-pinning, f = 0.05
Holmes, f = 0.05
Spring, f = 0.05
Goryachev, f = 0.025
0 < d ≤ C/6
d > C/6
Takeover probability (20τ)
Circumference C

Figure 4. Enlargement extends polarity heading memory under fluctuating input

Model panels use positive lognormal input with mean f times the native reference: Wave-pinning 0.1, Holmes 1, Spring 4 and Goryachev 6.4. The Gaussian driving field has unit variance, lognormal amplitude σ = 1, spatial parameter 1.5 and OU correlation time 50 (Spring basal-physical 0.15 and 5). Spring input is stationary in material coordinate, so its physical correlation length scales with circumference. Native clocks are not calibrated between models. Unless stated otherwise, error bars are 95% whole-history bootstrap intervals.

A, Detector-free null kymographs at C = 20 and 60: identical physical diffusion, size-independent jumping, or perimeter-proportional jumping. Each facet shows the same activity measure.

B, Null heading autocorrelation; persistence size exponents are 2, 0 and −1. C = 20 is light and C = 60 dark.

C, Native-model kymographs at f = 0.03 (Wave-pinning), 0.02 (Holmes), 0.005 (Spring) and 0.01 (Goryachev). Arc position is centred at the initial heading; time is elapsed after burn-in. Goryachev activity ranges are labelled separately for each size.

D, Model heading autocorrelation at the inputs in C, for the smallest and largest rings (Spring C = 1 and 3). Solid curves show the smallest ring and dotted curves the largest. Endpoint eligibility requires polarity coherence $|P| \geq 0.1$. These direct curves show longer heading memory in larger rings.

E, Local persistence exponent α versus input fraction f, using three neighbouring sizes (two at endpoints), with whole-history 95% intervals. In E and G–I, sizes are C = 20, 30, 40, 50 and 60 (Spring 1, 1.5, 2, 2.5 and 3), light to dark. Open circles in D, E, G and I mark the declared combined forcing-control flags; unavailable rates are omitted.

F, Noise-tolerance fold at heading-correlation lag 2000 (Spring 200), relative to the smallest ring. Spring circumference is multiplied by 20 for display. The dashed grey guide is proportional to circumference.

G, Physical within-tenure drift coefficient normalized by the smallest size at the same input, using five-frame persistent dominant tenures. Goryachev uses a different ordinate range, 10^{-7}–2; the other models use 0.2–8.

H, Drift plus counted five-frame-dominant takeovers, divided by observed heading decorrelation rate, at every designed input. Solid lower and dashed upper limits bound every unscored takeover loss between 0 and 2. Shading denotes identification intervals, not confidence intervals. Grey line: unity. Open upward triangles with arrows mark an unbounded upper limit where drift is unavailable. Only the unknown drift contribution has lower bound zero; counted-takeover lower contributions are retained. Crosses at the top mark unavailable observed rates.

I, Signed remainder share versus input, using point regions with native-amplitude scaling. R includes amplitude, within-region shape, unresolved activity and background contributions; its sign and magnitude depend on the accounting convention. This convention retains measured native coherence, avoids the unit-modulus single-region normalization artefact and gives a common comparison across inputs and sizes. The decomposition is descriptive. Exact own-mask

Figure 4. Enlargement extends polarity heading memory under fluctuating input · continued

and normalization sensitivities appear in S4.5; complete-mask reconstruction is an algebraic identity, not causal validation.

J, Probability that a certified persistent secondary birth becomes globally dominant within 20 input correlation times, at each model's labelled input. Filled circles: 0 < d ≤ C/6; open squares: d > C/6, where d is estimated edge-to-edge birth–incumbent separation. Probabilities concern the certified positive-distance subset; a [0,1] interval is uninformative. Lower finite-horizon incidence can reflect slower or rarer takeover; lifetime takeover probability is unidentified. Input selection, certification denominators and interval procedures are in Methods.

Scope/limitations: The strong-input regime remains unexplained.

$$Q_t = \frac{\sum_{i \in I_t} A_{i,t} e^{i\theta_{i,t}}}{\sum_{i \in I_t} A_{i,t}}, \qquad r_t = |P_t|, \quad \widetilde{Q}_t = r_t Q_t \tag{4.1}$$

Q is the normalized point-region moment; P=M₁/M₀ is measured from the native field. Its amplitude r prevents the unit-modulus single-region normalization artefact.

$$\Delta P_j = \bar{r}\,\Delta Q_j \ (j = \mathrm{M}, \mathrm{A}, \mathrm{B}), \qquad \Delta P_{\mathrm{R}} = \Delta P - \bar{r}\,\Delta Q \tag{4.2}$$

The three ΔQ components use the symmetric Shapley convention for motion, coexisting amount and membership; r̄ is mean endpoint native amplitude. R closes the native moment change and includes amplitude variation, shape and background. This is an accounting convention, not causal attribution.

$$L_j = \frac{\mathrm{Re}\left[\Delta P_j \overline{(H_1 - H_0)}\right]}{|P_0| + |P_1|}, \quad H_t = \frac{P_t}{|P_t|}, \quad \sum_j L_j = 1 - \cos(\theta_1 - \theta_0) \tag{4.3}$$

Pooled signed fractions are sums of L_j divided by total measured heading loss. They retain cancellation, can exceed one, and depend on the stated accounting convention; they are not causal shares.

$$k(\ell) = -\frac{\ln C(\ell)}{\ell}, \qquad k \propto \bar{C}^{-\alpha} \tag{4.4}$$

C(ℓ) is activity-heading autocorrelation; mean C is actual circumference. Endpoint eligibility uses |P|≥0.1. Main H evaluates every designed input; unavailable observed rates remain explicit.

$$a_n(t) = \frac{1}{N} \sum_{j=1}^{N} u_j(t) e^{in\theta_j}, \qquad P(t) = \frac{a_1(t)}{a_0(t)}, \qquad q_n = \frac{2\pi n}{\bar{C}} \tag{4.5}$$

Figure 4. Enlargement extends polarity heading memory under fluctuating input · continued

Fourier modes use cell-centre material phases and the positive convention matching the native first moment. Heading is the phase of P; Spring physical wavenumber uses mean circumference.

$$\gamma_n = -\frac{\ln C_n(\ell_*)}{\ell_*}, \qquad k(C) = \gamma(2\pi/C), \qquad \alpha = \frac{\mathrm{d}\ln\gamma}{\mathrm{d}\ln q} \tag{4.6}$$

The last two equalities are the detector-free hypothesis under a common size-independent mode curve, not an identity assumed in analysis. Complex correlations use the nearest-half lag rule. Half-crossing status is exported; primary collapse and slope comparisons use observed crossings. The declared descriptive screens and stencil sensitivities are reported model by model.

$$k^{\mathrm{lo}}_{\mathrm{pred}} = q_1^2 D_s + \frac{\sum_{\mathrm{scored}} (1-\cos\Delta\theta)}{T}, \qquad k^{\mathrm{hi}}_{\mathrm{pred}} = k^{\mathrm{lo}}_{\mathrm{pred}} + \frac{2N_{\mathrm{unscored}}}{T} \tag{4.7}$$

All counted takeovers enter the interval: each unscored angular loss lies between 0 and 2. T is eligible exposure. Missing drift has a nonnegative lower bound and unbounded upper limit; an unavailable observed heading rate prevents a numerical ratio.