

A
Preparation cue
Position / length
Time from patch onset
Native active profiles
Scaled active species
0
0.773
30.8
251
1.3e+04
Measured polarity
Orientation
Cue duration 0.773
Total establishment 116
B
Fixed width 3
L=15
L=30
Cue / reference peak
Position x / L
Relative width 20%
Constant physical slope
C
Fixed width 3
Relative width 20%
Constant physical slope
Wave-pinning
Goryachev
deBelly
Holmes 3
Spring
+1.00 n=19
+1.00 n=16
+1.00 n=20
+1.00 n=20
+0.61 n=20
+0.63 n=20
+1.00 n=21
+1.00 n=21
+1.00 n=21
+1.00 n=21
+1.00 n=21
+1.00 n=21
+0.41 n=12
+0.29 n=12
+0.55 n=12
+0.48 n=12
+0.43 n=12
+0.38 n=12
+1.00 n=10
+1.00 n=7
+1.00 n=23
+1.00 n=11
+0.05 n=23
+0.41 n=23
+1.00 n=21
+1.00 n=21
+1.00 n=21
+1.00 n=21
-0.87 n=21
-0.87 n=21
Orientation
Strength
Spearman correlation

Supplement 1 · Length versus finite cue duration

Five models × three tests · exact shared relative-20% basal checkpoint · unchanged scalar gates and recovery

deBelly uses fresh corrected preparations and independent endpoint commands

Orientation gate
Polarity strength within 1%
No passing pulse within tested range
Ineligible preparation (below axis)
Scalar success; native-field flag fails

Fixed width 3
Spring: native width 0.2

Relative width 20%

Constant physical slope

Goryachev

deBelly

Holmes 3

Spring

Shortest passing tested cue duration

Nominal cell length (native units)

Native resting length L0

1 ineligible

2 ineligible

10 ineligible

0 ineligible

Both patches use a constant peak across all lengths; Goryachev K=64, peak=6.4. Gradients retain one constant physical slope per model.
Each positive duration restarts the original checkpoint. Full strength adds the 1% reversed initial-polarity band to the orientation gate.
Open markers retain failing field-tail flags. Censoring marks explicit tested limits; missing endpoints break lines. Native times and scales differ.
Data generated 2026-10-08T19:26:33.200718+00:00; producer finished. Every imported and fresh trial passed independent native-array audit.