Manuscript outline

Figure 2 — round 3

Size responses and pulse-memory signatures

22 October 2026 · Figure 2 round 3 integrated; other main figures retain completed round-2 content

Scientific argument and manuscript plan

Proposed main figures

- Figure 2. Replacement resistance in enlarged cells has distinct size and pulse-memory signatures



Goryachev
Wave-pinning
Holmes
deBelly
Spring
A
Amount / reference
Length / reference
Extent / reference
Peak / reference
B
Initiation / L19
Native time
Length L
C
w = 0.95
w = 1.9
w = 3.8
w = 7.6
w = 15.2
> horizon
Area 0.5×
Area 1×
Area 2×
Replacement time (native units)
D
Gap 1 retained share
Below activity gate: open circle at x = 0
L19
L39
L19, gap 16
Pulse-2 gain / headroom
Gap-end / pulse 1
E
lost
kept
lost then commits
Challenger share
Time / pulse
F
Predictions (schematic)
WGD−
WGD+
Initiation-limited
Saturating activation
Slow non-activity state
Biosensor
Replace time
narrow
broad
Width at fixed dose

Figure 2. Replacement resistance in enlarged cells has distinct size and pulse-memory signatures

Separate size-dependent initiation and cue-width responses from observable pulse-memory classes.

Detailed legend

Enlarged cells resist replacement through distinct initiation, cue-width and pulse-memory signatures.

A, Incumbent excess amount, extent and peak, normalized to each model's reference, across 97 retained states.

B, Initiation is the time from cue onset until maximum in-cue activity minus the profile minimum exceeds 10% of the model's L19 incumbent excess peak. Area 1, width 3.8; times are normalized to L19. All six deBelly upper bounds are 0.5 native units (downward arrows); first-update sampling leaves its size dependence unresolved.

C, Replacement time (first eligible challenger share ≥ 0.75) versus length, with a line per width and rows for areas 0.5, 1 and 2. Upward triangles/arrows are lower bounds: the cue never replaced within the horizon (6,000 Wave-pinning; 10,000 Goryachev/Holmes; 2,000 deBelly; 100 Spring, native units). Broad cues rescue Holmes across all lengths; narrow-cue censoring does not establish a growing penalty. Spring native widths equal the displayed widths divided by 38.

D, Gap-1 activity kept versus pulse-2 gain as a fraction of headroom (Eq. 1). Four equal pulses, total ON 1.5 times the model-length commitment calibration; gaps 0–16 pulse durations. Holmes L31–39 uses the L27 calibration. Other models are grey; lightness indicates L. Open circles at x = 0 code activity below the existing visibility gate. At the left wall, deBelly/Spring gain on pulse 2 despite no visible activity; Goryachev never does. The deBelly L19/gap16 first-gap endpoint was checked against native fields.

E, L39 gap4 challenger-share histories, labelled kept / lost / lost then commits; shading marks cue ON. deBelly includes full recovery; others show one post-train pulse duration. Continuous raw shares include below-gate values. deBelly's later timing and fields remain numerically sensitive (supplement).

F, Schematic biosensor predictions for the same pulse train and equal-dose width series in WGD− and enlarged WGD+ cells: delayed rises and resets; saturation with gap retention and strong width rescue; or gap collapse followed by commitment, with weaker width rescue. Predictions are prospective. Model kinetics are unchanged.

Narrative points and conclusions

- Enlarged cells resist replacement through distinct initiation, cue-width and pulse-memory signatures.

Figure 2. Replacement resistance in enlarged cells has distinct size and pulse-memory signatures · continued

- The full width series shows that Holmes can replace at every tested length when the cue is broad enough; narrow-cue censoring cannot establish a growing width-penalty ratio.
- Pulse-2 headroom gain separates low visible retention with residual progress from the initiation-limited reset response. Saturating activation retains visible activity through gaps.

Proposed measurements and experimental follow-up

Measure first in-cue biosensor rise, then vary width at fixed integrated dose.

Measure pulse-1, gap-1 and pulse-2 challenger shares under the same pulse train, followed by eventual commitment.

Scope, limitations and pending validation

Commanded input areas and native clocks are not calibrated across models.

deBelly initiation is unresolved within the first native update; later-gap timing and endpoint fields remain numerically sensitive.

D codes known below-gate activity at x=0; the exact raw ratio remains in the plotted-value CSV.

Holmes large-length trains have no target-length commitment calibration; visible retention differs from eventual replacement.

The three classes describe selected deterministic regimes; WGD biosensor predictions are prospective.

Model definitions and readouts

$$x = \frac{s_{\mathrm{gap\,1}}}{s_1}, \qquad y = \frac{s_2 - s_1}{1 - s_1} \tag{1}$$

s_1 and s_2 are challenger shares at pulse ends 1 and 2; s_gap1 is the first-gap-end share. An observed endpoint below the existing visibility gate is coded as zero visible share in D; its raw value remains in the CSV.

Sources and reproducibility

All 450 per-width replacement times

All 180 gap1/pulse2 coordinates and raw shares

deBelly L19/gap16 native-phase audit

Preserved prior main panels and complete supplement

Supplementary material and supporting analyses

Complete supplementary evidence

Superseded main panels and all previous supplements are preserved.

Full Figure 2 supplement