A

Drifting peak (same physical D_s) | Jumping peak (same rate) | Jumping peak (rate ∝ perimeter)

Time (×10³) — C20 — C60

Heading $C(\ell)$ — Lag (×10³)

$\alpha = 2$ (C = 20, C = 60) | $\alpha = 0$ (curves coincide) | $\alpha = -1$

B

C: 20 30 40 50 60

Drift only | Drift + same-rate jumps | Drift + perimeter-rate jumps

Local exponent α — Input fraction f

drift | same-rate jumps | perimeter-rate jumps

$f_{1/2}/f_{1/2}(C_{20})$ — Circumference C

Drift | + same-rate jumps | + perimeter-rate jumps

$\propto C$

C

Wave-pinning: 20 30 40 50 60 | Holmes: 20 30 40 50 60 | Spring: 1 1.5 2 2.5 3 | Goryachev: 20 30 40 50 60

C; Spring ×20 in fold panel

Wave-pinning | Holmes | Spring | Goryachev

Local exponent α — Input fraction f

$f_{1/2}/f_{1/2}(C_{20})$ — Circumference C

Wave-pinning | Holmes | Spring | Goryachev

$\propto C$

D

Wave-pinning, $f = 0.03$ | Holmes, $f = 0.02$ | Spring, $f = 0.005$ | Goryachev, $f = 0.01$

Time (×10³) — switch — C20 — C60 (C1 — C3 for Spring)

Heading $C(\ell)$ — Lag (×10³)

$\alpha_{20,60} = 2.1$, 95% CI [1.9, 2.3] (C = 20, C = 60)

$\alpha_{20,60} = 1.4$, 95% CI [1.3, 1.5]

$\alpha_{20,60} = 1.8$, 95% CI [1.7, 1.9]

$\alpha_{20,60}$ unavailable

E

Wave-pinning | Holmes | Spring | Goryachev

$D_s/D_s(C_{\min})$

λ_{eff} — Scored events — Mixture fit

$k_{\text{pred}}/k_{\text{obs}}$

Multiple-front fraction

Input fraction f