Figure 3 round 2 · decomposition and censoring

At matched edge gap, larger native size accounts for most of the increase in driven handover time. Extra accessible separation contributes relatively little. Cue-free competition has a stronger separation response and remains unresolved in several mass-conserved-model conditions.

For each model and assay, the matched gap is the largest gap within both measured size curves. In this dataset it is the small-size antipodal-placement endpoint. In B, the coordinate is the measured gap at withdrawal, not the initial cue gap. The endpoint called antipode is the trajectory seeded by the initial antipodal cue: its withdrawal region centers can shift.

A1 uses native continuous complete-region handover roots. B uses saved-sample lower and upper time brackets. Linear interpolation is restricted to adjacent measured eligible placements. Interval endpoints are interpolated separately; any censored upper endpoint stays infinite. Missing or topology-ineligible values are never converted into bounds.

A ratio with a censored numerator and resolved denominator has a lower bound. A resolved numerator and censored denominator gives an upper bound. Two censored times give no finite one-sided ratio bound; the result is marked indeterminate. In particular, dividing two horizon caps is invalid. Finite bars use the geometric center of the resulting interval, with the full interval recorded in CSV.

The exact product identity uses the same interpolated matched-gap time in both terms. For finite B intervals the representative values obey the identity; multiplying independently propagated marginal interval limits would ignore this shared denominator and numerator. Total intervals are computed directly from endpoint times.

Optional extensions awaiting user review: 9 Holmes ratio-1 histories (5 small, 4 middle), each completed to 20,000 and eligible for a proposed 80,000 horizon. They lie within the common withdrawal-gap range. No extension has been submitted. These extensions may still leave large-size matched-gap or antipodal times censored.

The following pages retain BOTH superseded main pages, including the complete translation page, then the entire seven-page existing supplement. All four native-profile appendix volumes remain linked in the existing reader; no model inputs, histories, or prior supplements are replaced.

All decomposition values · time ratios with interval bounds

Model	Assay	Size / small	Matched gap	Size term	Extra separation	Total
Wave-pinning	A1	37.8 / 30	7.719	2.14	1.01	2.17
Wave-pinning	A1	47.6 / 30	7.719	5.59	1.05	5.86
Goryachev	A1	37.8 / 30	12.05	1.84	1.04	1.91
Goryachev	A1	47.6 / 30	12.05	3.5	1.15	4.02
Holmes	A1	37.8 / 30	6.29	3.59	1	3.6
Holmes	A1	47.6 / 30	6.29	23.9	1.01	24.2
Spring	A1	3.15 / 2.5	0.4296	2.44	1	2.44
Spring	A1	3.97 / 2.5	0.4296	8.45	1.01	8.52
Wave-pinning	B	37.8 / 30	8.378	5.67–5.77	1.02–1.03	5.81–5.92
Wave-pinning	B	47.6 / 30	8.378	≥15.9	indeterminate (both censored)	≥15.9
Goryachev	B	37.8 / 30	11.87	1.15–1.16	1.01–1.02	1.16–1.18
Goryachev	B	47.6 / 30	11.87	1.44–1.47	1.06–1.09	1.54–1.58
Holmes	B	37.8 / 30	7.609	indeterminate (both ce	indeterminate (both censored)	indeterminate (both censored)
Holmes	B	47.6 / 30	7.609	indeterminate (both ce	indeterminate (both censored)	indeterminate (both censored)
Spring	B	3.15 / 2.5	0.5456	4.2–4.38	1.73–1.77	7.31–7.71
Spring	B	3.97 / 2.5	0.5456	≥6.87	indeterminate (both censored)	≥13.8

Intervals reflect saved-time brackets, not biological uncertainty. ≥ is a one-sided lower bound.
All interval endpoints, interpolation weights and source slots appear in the companion CSVs.