
Excess active amount
Core width
Peak concentration
A Incumbent organization
Wave-pinning 19→39
Goryachev 19→39
deBelly 19→39
Holmes 3 20→40
Spring L₀ 1→2
Larger / smaller value
Wave-pinning
Goryachev
deBelly
Holmes 3
Spring
B Initial opposite-end input gain: all 97 basals
Refs: L19; Holmes L20; Spring resting L₀ = 1
Relative nominal size is not a physical calibration.
Cue-weighted gain / own reference (log₂ scale)
Nominal size / own reference size
C Restored local deficit over the model-specific early window
Exactly three measured pilot sizes per model; common deactivation rule, different fixed native clocks.
Wave-pinning τ = 0.1778 native
Goryachev τ = 0.005623 native
deBelly τ = 5 native
Holmes 3 τ = 0.1778 native
Spring τ = 0.01778 native
Initial deficit restored (fraction)
Native pilot sizes (Spring: resting L₀). Points are unjoined; compare size dependence within each model.
Bars show empirical numerical sensitivity (3E), not biological confidence intervals.
D Native basal profiles: the same small/large pairs as A
Smaller
Larger
Wave-pinning
Goryachev
deBelly
Holmes 3
Spring
Active concentration (native)
Physical interval coordinate (native units); profile concentration scales differ across models.
All panels use linear 1D states. Spring profiles use measured physical domains 1.049→2.049.