
Figure 2 · Ring correspondence and local response

Continued review sheet 1 of 2 · retained measurements plus an explicit recovery-map gap

A Ring / interval observable	Expected	Wave-pinning	Goryachev	Holmes 3
Full half-contrast width	2	1.9999–2.0000	1.9979–2.0046	2.0000–2.0000
Full excess-active core amount	2	1.9999–2.0001	1.9959–2.0030	1.9999–2.0000
Absolute primary peak	1	1.0000–1.0000	1.0009–1.0020	1.0000–1.0000
Matched cue-weighted input gain	1	1.0000–1.0000	0.9995–1.0024	1.0000–1.0000

13 independently prepared pairs at C = 2L: width/amount double; peak and matched-cue gain agree.
Baseline input matching: endpoint support 3 ↔ full ring support 6. The retained finite challenges below use full support 3.



One fixed 0–3 scale: within-size gain varies by <0.09% across eligible placements. Numbered cells link to sheet 2.
Rows are discrete sampled circumferences; × = 198 cue/core overlaps; dark gray = 17 unavailable Holmes C80 basals.
Native input/time units are not calibrated across models.

C Common active-deactivation recovery baseline and separation maps: not measured yet

This checkpoint contains the 15-case linear clock pilot only. No full linear or ring recovery grid is inferred.
The basal ring coefficient has no future separation; a separation map needs the same kick on declared challenged states.