Shared boundary translation and the explicit size comparison

The deterministic data, measured drift estimates, pulse-gain calibration and exact forcing covariance are unchanged. Filled main-panel D markers now retain the shared neutral translation; open markers preserve the tested independent-edge candidate. No correction is fitted to measured drift. Near-unity late-lag boundary co-motion refutes independence in Wave-pinning, Holmes and Spring.

The second D row compares predicted and measured size ratios at each fixed input. The same model-specific smallest size is used in both denominators. This removes the dominant f-squared scaling from the display. In Wave-pinning at f=0.001, the C60/C20 ratio is 0.4939 predicted and 0.4732 measured; the measured terminal susceptibility ratio is 0.7007. This is a concrete size test, not an inference from the absolute five-decade span.

C insets show finite integrated displacement divided by the smallest-size value, on identical zero-based 0–1.1 axes. The terminal gain falls strongly in Goryachev and moderately in Wave-pinning, while Holmes and Spring change much less. The two perturbation assays are distinct: varying-offset continuous photoactivation tests sliding versus takeover and half-turn timing; a weak brief boundary pulse tests small-signal boundary displacement and common translation. The edge assay does not uniquely identify the Figure 2 mechanism classes.

Goryachev C20 and C30 at f=0.01 lack a stationary polarized boundary at the mean input. Their absolute predictions are undefined. All five Goryachev size-normalized predictions at that f are undefined because the fixed C20 reference is undefined; a larger size is not substituted. Goryachev disagreement remains a result, and physical tracked domain drift does not explain its entire heading decay.

The historical reviewed main page and complete 25-page reviewed supplement follow these current supplementary pages. They remain dated records of the independent-edge presentation at commit 49754ab. The current main and legend supersede their interpretation, while all numerical source values and scientific limitations are retained.

Absolute agreement and size dependence; no fitted correction



Bottom: solid prediction, dashed observation, one curve per input. Source tables retain condition-specific values, missing references and exposure.