
Larger incumbents: more material, organized differently

Prior all-five evidence retained unchanged: excess amount, physical extent and native peak concentration.



Replacing an incumbent requires more than activating a new region

A wider cue can recruit more distal activity and weaken the old region more, yet leave a challenger that fails after withdrawal.


The next sheet shows this directly in Goryachev L9: same basal, same requested peak, same pulse; only physical width changes.

This illustrates coupled competition. Larger-cell and WP examples provide context; it is not a universal width rule or a unique molecular signature.


The organization comparison is observational. The challenge changes both spatial support and integrated input; it does not isolate incumbent height or width causally.


Greater incumbent weakening can accompany challenger failure


Goryachev L9: exact shared basal, requested native peak 6.4, pulse 74, widths 1.8 / 3; then 10000 cue-free units.



Actual saved fields; time is linear through 74 and logarithmic later. Cyan marks the physical cue-support edge during stimulation.







Native mechanism and hypothesis

$$J = (ae_c u^2 + be_c u + S)v - cu$$

Stimulation recruits from inactive v. Withdrawal removes S; nonlinear local maintenance and transport continue. Distributed activation may fail to sustain a concentrated challenger in this case.

Supported: the broader response forms with more distal active amount, then disappears. The shared-pool/local-maintenance account is a hypothesis, not a unique causal attribution.

Larger Goryachev and WP width contrasts differ; this is not a universal peak or depletion threshold. Native clocks/input are uncalibrated; N = 201; no spatial-convergence claim.