

Figure 2 discussion: larger incumbents and cue trade-offs

Prior organization result + existing feasibility evidence + proposed campaign · author-review draft · 19 October 2026



E–G above are prior work, reused unchanged. The cue experiment below is a proposal for discussion.

PROPOSED CAMPAIGN: vary peak, support and pulse duration independently



Regional movies across this grid have not been collected. Pulse duration T and cue-on event latency are different measurements.


For author input: optical actuator/native-channel mapping, cue levels, weakening criterion and retention criterion.

Existing two-width comparisons motivate a targeted cue campaign

Same exact basal and requested peak within each model/length · right end-to-zero raised-cosine patches · no new simulations

Native requested peaks: WP 0.1; Goryachev 6.4. These are not matched optical doses.

Selected qualified anchors



Point = passing-pulse ratio; whiskers use both tested brackets.
Bracket envelope, not CI. Exact construction is in the caption.

Actual tested native pulse brackets

Model; size	Support	Narrow (F, P]	Broad (F, P]
WP; L9	1.8 → 3	(19.75, 19.875]	(12.125, 12.1875]
WP; L19	3 → 3.8	(73, 73.5]	(51.5, 52]
Gory; L9	1.8 → 3	(70, 70.5]	(78, 78.5]
Gory; L19	3 → 3.8	(375, 377.5]	(375, 377.5]
Gory; L21	3 → 4.2	(495, 500]	(492.5, 495]
Gory; L39	3 → 7.8	(3520, 3540]	(2660, 2680]

Every shown endpoint: native field-tail PASS.

Both scalar gates give the same displayed brackets.
F = last tested fail; P = first tested pass.

Existing feasibility
Broader helps WP at the selected sizes; it delays smaller Gory, shares the L19 bracket, then helps by L21 and at larger sizes.

Measurement gap
Threshold searches retain scalar histories, tail summaries and final fields, rather than full regional movies across the grid.

Increasing support at fixed requested peak also increases integrated cue. These comparisons sample two widths per size, rather than a complete width scan.

Existing success: persistent reversed scalar polarity near its initial magnitude after 10,000 cue-free units.

Field-tail qualification is separate; whole-field replacement is not implied. N = 201; no spatial-convergence claim.