


Wave-pinning: what changes when the same-peak cue is broader?


New full-field experiments from exact shared basals; two widths at the same pulse within each size.


Solid = original-core activity / initial amount (left axis). Dashed = native active peak inside actual cue support (right axis).














L9: w1.8: scalar-strength FAIL, field-tail PASS; w3: scalar-strength PASS, field-tail PASS

L19: w3: scalar-strength FAIL, field-tail PASS; w3.8: scalar-strength PASS, field-tail PASS


Regional activity loss is not identified molecular damage. Width also changes integrated input; native stimulus × inactive is not a calibrated optical actuator.


Same pulse within each width pair; pulses differ across the two sizes. N = 201; no spatial-convergence claim. Scalar gates and regional profiles are separate observations.


Goryachev: what changes when the same-peak cue is broader?


New full-field experiments from exact shared basals; two widths at the same pulse within each size.


Solid = original-core activity / initial amount (left axis). Dashed = native active peak inside actual cue support (right axis).














L9: w1.8: scalar-strength PASS, field-tail PASS; w3: scalar-strength FAIL, field-tail PASS

L25: w3: scalar-strength FAIL, field-tail PASS; w5: scalar-strength PASS, field-tail PASS

Regional activity loss is not identified molecular damage. Width also changes integrated input; native stimulus × inactive is not a calibrated optical actuator.

Same pulse within each width pair; pulses differ across the two sizes. N = 201; no spatial-convergence claim. Scalar gates and regional profiles are separate observations.