



Holmes · Size changes cue-on reorientation

Cue-on response for 100 native units. Rate is measured from the native RHS at each saved state; no reintegration.










Core width 8.223; cue overlap 20.4%.
Withdrawal 65.50°; after 13,000 cue-free units 60.82°.



Core width 16.538; cue overlap 100.0%.
Withdrawal 5.70°; after 13,000 cue-free units 1.78°.


Native cue: full physical support 3, prescribed peak 1; matched input-coefficient integral 1.5, not activated amount.


Cue-on time axes: linear from 0–1, logarithmic thereafter. Rates are angular activity-vector rates; they are not peak travel or cell velocity.


Weak R < 0.1 would mask heading and rate independently; no weak interval occurs here. All selected histories pass the finite full-field tail check.


Wave-pinning · Placement changes cue-on reorientation

Cue-on response for 300 native units. Rate is measured from the native RHS at each saved state; no reintegration.










Core width 12.280; cue overlap 100.0%.
Withdrawal 1.64°; after 13,000 cue-free units 1.09°.



Core width 12.280; cue overlap 0.0%.
Withdrawal 50.94°; after 13,000 cue-free units 46.97°.


Native cue: full physical support 3, prescribed peak 0.1; matched input-coefficient integral 0.15, not activated amount.


Cue-on time axes: linear from 0–1, logarithmic thereafter. Rates are angular activity-vector rates; they are not peak travel or cell velocity.


Weak R < 0.1 would mask heading and rate independently; no weak interval occurs here. All selected histories pass the finite full-field tail check.


Goryachev · Near and far cues follow different time courses

Cue-on response for 300 native units. Rate is measured from the native RHS at each saved state; no reintegration.










Core width 1.434; cue overlap 0.0%.
Withdrawal 29.59°; after 13,000 cue-free units 44.10°.



Core width 1.434; cue overlap 0.0%.
Withdrawal 33.48°; after 13,000 cue-free units -0.03°.

Native cue: full physical support 3, prescribed peak 6.4; matched input-coefficient integral 9.6, not activated amount.

Cue-on time axes: linear from 0–1, logarithmic thereafter. Rates are angular activity-vector rates; they are not peak travel or cell velocity.

Weak R < 0.1 would mask heading and rate independently; no weak interval occurs here. All selected histories pass the finite full-field tail check.