
Holmes · Every pulse: withdrawal and retained activity vectors

Orange: cue withdrawal. Black: after 13,000 cue-free units. Each arrow is the actual P; one common scale in every cell.

Background: fraction of initial target-angle error removed at the retained endpoint (colour clipped to 0–1 only). Grey: pending / weak final R.




All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.


Rightward guide is P(0)=1; circle = measured R<0.1 without a heading. ? full-field tail unavailable; ! failed finite-tail gate. No minimum-pulse inference.

Holmes · Every pulse: cue-on response and sampled coverage

Five tested cue durations × every prescribed distance × three exact qualified basals. Grey = pending or the specific quantity is unavailable.








All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.


Error-closure colours clip to 0–1; exact raw values remain in CSV. Fixed original-region loss is redistribution-compatible, not site ancestry or destruction.


Holmes · Every pulse: vector magnitude, active amount and fixed-region response

Five tested cue durations × every prescribed distance × three exact qualified basals. Grey = pending or the specific quantity is unavailable.










All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC

I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Error-closure colours clip to 0–1; exact raw values remain in CSV. Fixed original-region loss is redistribution-compatible, not site ancestry or destruction.

Holmes · Every pulse: finite cue-free endpoint organization

Five tested cue durations × every prescribed distance × three exact qualified basals. Grey = pending or the specific quantity is unavailable.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC

I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Error-closure colours clip to 0–1; exact raw values remain in CSV. Fixed original-region loss is redistribution-compatible, not site ancestry or destruction.

Holmes · Every placement: full longest-cue time courses

d₀=0.5 d₀=1.25 d₀=2.5 d₀=5 d₀=7.5 d₀=9 d₀=10 d₀=11 d₀=12 d₀=13 d₀=14

C=30 · 11/11 histories

C=35 · 11/11 histories

C=40 · 11/11 histories

Heading (degrees)

R = |P|

Total active amount / initial

Fixed original-region excess

Sampled phase secant (degrees / native unit)

Cue-on time (native units)

All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC

I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Solid/dashed/dotted = I/S/O at initial 10% edge. Time linear to 0.1 then log; rates linear near ±0.001 then log. Each rate segment spans its actual saved interval.