
Spring · Every pulse: withdrawal and retained activity vectors

Orange: cue withdrawal. Black: after 13,000 cue-free units. Each arrow is the actual P; one common scale in every cell.

Background: fraction of initial target-angle error removed at the retained endpoint (colour clipped to 0–1 only). Grey: pending / weak final R.




All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.


Rightward guide is P(0)=1; circle = measured R<0.1 without a heading. ? full-field tail unavailable; ! failed finite-tail gate. No minimum-pulse inference.


Spring · Every pulse: cue-on response and sampled coverage

Five tested cue durations × every prescribed distance × three exact qualified basals. Grey = pending or the specific quantity is unavailable.









All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.


Error-closure colours clip to 0–1; exact raw values remain in CSV. Fixed original-region loss is redistribution-compatible, not site ancestry or destruction.

Spring · Every pulse: vector magnitude, active amount and fixed-region response

Five tested cue durations × every prescribed distance × three exact qualified basals. Grey = pending or the specific quantity is unavailable.










All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.


Error-closure colours clip to 0–1; exact raw values remain in CSV. Fixed original-region loss is redistribution-compatible, not site ancestry or destruction.


Spring · Every pulse: finite cue-free endpoint organization

Five tested cue durations × every prescribed distance × three exact qualified basals. Grey = pending or the specific quantity is unavailable.









All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC

I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Error-closure colours clip to 0–1; exact raw values remain in CSV. Fixed original-region loss is redistribution-compatible, not site ancestry or destruction.

Spring · Every placement: full longest-cue time courses

d₀=0.065088 d₀=0.13018 d₀=0.26035 d₀=0.5207 d₀=0.7 d₀=0.85 d₀=0.95 d₀=1.05 d₀=1.15 d₀=1.25

C₀=2.5 · 10/10 histories

C₀=2.75 · 10/10 histories

C₀=3 · 10/10 histories

Heading (degrees)

R = |P|

Total active amount / initial

Fixed original-region excess

Sampled phase secant (degrees / native unit)

Cue-on time (native units)

All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Solid/dashed/dotted = I/S/O at initial 10% edge. Time linear to 0.1 then log; rates linear near ±0.001 then log. Each rate segment spans its actual saved interval.

Spring · Every final profile · C₀=2.5 · cue T=0.1

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=2.5 · cue T=1

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=2.5 · cue T=10

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=2.5 · cue T=100

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=2.5 · cue T=1000

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=2.75 · cue T=0.1

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=2.75 · cue T=1

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=2.75 · cue T=10

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=2.75 · cue T=100

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=2.75 · cue T=1000

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=3 · cue T=0.1

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=3 · cue T=1

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC

I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=3 · cue T=10

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.

All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=3 · cue T=100

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC


I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.

Spring · Every final profile · C₀=3 · cue T=1000

Dotted grey: initial native basal. Black: measured final active field. Blue: basal rotated by final first-moment phase; this is a shape comparison, not lineage.

Coloured support bands show the prescribed initial footprint (I/S/O at the 10% edge). Full-ring coordinate is physical arc on the initial basal, also for Spring.



All requested placements and durations retained · 645/645 conditions available · data 2026-10-19T20:51:57 UTC

I / S / O = entire initial cue footprint inside / straddling / outside the 10%-contrast region. Centre membership is recorded separately.

Endpoint is 13,000 units after withdrawal; final orientation alone does not establish stationarity. All profiles use common native-amplitude limits within this model.