Existing-law strength sweep: exploratory organization readouts


Positive input mean and fluctuation amplitude scale together; each model keeps its native law and three original sizes.

Native status observed 2026-10-19 18:55:13 UTC: running · 229/256 new outputs · 0 failures


New levels: two-seed target (exploratory). Original 0.001/0.01: twelve seeds. Whiskers show observed seed range, not confidence intervals.

solid: raw regions / noisy R / amount A dashed: qualified regions / mean-control R / coherence Q


A1 Wave-pinning: multiple regions




A2 First-moment magnitude




A3 Active amount and coherence




B1 Goryachev: multiple regions




B2 First-moment magnitude




B3 Active amount and coherence




C1 Holmes 3: multiple regions




C2 First-moment magnitude




C3 Active amount and coherence




D1 Spring ring: multiple regions



D2 First-moment magnitude



D3 Active amount and coherence



Regions: connected arcs above current minimum + half current contrast. Dashed occupancy also requires contrast ≥10% of own basal contrast.

$A = M_0/M_0(0)$; $Q = q_1/q_1(0)$; $R = AQ$ at each frame. Mean R is the mean of the paired product, not the product of separate means.

Filled circles: complete n=2; squares: n=12. Open data markers: partial n=1; open markers below occupancy panels: no noisy path available.


Full windows: chemical 0–32000 (cadence 5); Spring 0–3200 (cadence 0.5). Connecting segments guide the eye. Native units are not calibrated across models.

What multiple threshold regions look like in native profiles

C=20, f=1; first prespecified pilot path per model. Snapshot conditioning is diagnostic, not representative sampling.


Wave-pinning · native t=16040

2 threshold regions; R=0.885, A=1.922, Q=0.461



Goryachev · native t=16040

2 threshold regions; R=0.072, A=1.091, Q=0.066



WP selection: among its saved frames with >1 current half-contrast region, contrast ≥10% of basal, and R≥0.75, choose nearest the window midpoint.
Gory uses the same native time on its own independent path. The two models have separate native concentration scales; neither profile is peak-normalized.
The matched mean control receives the declared ensemble mean at every location. A low first moment or many threshold regions alone does not identify a mechanism.
Threshold arcs are morphology descriptors: they do not establish equal-strength peaks, functional competition, site lineage, or an event class.