Existing-law strength sweep: exploratory organization readouts


Positive input mean and fluctuation amplitude scale together; each model keeps its native law and three original sizes.

Native status observed 2026-10-19 18:55:13 UTC: running · 229/256 new outputs · 0 failures


New levels: two-seed target (exploratory). Original 0.001/0.01: twelve seeds. Whiskers show observed seed range, not confidence intervals.

solid: raw regions / noisy R / amount A dashed: qualified regions / mean-control R / coherence Q


A1 Wave-pinning: multiple regions




A2 First-moment magnitude




A3 Active amount and coherence




B1 Goryachev: multiple regions




B2 First-moment magnitude




B3 Active amount and coherence




C1 Holmes 3: multiple regions




C2 First-moment magnitude




C3 Active amount and coherence




D1 Spring ring: multiple regions



D2 First-moment magnitude



D3 Active amount and coherence



Regions: connected arcs above current minimum + half current contrast. Dashed occupancy also requires contrast ≥10% of own basal contrast.

$A = M_0/M_0(0)$; $Q = q_1/q_1(0)$; $R = AQ$ at each frame. Mean R is the mean of the paired product, not the product of separate means.

Filled circles: complete n=2; squares: n=12. Open data markers: partial n=1; open markers below occupancy panels: no noisy path available.

Full windows: chemical 0–32000 (cadence 5); Spring 0–3200 (cadence 0.5). Connecting segments guide the eye. Native units are not calibrated across models.