
Late-quarter organization sensitivity

Quarter4 minus quarter3: chemicals24000–32000 minus16000–24000; Spring2400–3200 minus1600–2400.

Qualified multiple-region occupancy Low-contrast occupancy

Filled squares:n=12; open circles:n=2. Bars:95% whole-seed intervals where n≥3 and direction support≥95%; unavailable directions stay absent.



These are finite-window differences, not a stationarity pass/fail test. Intervals resample independent whole seeds with quarter pairing preserved.

Quarter direction uses shorter native lag2000/200; main lag8000/800 leaves only one pair per seed in a quarter and cannot meet the reporting support gate.


Use the exported quarter-specific pair coverage and seed counts. No continuous region lifetime or migration-velocity inference is made.

Late-quarter magnitude and direction sensitivity

Quarter4 minus quarter3: chemicals24000–32000 minus16000–24000; Spring2400–3200 minus1600–2400.

Mean R Direction cosine at native lag2000/200

Filled squares:n=12; open circles:n=2. Bars:95% whole-seed intervals where n≥3 and direction support≥95%; unavailable directions stay absent.



These are finite-window differences, not a stationarity pass/fail test. Intervals resample independent whole seeds with quarter pairing preserved.

Quarter direction uses shorter native lag2000/200; main lag8000/800 leaves only one pair per seed in a quarter and cannot meet the reporting support gate.

Use the exported quarter-specific pair coverage and seed counts. No continuous region lifetime or migration-velocity inference is made.