


Local organization during a short second-half window


Largest original size; f=0.1. Fixed display window begins at half the run; white line marks the native profile shown below.

Morphology-descriptor medoid from12 independent paths per condition; fixed snapshot time is independent of field values or region counts.










A region is a connected arc above half current contrast; counts need amplitude context. Goryachev's high-input noisy AND mean-control profiles have low contrast.

Snapshots are fixed in advance within the displayed windows, not selected for extra peaks. Regions do not establish competition or site lineage; no duration classifier.

Native activity; no alignment/peak normalization. Saved fields: chemicals40, Spring4 native units; actual C(t) enters Spring M_0/M_1. Laws/clocks differ.

Size effects on activity-heading memory depend on input regime

Second-half windows: chemicals16000–32000, Spring1600–3200. First half discarded; stationarity is not assumed.



Every displayed condition:12 independent seeds. Whiskers:95% whole-seed percentile intervals (2000 paired resamples).

Direction requires both endpoint R≥0.1; usable coverage retains all available lag pairs. Multiple regions require frame contrast≥10% of basal; low contrast is below that threshold.

Positive mean and fluctuation amplitude scale together. Native laws/clocks differ. Activity heading is not cell migration; size effects do not isolate physical site motion from geometry.

Read with the native histories and full temporal/window/organization supplements. Spring sampled run-duration inference is excluded from the accepted primary scope.