


A


Drift
Same-rate jumps
Perimeter-rate jumps
C = 20
C = 60
Elapsed time / 1000
Relative arc position

B


Drift, α = 2
Same-rate jumps, α = 0
Perimeter-rate jumps, α = −1
C = 20
C = 60
Heading correlation
Lag / 1000

C


Wave-pinning, f = 0.005
Holmes, f = 0.005
Spring, f = 0.005
Goryachev, f = 0.005
Wave-pinning, f = 0.3
Holmes, f = 0.3
Spring, f = 0.3
Goryachev, f = 0.3
C = 20
C = 40
C = 60
C = 1
C = 2
C = 3
Heading correlation
Lag / τ

D

Wave-pinning, C = 20, f = 0.005
Wave-pinning, C = 40, f = 0.005
Wave-pinning, C = 60, f = 0.005
Wave-pinning, C = 20, f = 0.3
Wave-pinning, C = 40, f = 0.3
Wave-pinning, C = 60, f = 0.3
Elapsed time / 1000
Relative arc position
Activity u

E



Wave-pinning
Holmes
Spring
Goryachev
C = 20
C = 40
C = 60
C = 1
C = 2
C = 3
Local size exponent α
Mean input fraction f

Figure 4. Larger rings retain coherent heading longer at weak input

At weak input, enlargement increases native activity-heading persistence, with an exponent near the geometric drift baseline in Holmes and Spring and above it in Wave-pinning. At strong input, the readout is orientation of whole-ring activity imbalance; Spring curves nearly coincide and the size advantage is reduced.

A, Synthetic activity kymographs for a single drifting peak, same-rate random relocations and relocations whose rate scales with circumference, at C20 and C60. Position is relative arc length; grayscale denotes activity from 0 (white) to 1 (black). Time increases upward. These pedagogical baselines have no biochemical kinetics.

B, Heading correlation in the same three baselines, from 1000 seeds; solid and dashed lines indicate C20 and C60, dotted lines the analytic curves. A fixed physical drift diffusivity gives k proportional to C^{-2} ($\alpha = 2$); size-independent jumps give $\alpha = 0$, and jumps per unit perimeter give $\alpha = -1$. The same-rate curves coincide.

C, Native-model heading correlation at common weak f = 0.005 and strong f = 0.3, for C20, C40 and C60 (Spring 1, 2 and 3). Size is a light-to-dark grayscale ramp within each facet. Shading is the 95% confidence interval from 2000 seed-bootstrap resamples of the complete retained native histories, recomputed on a dense early-lag grid. Weak-input facets show 0–320τ; strong-input facets show 0–12τ, Goryachev 0–80τ, to resolve their different correlation scales. $\tau = 50$ native time units, Spring 5. Heading correlation is the mean cosine of the change in arg(P), where P is the first Fourier activity moment divided by total activity. These estimates require $|P| \geq 0.1$ at both lag endpoints; their coverage is in condition_lag_metrics.csv. At weak input, Goryachev C40 and C60 remain near correlation 1 throughout the window, so their ordering is unresolved. At strong input, Spring curves nearly coincide. In patch fields, heading denotes the orientation of whole-ring activity imbalance, rather than persistence of an established polarity domain. Figure5D shows the ungated distributions.

D, Native Wave-pinning examples at f = 0.005 and 0.3 and C20, C40 and C60, selected as replicate 0 without a seed search. Every panel shows the entire retained post-burn-in interval 16000–32000, on relative arc position. Activity scales are common across sizes at each input; the color bars state the separate weak- and strong-input ranges. Weak input mainly shifts a coherent domain; strong input produces a changing patch field. The kymographs are illustrative, not an event classification.

E, Local α, the negative fitted slope of log heading-decorrelation rate against log circumference in adjacent-size neighborhoods, against mean input f. Lines are size neighborhoods centered on C20, C30, C40, C50 and C60 (Spring 1, 1.5, 2, 2.5 and 3); light to dark follows size. Vertical bars are seed-bootstrap 95% CIs. Missing estimates interrupt curves. Horizontal guides are $\alpha = 2$, 0 and −1. Native f multiplies reference input 0.1, 1, 4 and 6.4 in Wave-pinning, Holmes, Spring and Goryachev respectively; doses and clocks are not calibrated across models.

All simulations reuse the complete, frozen round6 compact campaign (96–384 independent seeds per condition). Local kinetics, basal preparations and solver outputs are unchanged. α and activity-heading persistence are model observables, distinct from migration persistence or fitted OU velocity-memory parameters.