




Figure 6 | Migration summaries and conditional OU dynamics















Points sample digitized published curves; original experimental lag grid unknown.

B slopes describe curve shape, not independent measurements or OU memory times. C tracks are simulations. D/F show conditional objective/prior sensitivity.

No new fit or simulation; A and C–F retain their original numerical evidence. See the enlarged B detail for readable curve-shape comparison.

Figure 6B | Digitized population MSD and curve shape



Points sample digitized published curves;
original experimental lag grid unknown.

Lower points: newly estimated window slopes. Correlated raster samples; no biological error bars.

Figure 6. Published migration summaries support a persistence hypothesis with conditional OU interpretation

A, Published/digitized enlarged or polyploid-group to reference-group ratios for nine migration summaries, alongside primary Ornstein–Uhlenbeck (OU) fit predictions. Xuan contributes sampled mean speed and population mean-squared displacement (MSD) at the last shared lag, approximately 10.95 h; Hosny summaries are group medians. Mallin central-bar statistic is not specified by the source and retains the historical mean approximation. These distinct source summaries are not interchangeable measures of persistence.

B, Upper: twelve chosen display samples from each digitized Xuan population MSD curve (small points) and the unchanged fitted analytical OU expectation (solid lines), on logarithmic lag and MSD axes. These are samples of the published raster curves, not recovered experimental lag positions or independent cell observations; the original experimental lag grid is unknown. The open circle identifies the crossover of the full digitized traces at approximately 2.32 h, computed by linear interpolation in native lag and MSD. The open diamond identifies the different fitted OU crossover at approximately 1.11 h. Lower: descriptive slopes of log(MSD) versus log(lag) fitted in six equal log-lag windows of the full digitized curves (points), and matching-window moving OU slopes (solid lines). Each window spans approximately 0.304 decades; both observed and model slopes use the same full window support. Observed raster columns remain correlated and give no biological sample size or confidence interval. These slopes are newly calculated curve-shape summaries, not author-reported local exponents or fitted OU memory times. A constant power-law exponent would appear horizontal; reference values 1 and 2 indicate diffusive and straight-line scaling. The model does not reproduce the complete observed curvature or the observed crossover. The analytical fit, all fitting targets and numerical inputs to A and C–F are unchanged.

C, Representative Xuan primary-fit model-generated tracks, with sampled speed, total path length and directness (net displacement divided by path length). Both groups use identical spatial scales. The tracks are simulations, not published individual-cell trajectories; a slower sampled trajectory can nevertheless have greater endpoint directness. Dashed chords illustrate the contribution of finite-time displacements.

D, With reference-group parameters fixed, vary enlarged-group persistence and velocity-noise ratios. Separate surfaces show change in the descriptive data-misfit score and in the total regularized objective; the selected fit and equal-group reference are marked. The heuristic score is not a calibrated likelihood, and the surfaces are not confidence regions.

E, Primary paired persistence and velocity-noise estimates for Xuan 2018, Mallin 2023, Jemaà 2017, Jemaà 2023 and the Hosny hypoxic baseline. F, Enlarged/reference persistence folds under the primary 10-minute prior, 5- and 20-minute alternatives and stronger/weaker pooling, with the pooled geometric mean. Priors and pooling define conditional sensitivity scenarios, not uncertainty intervals. The Jemaà comparisons constrain speed alone, leaving persistence/noise tradeoffs dependent on hierarchy and priors.

The fitted population contrasts span stable tetraploids, giant polyploids and treatment-induced polyaneuploid cancer cells (PACCs). The inherited WGD+/WGD− labels in the artwork are comparison shorthand and do not establish matched WGD-only interventions in every study.

Figure 6. Published migration summaries support a persistence hypothesis with conditional OU interpretation · continued

Hosny observations derive from the recovered preprint; equivalence to the journal version remains unresolved.