Figure 3 round 2 · decomposition and censoring

Separation slows the resolution of coexisting regions at short range; at matched separation, cell size dominates both driven takeover and resolution. Extra separation adds at most 1.15-fold to driven handover. In resolved cue-free comparisons it adds about 1.01-1.07-fold in Wave-pinning and Goryachev; Spring middle size is the supported exception, about 1.75-fold. Censored comparisons do not establish a numerical extra-separation effect.

For each model and assay, the matched gap is the largest gap within both measured size curves. In this dataset it is the small-size antipodal-placement endpoint. In B, the coordinate is the measured gap at withdrawal, not the initial cue gap. The endpoint called antipode is the trajectory seeded by the initial antipodal cue: its withdrawal region centers can shift.

A1 uses native continuous complete-region handover roots. B uses saved-sample lower and upper time brackets. Linear interpolation is restricted to adjacent measured eligible placements. Interval endpoints are interpolated separately; any censored upper endpoint stays infinite. Missing or topology-ineligible values are never converted into bounds.

A ratio with a censored numerator and resolved denominator has a lower bound. A resolved numerator and censored denominator gives an upper bound. Two censored times give no finite one-sided ratio bound; the result is marked indeterminate. In particular, dividing two horizon caps is invalid. Finite bars use the geometric center of the resulting interval, with the full interval recorded in CSV.

The exact product identity uses the same interpolated matched-gap time in both terms. For finite B intervals the representative values obey the identity; multiplying independently propagated marginal interval limits would ignore this shared denominator and numerator. Total intervals are computed directly from endpoint times.

All 9 author-approved Holmes ratio-1 extensions (5 small, 4 middle) completed to 80,000 from exact archived 20,000 endpoints. Six resolve and three remain censored. Other Holmes conditions retain their original 20,000 horizon; different lower bounds reflect observation duration, not a decline in resolution time. No cue, kinetic parameter, mesh or resolution definition changed.

The following pages retain BOTH superseded main pages, including the complete translation page, then the entire seven-page existing supplement. All four native-profile appendix volumes remain linked in the existing reader; no model inputs, histories, or prior supplements are replaced.

Approved Holmes cue-free extensions: every trajectory continued to 80,000

Circumference	Withdrawal gap	Resolution time	Final native state	Field steady
30	4.3786	29,420–29,440	single region incumbent associated	True
30	5.3610	32,160–32,180	single region incumbent associated	True
30	6.3425	32,280–32,300	single region incumbent associated	True
30	7.3241	32,320–32,340	single region incumbent associated	True
30	7.6088	32,320–32,340	single region incumbent associated	True
37.8	4.5702	34,380–34,400	single region incumbent associated	True
37.8	5.5498	≥80,000	two resolved spatially associated regions	False
37.8	6.5270	≥80,000	two resolved spatially associated regions	False
37.8	7.5037	≥80,000	two resolved spatially associated regions	True

Full native fields and complete-region readouts are retained. Fate and full-field stationarity are independent.
Six observed resolutions; three one-sided lower bounds. Original campaigns are unchanged.
Mixed 20,000/80,000 bounds on the main curves identify observation duration, not point estimates.

All decomposition values · time ratios with interval bounds

Model	Assay	Size / small	Matched gap	Size term	Extra separation	Total
Wave-pinning	A1	37.8 / 30	7.719	2.14	1.01	2.17
Wave-pinning	A1	47.6 / 30	7.719	5.59	1.05	5.86
Goryachev	A1	37.8 / 30	12.05	1.84	1.04	1.91
Goryachev	A1	47.6 / 30	12.05	3.5	1.15	4.02
Holmes	A1	37.8 / 30	6.29	3.59	1	3.6
Holmes	A1	47.6 / 30	6.29	23.9	1.01	24.2
Spring	A1	3.15 / 2.5	0.4296	2.44	1	2.44
Spring	A1	3.97 / 2.5	0.4296	8.45	1.01	8.52
Wave-pinning	B	37.8 / 30	8.378	5.67–5.77	1.02–1.03	5.81–5.92
Wave-pinning	B	47.6 / 30	8.378	≥15.9	indeterminate (both censored)	≥15.9
Goryachev	B	37.8 / 30	11.87	1.15–1.16	1.01–1.02	1.16–1.18
Goryachev	B	47.6 / 30	11.87	1.44–1.47	1.06–1.09	1.54–1.58
Holmes	B	37.8 / 30	7.609	≥2.37	indeterminate (both censored)	≥0.618
Holmes	B	47.6 / 30	7.609	≥0.618	indeterminate (both censored)	≥0.618
Spring	B	3.15 / 2.5	0.5456	4.2–4.38	1.73–1.77	7.31–7.71
Spring	B	3.97 / 2.5	0.5456	≥6.87	indeterminate (both censored)	≥13.8

Intervals reflect saved-time brackets, not biological uncertainty. ≥ is a one-sided lower bound.
All interval endpoints, interpolation weights and source slots appear in the companion CSVs.