




S1 Wave-pinning • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Wave-pinning • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Wave-pinning • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Wave-pinning • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Wave-pinning • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Wave-pinning • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Wave-pinning • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Wave-pinning • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Wave-pinning • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Goryachev • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Goryachev • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Goryachev • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Goryachev • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Goryachev • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Goryachev • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Goryachev • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Goryachev • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Goryachev • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 deBelly • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 deBelly • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 deBelly • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 deBelly • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 deBelly • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 deBelly • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 deBelly • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 deBelly • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 deBelly • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Holmes 3 • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Holmes 3 • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Holmes 3 • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Holmes 3 • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Holmes 3 • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Holmes 3 • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Holmes 3 • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Holmes 3 • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Holmes 3 • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Spring • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Spring • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Spring • commanded area 0.5× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Spring • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Spring • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Spring • commanded area 1× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Spring • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Spring • commanded area 2× • native kinetics


















Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.


S1 Spring • commanded area 2× • native kinetics














Triangles are finite lower bounds. Regional excess uses the instantaneous spatial minimum; ROI conventions can overlap the initial incumbent and do not define the primary disjoint-half D. Every field-tail flag remains in the source table.