

# S9 · E0 provisional  Region births per native time per physical perimeter





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_001_births_per_time_per_perimeter.csv

# S9 · E0 provisional  Region births per native time per physical perimeter · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_002_births_per_time_per_perimeter_sensitivity.csv

# S9 · E0 provisional  Fraction of trajectories with an unscorable takeover track





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_003_ever_unscorable_takeover_fraction.csv

# S9 · E0 provisional  Fraction of trajectories with an unscorable takeover track · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_004_ever_unscorable_takeover_fraction_sensitivity.csv

# S9 · E0 provisional  Time fraction below the contrast gate





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_005_low_contrast_time_fraction.csv

# S9 · E0 provisional  Time fraction below the contrast gate · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_006_low_contrast_time_fraction_sensitivity.csv

# S9 · E0 provisional  Mean absolute dominant-heading turn over 1 observation frames





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_007_mean_absolute_turn_1_frames.csv

# S9 · E0 provisional  Mean absolute dominant-heading turn over 1 observation frames · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_008_mean_absolute_turn_1_frames_sensitivity.csv

# S9 · E0 provisional  Mean absolute dominant-heading turn over 2 observation frames





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_009_mean_absolute_turn_2_frames.csv

# S9 · E0 provisional  Mean absolute dominant-heading turn over 2 observation frames · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_010_mean_absolute_turn_2_frames_sensitivity.csv

# S9 · E0 provisional  Mean absolute dominant-heading turn over 4 observation frames





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_011_mean_absolute_turn_4_frames.csv

# S9 · E0 provisional  Mean absolute dominant-heading turn over 4 observation frames · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_012_mean_absolute_turn_4_frames_sensitivity.csv

# S9 · E0 provisional  Mean absolute dominant-heading turn over 8 observation frames





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_013_mean_absolute_turn_8_frames.csv

# S9 · E0 provisional  Mean absolute dominant-heading turn over 8 observation frames · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_014_mean_absolute_turn_8_frames_sensitivity.csv

# S9 · E0 provisional  Mean newborn angle from the previous dominant region





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_015_mean_birth_angle.csv

# S9 · E0 provisional  Mean newborn angle from the previous dominant region · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_016_mean_birth_angle_sensitivity.csv

# S9 · E0 provisional  Mean newborn physical arc distance





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_017_mean_birth_physical_distance.csv

# S9 · E0 provisional  Mean newborn physical arc distance · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_018_mean_birth_physical_distance_sensitivity.csv

# S9 · E0 provisional  Mean physical region width





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_019_mean_region_width.csv

# S9 · E0 provisional  Mean physical region width · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_020_mean_region_width_sensitivity.csv

# S9 · E0 provisional  Mean region width / actual circumference





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_021_mean_region_width_fraction.csv

# S9 · E0 provisional  Mean region width / actual circumference · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_022_mean_region_width_fraction_sensitivity.csv

# S9 · E0 provisional  Time fraction with multiple qualified regions





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_023_multiple_front_fraction.csv

# S9 · E0 provisional  Time fraction with multiple qualified regions · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_024_multiple_front_fraction_sensitivity.csv

# S9 · E0 provisional  Observed region births per trajectory





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_025_observed_births.csv

# S9 · E0 provisional  Observed region births per trajectory · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_026_observed_births_sensitivity.csv

# S9 · E0 provisional  Literal pairwise takeover sensitivity per trajectory





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Literal pairwise events are a separate sensitivity and can occur while a third region remains dominant. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_027_observed_pairwise_takeovers.csv

# S9 · E0 provisional  Literal pairwise takeover sensitivity per trajectory · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_028_observed_pairwise_takeovers_sensitivity.csv

# S9 · E0 provisional  Literal pairwise takeover sensitivity per birth





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Literal pairwise events are a separate sensitivity and can occur while a third region remains dominant. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_029_observed_pairwise_takeovers_per_birth.csv

# S9 · E0 provisional  Literal pairwise takeover sensitivity per birth · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_030_observed_pairwise_takeovers_per_birth_sensitivity.csv

# S9 · E0 provisional  Observed globally dominant newborn takeovers per trajectory





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_031_observed_takeovers.csv

# S9 · E0 provisional  Observed globally dominant newborn takeovers per trajectory · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_032_observed_takeovers_sensitivity.csv

# S9 · E0 provisional  Globally dominant newborn takeovers per birth





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_033_observed_takeovers_per_birth.csv

# S9 · E0 provisional  Globally dominant newborn takeovers per birth · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_034_observed_takeovers_per_birth_sensitivity.csv

# S9 · E0 provisional  Fraction of births opposite the previous dominant region





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_035_opposite_birth_fraction.csv

# S9 · E0 provisional  Fraction of births opposite the previous dominant region · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_036_opposite_birth_fraction_sensitivity.csv

# S9 · E0 provisional  Mean simultaneous qualified-region count





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_037_region_count_mean.csv

# S9 · E0 provisional  Mean simultaneous qualified-region count · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_038_region_count_mean_sensitivity.csv

# S9 · E0 provisional  Lower-censoring restricted mean lifetime





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_039_restricted_mean_lifetime_lower.csv

# S9 · E0 provisional  Lower-censoring restricted mean lifetime · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_040_restricted_mean_lifetime_lower_sensitivity.csv

# S9 · E0 provisional  Observed-frame restricted mean lifetime





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_041_restricted_mean_lifetime_observed.csv

# S9 · E0 provisional  Observed-frame restricted mean lifetime · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_042_restricted_mean_lifetime_observed_sensitivity.csv

# S9 · E0 provisional  Upper-censoring restricted mean lifetime





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_043_restricted_mean_lifetime_upper.csv

# S9 · E0 provisional  Upper-censoring restricted mean lifetime · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_044_restricted_mean_lifetime_upper_sensitivity.csv

# S9 · E0 provisional  Time fraction with one qualified region





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_045_single_front_fraction.csv

# S9 · E0 provisional  Time fraction with one qualified region · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_046_single_front_fraction_sensitivity.csv

# S9 · E0 provisional  Tracks whose takeover cannot be scored





E0 only; later fine-profile experiments pending. Primary optics: threshold 0.5; fixed physical Gaussian σ=1.5 (Spring 0.15); common cadence 80 (Spring 8). Grey points are forced conditions. Dashed mean controls and dotted zero-input references remain shown. CIs resample whole seeds; open triangles preserve identified lifetime bounds and × marks no identified estimate/bound. Primary takeover requires the newborn to be globally dominant among qualified regions and have pairwise D>0.5 against the previous dominant for two observed frames. Complete native peak basins define amounts; lifetimes remain cadence/censoring limited. Units are model-native; absolute cross-model comparison is not used.
CSV: figure4_S9_E0_047_unscorable_takeover_tracks.csv

# S9 · E0 provisional  Tracks whose takeover cannot be scored · sensitivity





E0 only; later fine-profile experiments pending. Size colours are fixed. One-factor contrasts: threshold 0.3/0.5/0.7 uses solid/dashed/dotted; native-profile downsampling ×1/×2/×4/×8 uses solid/dashed/dotted/dash-dot; optics raw/smoothed uses solid/dashed. Other factors use the common primary. The full 24-way factorial, controls, CIs and unavailable estimands remain in this page's CSV; no combined-frame bootstrap is used.
CSV: figure4_S9_E0_048_unscorable_takeover_tracks_sensitivity.csv

# S9 · E0 provisional  birth angle from previous dominant





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_049_birth_angle_from_previous_dominant_None_0.csv

# S9 · E0 provisional  birth angle from previous dominant





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_050_birth_angle_from_previous_dominant_None_3.csv

# S9 · E0 provisional  birth angle from previous dominant





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_051_birth_angle_from_previous_dominant_None_6.csv

# S9 · E0 provisional  birth angle from previous dominant





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_052_birth_angle_from_previous_dominant_None_9.csv

# S9 · E0 provisional  dominant heading turn · 1 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_053_dominant_heading_turn_1.0_0.csv

# S9 · E0 provisional  dominant heading turn · 1 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_054_dominant_heading_turn_1.0_3.csv

# S9 · E0 provisional  dominant heading turn · 1 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_055_dominant_heading_turn_1.0_6.csv

# S9 · E0 provisional  dominant heading turn · 1 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_056_dominant_heading_turn_1.0_9.csv

# S9 · E0 provisional  dominant heading turn · 2 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_057_dominant_heading_turn_2.0_0.csv

# S9 · E0 provisional  dominant heading turn · 2 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_058_dominant_heading_turn_2.0_3.csv

# S9 · E0 provisional  dominant heading turn · 2 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_059_dominant_heading_turn_2.0_6.csv

# S9 · E0 provisional  dominant heading turn · 2 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_060_dominant_heading_turn_2.0_9.csv

# S9 · E0 provisional  dominant heading turn · 4 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_061_dominant_heading_turn_4.0_0.csv

# S9 · E0 provisional  dominant heading turn · 4 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_062_dominant_heading_turn_4.0_3.csv

# S9 · E0 provisional  dominant heading turn · 4 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_063_dominant_heading_turn_4.0_6.csv

# S9 · E0 provisional  dominant heading turn · 4 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_064_dominant_heading_turn_4.0_9.csv

# S9 · E0 provisional  dominant heading turn · 8 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_065_dominant_heading_turn_8.0_0.csv

# S9 · E0 provisional  dominant heading turn · 8 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_066_dominant_heading_turn_8.0_3.csv

# S9 · E0 provisional  dominant heading turn · 8 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_067_dominant_heading_turn_8.0_6.csv

# S9 · E0 provisional  dominant heading turn · 8 observation frames





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_068_dominant_heading_turn_8.0_9.csv

# S9 · E0 provisional  simultaneous region count





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_069_simultaneous_region_count_None_0.csv

# S9 · E0 provisional  simultaneous region count





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_070_simultaneous_region_count_None_3.csv

# S9 · E0 provisional  simultaneous region count





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_071_simultaneous_region_count_None_6.csv

# S9 · E0 provisional  simultaneous region count





E0 only; later fine-profile experiments pending. Imaging primary only; size colours match S1 and forced conditions are grey. Solid/dashed/dotted also distinguish small/intermediate/large sizes. Histogram bin centres are displayed; exact bins, finite-seed uncertainty, observed totals and frame intervals remain in CSV. Native coarsely saved profiles limit birth/turn observability. Missing densities remain missing, not zero.
CSV: figure4_S9_E0_072_simultaneous_region_count_None_9.csv


# S9 · E0 provisional  Incident-region lifetime survival · lower observation clock




E0 only; later fine-profile experiments pending. Kaplan–Meier curves use incident tracks; preexisting left-censored regions are excluded. End-of-window and unresolved states retain right censoring. Lower/observed/upper clocks preserve cadence-limited times; symlog lifetime axes are linear through one observation cadence. Solid/dashed/dotted distinguish small/intermediate/large; forced conditions are grey. Whole-seed CIs and at-risk/censoring counts remain in CSV. Unidentifiable tails are not extrapolated.
CSV: figure4_S9_E0_073_lifetime_lower_0.csv





E0 only; later fine-profile experiments pending. Kaplan–Meier curves use incident tracks; preexisting left-censored regions are excluded. End-of-window and unresolved states retain right censoring. Lower/observed/upper clocks preserve cadence-limited times; symlog lifetime axes are linear through one observation cadence. Solid/dashed/dotted distinguish small/intermediate/large; forced conditions are grey. Whole-seed CIs and at-risk/censoring counts remain in CSV. Unidentifiable tails are not extrapolated.
CSV: figure4_S9_E0_074_lifetime_lower_3.csv





E0 only; later fine-profile experiments pending. Kaplan–Meier curves use incident tracks; preexisting left-censored regions are excluded. End-of-window and unresolved states retain right censoring. Lower/observed/upper clocks preserve cadence-limited times; symlog lifetime axes are linear through one observation cadence. Solid/dashed/dotted distinguish small/intermediate/large; forced conditions are grey. Whole-seed CIs and at-risk/censoring counts remain in CSV. Unidentifiable tails are not extrapolated.
CSV: figure4_S9_E0_075_lifetime_lower_6.csv

# S9 · E0 provisional  Incident-region lifetime survival · lower observation clock





E0 only; later fine-profile experiments pending. Kaplan–Meier curves use incident tracks; preexisting left-censored regions are excluded. End-of-window and unresolved states retain right censoring. Lower/observed/upper clocks preserve cadence-limited times; symlog lifetime axes are linear through one observation cadence. Solid/dashed/dotted distinguish small/intermediate/large; forced conditions are grey. Whole-seed CIs and at-risk/censoring counts remain in CSV. Unidentifiable tails are not extrapolated.
CSV: figure4_S9_E0_076_lifetime_lower_9.csv

# S9 · E0 provisional  Incident-region lifetime survival · observed observation clock





E0 only; later fine-profile experiments pending. Kaplan–Meier curves use incident tracks; preexisting left-censored regions are excluded. End-of-window and unresolved states retain right censoring. Lower/observed/upper clocks preserve cadence-limited times; symlog lifetime axes are linear through one observation cadence. Solid/dashed/dotted distinguish small/intermediate/large; forced conditions are grey. Whole-seed CIs and at-risk/censoring counts remain in CSV. Unidentifiable tails are not extrapolated.
CSV: figure4_S9_E0_077_lifetime_observed_0.csv

# S9 · E0 provisional  Incident-region lifetime survival · observed observation clock





E0 only; later fine-profile experiments pending. Kaplan–Meier curves use incident tracks; preexisting left-censored regions are excluded. End-of-window and unresolved states retain right censoring. Lower/observed/upper clocks preserve cadence-limited times; symlog lifetime axes are linear through one observation cadence. Solid/dashed/dotted distinguish small/intermediate/large; forced conditions are grey. Whole-seed CIs and at-risk/censoring counts remain in CSV. Unidentifiable tails are not extrapolated.
CSV: figure4_S9_E0_078_lifetime_observed_3.csv





E0 only; later fine-profile experiments pending. Kaplan–Meier curves use incident tracks; preexisting left-censored regions are excluded. End-of-window and unresolved states retain right censoring. Lower/observed/upper clocks preserve cadence-limited times; symlog lifetime axes are linear through one observation cadence. Solid/dashed/dotted distinguish small/intermediate/large; forced conditions are grey. Whole-seed CIs and at-risk/censoring counts remain in CSV. Unidentifiable tails are not extrapolated.
CSV: figure4_S9_E0_079_lifetime_observed_6.csv

# S9 · E0 provisional  Incident-region lifetime survival · observed observation clock





E0 only; later fine-profile experiments pending. Kaplan–Meier curves use incident tracks; preexisting left-censored regions are excluded. End-of-window and unresolved states retain right censoring. Lower/observed/upper clocks preserve cadence-limited times; symlog lifetime axes are linear through one observation cadence. Solid/dashed/dotted distinguish small/intermediate/large; forced conditions are grey. Whole-seed CIs and at-risk/censoring counts remain in CSV. Unidentifiable tails are not extrapolated.
CSV: figure4_S9_E0_080_lifetime_observed_9.csv


# S9 · E0 provisional  Incident-region lifetime survival · upper observation clock




E0 only; later fine-profile experiments pending. Kaplan–Meier curves use incident tracks; preexisting left-censored regions are excluded. End-of-window and unresolved states retain right censoring. Lower/observed/upper clocks preserve cadence-limited times; symlog lifetime axes are linear through one observation cadence. Solid/dashed/dotted distinguish small/intermediate/large; forced conditions are grey. Whole-seed CIs and at-risk/censoring counts remain in CSV. Unidentifiable tails are not extrapolated.
CSV: figure4_S9_E0_081_lifetime_upper_0.csv





E0 only; later fine-profile experiments pending. Kaplan–Meier curves use incident tracks; preexisting left-censored regions are excluded. End-of-window and unresolved states retain right censoring. Lower/observed/upper clocks preserve cadence-limited times; symlog lifetime axes are linear through one observation cadence. Solid/dashed/dotted distinguish small/intermediate/large; forced conditions are grey. Whole-seed CIs and at-risk/censoring counts remain in CSV. Unidentifiable tails are not extrapolated.
CSV: figure4_S9_E0_082_lifetime_upper_3.csv




E0 only; later fine-profile experiments pending. Kaplan–Meier curves use incident tracks; preexisting left-censored regions are excluded. End-of-window and unresolved states retain right censoring. Lower/observed/upper clocks preserve cadence-limited times; symlog lifetime axes are linear through one observation cadence. Solid/dashed/dotted distinguish small/intermediate/large; forced conditions are grey. Whole-seed CIs and at-risk/censoring counts remain in CSV. Unidentifiable tails are not extrapolated.
CSV: figure4_S9_E0_083_lifetime_upper_6.csv

# S9 · E0 provisional  Incident-region lifetime survival · upper observation clock




E0 only; later fine-profile experiments pending. Kaplan–Meier curves use incident tracks; preexisting left-censored regions are excluded. End-of-window and unresolved states retain right censoring. Lower/observed/upper clocks preserve cadence-limited times; symlog lifetime axes are linear through one observation cadence. Solid/dashed/dotted distinguish small/intermediate/large; forced conditions are grey. Whole-seed CIs and at-risk/censoring counts remain in CSV. Unidentifiable tails are not extrapolated.
CSV: figure4_S9_E0_084_lifetime_upper_9.csv